

Determination of the chemical compositions of *Viscum album* ssp. *album* and soft core hosts

Tamer Üstüner^{1*} and Sema Düzenli²

¹Department of Plant Protection, Faculty of Agriculture, University of Kahramanmaraş Sutcu Imam, Avsar Campus, Kahramanmaraş-Türkiye

²Department of Biology, Faculty of science literature, University of Cukurova, Adana-Türkiye

(Received July 26, 2017; Revised October 31, 2017; Accepted November 01, 2017)

Abstract: In this study, leaf proportional water content, chlorophyll a and b, total chlorophyll, carotenoid, proline, total amino acid, ascorbic acid, glucose, fructose and sulphhydryl group (SH) were determined in both *Viscum album* ssp. *album* and its hosts. Physiological alterations in soft-core fruits such as Ankara pear, Braeburn apple and hawthorn when infected with *Viscum album* ssp. *album* were investigated. While proportional water content of *V. a.* ssp. *album* was found higher than all fruit trees with or without *V. album* (control), its chlorophyll a content was lower than only control and two hosts (Braeburn apple and Ankara pear). Chlorophyll b content, on the other hand, was similar in all experimental groups for Ankara pear and hawthorn whereas, in Braeburn apple, it was significantly higher in control. As for the total chlorophyll and carotenoid contents of *V. a.* ssp. *album*, in Braeburn and Ankara pear, they were found less than the controls and hosts whilst in hawthorn, the values were variable. Although proline content of *V. a.* ssp. *album* was higher compared to their controls and hosts, total amino acid, reduced and total ascorbic acid, glucose, fructose and sulphhydryl acid in *V. a.* ssp. *album* were determined to be lower than all uninfected fruits (control).

Keywords: *V. album* sp *album*; leaf proportional water; chlorophyll a-b; total chlorophyll; carotenoid.

© 2017 ACG Publications. All rights reserved.

1. Introduction

The semi parasitic plant *V. a.* ssp. *album* causes significant loss of yield and quality in fruit production in many countries such as Turkey. *Viscum* genus of Loranthaceae family in Santalales order includes three sub species living as semi parasite on host plants. The sub species, *V. album* L. ssp. *album*, *V. album* L. ssp. *abietis* (Wiesb.) Abromeit and *V. album* L. ssp. *austriacum* (Wiesb.) Vollmann, belonging to *Viscum album* (mistletoe), were reported by Miller [1].

V. album is a mostly globose perennial evergreen shrub with persistent haustoria in the host. This semi parasite is able to conduct the photosynthesis since it contains chlorophyll a and b in its leaves. It takes water and nutrients from the host and produces its own metabolites whenever it has access to light and carbon dioxide. *Viscum* species using xylem water and dissolved organic substances from their hosts weakens and eventually kills them. *V. album* L. subsp. *album*, white berry mistletoe (Loranthaceae), is one of the most important biological stress sources for host plants [2-11]. This semi parasite plant feeds on water and nutrients obtained from its host via its haustorium [12].

Subspecies of *Viscum album* in Turkey are often semi parasite plants living on various fruit trees such as wild pear, hawthorn, pear, quince, apple and apricot. *V. a.* ssp. *album* was reported to be

* Corresponding author: E-mail: TamerUstuner@ksu.edu.tr; Phone: 90(344) 3002060, Fax: 90(344) 3002002

the most commonly observed on such as fruit trees; Ankara pear, Braeburn apple, almond, apricot, whitehorn, hawthorn, plum and on such as park trees; populus and salix in Nigde, Turkey. The severity of the disease caused by *V. album* ssp. *album* was determined as the highest on Almond (*Amygdalus* spp.) 48.54%, Apricot (*Prunus armenica* L.) 34.98 % and Pear (*Pyrus communis* L.) 28.64%. [13-15]. Glucose, fructose and sucrose contents of mistletoe on populus have been shown to be higher in spring than autumn. Xylem of populus contained highest carbohydrate concentrations in spring, and much lower concentrations in the autumn and summer [16]. Moreover, Ziegler et al. [17] measured faster rates of stem flow and transpiration in the mistletoe compared with its woody hosts. The study of the host and mistletoe ecophysiology of *Tapinanthus bangwensis* on the two hosts, *Citrus sinensis* and *Irvingia gabonensis* shows that mistletoe thrives on its hosts on the strength of the available water, mineral nutrients, sugars (photosynthates) and the effective leaf chlorophyll content [18]. According to a study carried out with *Dendrophthoe falcata* on mango, chlorophyll pigments showed a decrease possibly the reduction in availability of water and minerals. The total protein, phenols, flavonoids also showed a defense related response in the plant but, the chlorophyll content showed a decrease in the infected plant [19].

The aim of this study is to determine the leaf proportional water content, chlorophyll a-b, total chlorophyll, carotenoid, proline, total amino acid, ascorbic acid, glucose, fructose and sulphhydryl group (SH) of *V. a. ssp. album* on Ankara pear, Braeburn apple and hawthorn grown in Nigde, Turkey.

2. Materials and Methods

Plant materials used in the study were collected from Nigde province of Turkey. These are the leaf of Braeburn apple, Ankara pear, hawthorn and *V. a. ssp. album* between the years of 2014 and 2015.

Based on the meteorological data of Nigde, annual mean temperature was 18.22 °C, relative humidity 47.99 g/m³, sunning duration 7.14 hours/day, mean wind speed 2.64m/s, mean precipitation 0.866 mm and surface soil temperature 4.44 °C in these years. The soil of the study area was arid, sandy, slightly alkaline and total salt was low. The pH of the soil was 7.4, the lime ratio was 7.2, phosphorous content was 3.1 mg/kg, the amount of the organic matter was 0.32% and the potassium content was 48 mg/kg.

Proportional water content, chlorophyll and carotenoid were calculated as described by Sairam et al. [20] and Arnon [21]. In addition, other parameters were estimated by using the method defined by Lichtenthaler and Wellburn [22]. Analysis of SH-group was made by using 5-5 dithiobis (2-nitrobenzoic acid) (DTNB) solution at the 5% meta- phosphoric acid extraction. Total and reduced ascorbic acid, amino acid and proline were analyzed as previously reported [23-25].

2.1. Statistical analysis

In order to determine whether there is a statistically significant difference among the obtained results for each parameter of *V. a. ssp. album* and its hosts, variance analyses were carried out using SPSS 20 software package. Differences between means were tested by Duncan test and ANOVA. The values with a p value lower than 0.05 ($p \leq 0.05$) were considered significantly different.

3. Results and discussion

3.1. Leaf proportional water content

In three fruits (Braeburn apple, Ankara pear and hawthorn), leaf proportional water content of *V. a. ssp. album* was determined to be statistically different from control and host plants. Leaf proportional water content of *V. a. ssp. album* was about 3-4 times higher than the control and the host (Table 1).

Table 1. Proportional water content (%) of control, hosts and *V. a. ssp. album*

Fruit trees	Proportional water content (%)			
	Control	Host	<i>V. a. ssp. album</i>	F and P
Breaburn apple	24.761±2.55 b	21.948±0.849 b	86.499±9.019 a	F _{2,9} =Infinity P<0.0001
Ankara pear	28.921±6.16 b	25.879±0.923 b	67.48±6.6502 a	F _{2,9} =Infinity P<0.0001
Hawthorn	16.66±5.03 b	14.217±1.744 b	89.266±5.141 a	F _{2,9} =803.18 P<0.0001

One-way analysis of variance (ANOVA) and DUNCAN tests were applied to the data and the differences between the means were calculated at the significance level of P<0.05.

3.2. Chlorophyll a-b

Chlorophyll a content of *V. a. ssp. album* was found to be lower than control and hosts for Breaburn apple and Ankara pear. However, Chlorophyll a content of *V. a. ssp. album* was found to be similar to the host for hawthorn (Table 2). On the other hand, Chlorophyll b content of *V. a. ssp. album* was lower than control and similar to the host for Breaburn apple while it was similar to the control and host for Ankara pear and hawthorn.

Table 2. Chlorophyll a-b contents (mg/g) of control, hosts and *V. a. ssp. album*

Fruit trees	Chlorophyll a content (mg/g)				Chlorophyll b content (mg/g)			
	Control	Host	<i>V. a. ssp. album</i>	F and P	Control	Host	<i>V. a. ssp. album</i>	F and P
Breaburn apple	15.5±1.0 a	13.8±0.8 a	3.7±0.4 b	F _{2,6} = 65.06 P<0.0001	4.3±0.5 a	1.5±0.3 b	1.0±0.18 b	F _{2,6} = 25.38 P<0.001
Ankara pear	23.5±0.8 a	14.3±2.4 b	5.3±0.8 c	F _{2,6} = 32.84 P<0.001	0.03±0.07 a	1.7±0.4 a	0.3±0.1 a	F _{2,6} = 3.10 P=0.1192
Hawthorn	22.1±2.3 a	9.2±1.6 b	10.3±0.3 b	F _{2,6} = 18.30 P<0.01	1.9±0.2 a	2.8±0.5 a	3.0±0.2 a	F _{2,6} = 2.39 P=0.1722

One-way analysis of variance (ANOVA) and DUNCAN tests were applied to the data and the differences between the means were calculated at the significance level of P<0.05.

3.3. Total chlorophyll and caroteneoid content

Although the total chlorophyll and carotenoid contents of *V. a. ssp. album* were found less than control and host for Breaburn and Ankara pear, the total chlorophyll of *V. a. ssp. album* was resembling the host for hawthorn. The carotenoid content of *V. a. ssp. album* was higher in host for hawthorn (Table 3).

Table 3. Total Chlorophyll and carotenoid contents of control, hosts and *V. a. ssp. album*

Fruit trees	Total Chlorophyll content (mg/g)				Carotenoid content (mg/g)			
	Control	Host	<i>V. a. ssp. album</i>	F and P	Control	Host	<i>V. a. ssp. album</i>	F and P
Breaburn apple	19.8±1.3 a	15.3±0.6 b	4.8±0.5 c	F _{2,6} =68.29 P<0.0001	6.1±0.4 a	3.0±0.1 b	1.5±0.1 c	F _{2,6} = 73.82 P<0.0001
Ankara pear	23.4±1.4 a	16.0±2.5 b	5.7±0.8 c	F _{2,6} = 24.48 P<0.01	5.2±0.2 a	3.1±0.4 b	1.1±0.2 c	F _{2,6} = 37.98 P<0.001
Hawthorn	24.1±2.2 a	12.0±2.1 b	13.3±0.6 b	F _{2,6} =12.8 P<0.01	5.0±0.4 a	2.2±0.2 c	3.4±0.1 b	F _{2,6} = 23.94 P<0.01

One-way analysis of variance (ANOVA) and DUNCAN tests were applied to the data and the differences between the means were calculated at the significance level of P<0.05.

3.4. Proline and total amino acid concentrations

In all three fruits, proline concentration of *V.a. ssp. album* was observed to be higher than control and host (Table 4).

Table 4. Proline concentration (mg/g) of control, hosts and *V. a. ssp. album*

Fruit trees	Proline concentration (mg/g)			
	Control	Host	<i>V. a. ssp. album</i>	F and P
Breaburn apple	0.02±0.05 b	0.04±0.001 b	0.11±0.001 a	F _{2,6} =7.69 P<0.05
Ankara pear	0.03±0.002 b	0.05±0.002 b	0.41±0.02 a	F _{2,6} =134.61 P<0.0001
Hawthorn	0.04±0.01 b	0.01±0.002 b	0.25±0.02 a	F _{2,6} =62.58 P<0.0001

One-way analysis of variance (ANOVA) and DUNCAN tests were applied to the data and the differences between the means were calculated at the significance level of P<0.05.

Total amino acid concentration of *V. a. ssp. album* was calculated to be less than control in three fruits, while it was lower than host for Breaburn apple, higher than host for Ankara pear and similar to the host for hawthorn (Table 5).

Table 5. Amino acid concentrations (mg/g) of control, hosts and *V. a. ssp. album*

Fruit trees	Total amino acid content (mg/g)			
	Control	Host	<i>V. a. ssp. album</i>	F and P
Breaburn apple	4.30±0.17 a	2.92±0.001 b	2.14±0.204 c	F _{2,6} = 37.17 P< 0.001
Ankara pear	3.95±0.37 a	0.430±0.051 c	2.630±0.219 b	F _{2,6} =Infinity P<0.0001
Hawthorn	4.45±0.01 a	1.550±0.106 b	1.59±0.206 b	F _{2,6} = 91.31 P<0.0001

One-way analysis of variance (ANOVA) and DUNCAN tests were applied to the data and the differences between the means were calculated at the significance level of P<0.05.

3.5. Ascorbic acid concentration (mg/g)

Reduced ascorbic acid concentration of *V. a. ssp. album* in three fruits was determined to be significantly lower than control (Table 6). Reduced ascorbic acid of *V. a. ssp. album* was found similar to the host in Breaburn and hawthorn whereas it was determined to be less than the host for Ankara pear.

Table 6. Ascorbic acid concentrations (mg/g) of control, hosts and *V.a. ssp. album*

Fruit trees	Reduced ascorbic acid (mg/mL)				Total ascorbic acid (mg/mL)			
	Control	Host	<i>V. a. ssp. album</i>	F and P	Control	Host	<i>V. a. ssp. album</i>	F and P
Breaburn apple	0.155±0.003 a	0.147±0.002 b	0.142±0.001 b	F _{2,6} = 7.94 P<0.05	0.148±0.001 a	0.144±0.001 b	0.142±0.004 c	F _{2,6} =Infinity P<0.0001
Ankara pear	0.152±0.006 a	0.143±0.001 b	0.135±0.001 c	F _{2,6} = 28.28 P<0.001	0.150±0.003 a	0.144±0.001 b	0.137±0.001 c	F _{2,6} = 97.00 P<0.0001
Hawthorn	0.144±0.002 a	0.135±0.001 b	0.134±0.001 b	F _{2,6} = 20.73 P<0.01	0.145±0.003 a	0.136±0.001 b	0.134±0.001 b	F _{2,6} = 18.05 P<0.01

One-way analysis of variance (ANOVA) and DUNCAN tests were applied to the data and the differences between the means were calculated at the significance level of P<0.05.

Total ascorbic acid concentration of *V. a. ssp. album* was less than control and host for Breaburn and Ankara pear. However, the value was alike in the host for hawthorn.

3.6. Glucose and fructose concentrations

Glucose concentration of *V. a. ssp. album* was lower than control and host in three fruit species (Table 7). However, the fructose concentration of *V. a. ssp. album* was less than control and host for Breaburn apple and Ankara pear. Fructose concentration of *V. a. ssp. album*, however, was found to be similar with control for hawthorn.

Table 7. Glucose and fructose concentrations (mg/g) of control, hosts and *V. a. ssp. album*

Fruit trees	Glucose content (mg/g)				Fructose content (mg/g)			
	Control	Host	<i>V.a. ssp. album</i>	F and P	Control	Host	<i>V.a. ssp. album</i>	F and P
Breaburn apple	0.567±0.02 a	0.469±0.131 b	0.117±0.011 c	F _{2,6} =2191.6 P<0.0001	0.688±0.007 a	0.568±0.024 b	0.201±0.03 c	F _{2,6} = 108.50 P<0.0001
Ankara pear	0.445±0.08 a	0.261±0.076 b	0.059±0.014 c	F _{2,6} =11.93 P<0.01	0.406±0.002 a	0.317±0.017 b	0.125±0.01 c	F _{2,6} = 103.9 P<0.0001
Hawthorn	0.225±0.01 a	0.173±0.092 b	0.098±0.002 c	F _{2,6} =173.62 P<0.0001	0.313±0.005 a	0.258±0.081 b	0.075±0.01 b	F _{2,6} = 6.76 P<0.05

One-way analysis of variance (ANOVA) and DUNCAN tests were applied to the data and the differences between the means were calculated at the significance level of P<0.05.

3.7. Sulfhydryl concentrations

Sulfhydryl (SH) concentration of *V. a. ssp. album* was found to be lower than control and host in three fruits (Table 8).

Table 8. SH concentration (ppm) of control, hosts and *V. a. ssp. album*

Fruit trees	SH concentration (ppm)			F and P
	Control	Host	<i>V. a. ssp. album</i>	
Breaburn apple	0.142±0.0001 a	0.139±0.0004 b	0.136±0.0001 c	F _{2,6} = 43.00 P< 0.001
Ankara pear	0.101±0.0001 a	0.099±0.0001 b	0.097±0 c	F _{2,6} = 37.00 P< 0.001
Hawthorn	0.102±0 a	0.098±0.0001 b	0.097±0 c	F _{2,6} = Infity P< 0.001

One-way analysis of variance (ANOVA) and DUNCAN tests were applied to the data and the differences between the means were calculated at the significance level of P<0.05.

According to the results of statistical analysis, leaf proportional water content of *V. a. ssp. album* were 3-4 times higher than control and host in three fruits. In addition, leaf proportional water contents of control and host were found to be similar to each other. Leaf water status is intimately related to several leaf physiological variables, such as leaf turgor, growth, stomatal conductance, transpiration, photosynthesis and respiration [26]. Hawksworth and Wiens [6] showed that Dwarf mistletoe has conductance in the parasite was typically from 2 to 5 times that of the host. Mistletoes most successfully compete for a share of the host's water. It has been previously found that mistletoe on trees has a more than 3 fold higher transpiration rate, calculated by leaf surface, compared to the host. Compared with the control, the leaf relative water content of mistletoe was found much higher than its host [18,19,27-29].

According to our findings, *V. a. ssp. album* seemed to have a lower chlorophyll a content than control and hosts for Breaburn apple and Ankara pear. Chlorophyll a content of *V. a. ssp. album* was resembling the one measured for the hosts in hawthorn. Chlorophyll b content of *V. a. ssp. album*, however, was found to be lower than control and similar to the host for Breaburn apple. On the other hand, it was similar to the control and host for Ankara pear and hawthorn. The negative effects of compulsory hemiparasitic weeds on photosynthesis were also observed on the hosts. *Viscum album* is able to produce its own assimilates in the process of photosynthesis. *V. album* has a high transpiration rate, similar to many other parasitic plants and a low photosynthetic rate. In previous studies, the value of chlorophyll a in the mistletoe has been shown to be significantly lower than hosts. The invasion of parasite in the mango (host) has also been reported to alter the ratio between chlorophyll a and b, probably affecting the photosynthetic efficiency [9, 12, 30, 31]. In our study, chlorophyll a content of mistletoe was found to be higher than chlorophyll b while Oyetunji and Edagbo [18] showed that both host (Citrus, Irvingia) and African mistletoe (*Tapinanthus bangwensis*) have higher chlorophyll b contents.

The total chlorophyll and carotenoid contents of *V. a. ssp. album* were less than control and host for Breaburn and Ankara pear. However the total chlorophyll was measured to be similar to the host for hawthorn and *V. a. ssp. album*. This probably resulted from the low level photosynthesis activity. Total chlorophyll of *V. album* in hawthorn was similar to its host while carotenoid content was higher than host. *V. album* contains all of the pigments, chlorophyll a and b as well as carotenoid that are necessary for photosynthesis [32]. In a report published by Oyetunji and Edagbo, total chlorophyll content of the mistletoe and hosts exhibited values in which the total chlorophyll of the *Citrus* was significantly higher while the total chlorophyll content for the mistletoe-*Irvingia* relationship was not statistically different [18]. Semi parasite plant has total chlorophyll and carotenoids more much than control plants [19].

Proline content of *V. a. ssp. album* was higher than control and host in three fruits. This is probably due to the stress arising from abiotic factors for *V. a. ssp. album*. The increase of proline concentration promotes proline oxidation while it inhibits protein synthesis [33]. It was reported that proline is a nitrogen compound that increased in stress condition, participated in detoxification of free O₂ and contained protective properties in the stress conditions [34]. Many authors stated that the contents of proline and relative chlorophyll increased with the effect of oxidative stress [20, 35, 36]. Mistletoe-host associations have shown that the potassium and proline contents of mistletoes were the

highest among other major nutrients and osmotically active solutes, thus playing a vital role in osmotica of the parasite [37, 38]. In another study, it has been demonstrated that the concentration of proline in the leaf of Washington and Valencia increased after pesticide application [39]. *Erwinia amylovora* is a biological stress factor causing increase in the content of proline and ascorbic peroxidase in apples and pears [40]. The results of the present study are in accordance with Murugan et al. [19] who showed that semi parasite plant has much more proline than control plants.

Total amino acid concentrations of *V. a. ssp. album* were found to be less than control in three fruits, while it was less than host for Breaburn apple, higher than host for Ankara pear. It was also found to be similar to the host for hawthorn. Since mistletoe takes the required nutrients from its host, amino acid content of fruit tree is reduced. A significantly lower content of leaf protein in most species of the infected trees was observed when compared to the uninfected trees [41]. *Dendrophthoe falcate*, which is also a semiparasitic plant, increased amino acid content in its host. While the total protein, phenols, flavonoids also showed a defense related response in the plant, the chlorophyll content showed a decrease in the infected plant [19]. In autumn, *V. album* collected from Crataegus had the lowest protein content whereas protein content in the leaves of mistletoe hosted by Acer, Populous, and Robinia did not change significantly in different seasons [42].

Total and reduced ascorbic acid content of *V. a. ssp. album* in three fruits were found to be lower than the uninfected plants. Their content in hosts were lower than control. This is most likely due to the taking of ascorbic acid by *V. a. ssp. album* from their host. It has been reported that some hosts of *Viscum album* subspecies, such as apples, cherries, pears and plums, contain ascorbic acid [43], and that ascorbic acid content decreases in stress conditions (salt and metal) in some plant species while some increase it as an antioxidant defense mechanism [44, 45].

Glucose and fructose contents of *V. a. ssp. album* in three fruits were found to be lower than control and host. This observation may be explained by the fact that glucose and fructose can be intaken by *V. a. ssp. album* from its host. Glucose, galactose, fructose, saccarose, galacturonic acid, and pectine were found in the leaf of mistletoe [46, 47]. The quantity of the fructose and glucose in the mistletoe that was parasitic on Irvingia were higher than that on Citrus. However, the host plants had more of these reducing sugars than the semiparasite. This further suggests that the semi parasite depends on the host for little sugar supply from its host to complement its weak photosynthetic activities [18]. In a study conducted by Murugan et al., reduction in soluble sugar contents were noted in the infected mango plant by the parasite *Dendrophthoe falcata*. Under parasitic stress conditions, total soluble sugar content decreased considerably. Parallely in the control, the total soluble sugar content was remarkably higher. A significant decrease in total soluble sugar content under biotic stress was observed in the infected host [19].

The content of SH in three fruits was highest in the leaf of control and they were at a lower level in their hosts while they were the lowest in the leaf of *V. a. ssp. album*. The fact that SH content of hosts were lower than controls was probably resulted from the mistletoe stress factor. SH content in the plant tissues increases in spring compared to winter months. SH content of the plants may be elevated by the cold and water stress in the photosynthesis reaction. Reactive oxygen species and free radicals oxidate nucleic acids, proteins and lipids in the oxidative stress period. It has also been determined by researchers that antioxidants may be reduced in the event of stress in different environments [48-50]. SH contents in the leaf of Washington and Valencia were found to be at higher levels before pesticid application [39]. *V. album* may have detrimental effect on leaf structure, physiology and chemical composition of highly infected host trees [51]. It is widely appreciated that the chemical composition of mistletoe is not stable and it depends not only on biosynthesis but also on the type of host plant and growth conditions, such as ambient temperature, carbon dioxide concentration and season of the year [52, 53].

4. Conclusion

Due to the fact mistletoe uses the water and nutrients of the host, the proportional water content, chlorophyll a-b, caratoneid, proline, total amino acid, ascorbic acid, glucose, fructose and SH contents of *V. a. ssp. album* showed a variation. At the same time, these parameters appear to be

affected by the morphological and biological characteristics of hosts as well as the soil properties and ecological properties of the region.

Acknowledgement

This project was supported by Kahramanmaraş Sutcu Imam University, Scientific Research Projects Unit with the project number of 2013/1-22M.

ORCID

Tamer Üstüner: [0000-0003-3584-4249](https://orcid.org/0000-0003-3584-4249)

Sema Düzenli: [0000-0002-2506-6415](https://orcid.org/0000-0002-2506-6415)

References

- [1] A.G. Miller (1982). *Arceuthobium* Bieb. and *Viscum album* L. (Flora of Turkey and the East Aegean Islands), University press, Edinburgh.
- [2] M. Calder and P. Bernhardt (1983). *The Biology of mistletoes*. Sydney, New York, London, Paris, San Diego, San Francisco, Sao Paulo, Tokyo, Toronto: Academic press.
- [3] D.B. Fischer (1983). Year round collection of willow sieve tube exudate, *Planta* **159**, 529-533.
- [4] J.R. Ehleringer, E.D. Schultze, H. Ziegler, O.L. Lange, G.D. Farquhar and I.R. Cowar (1985). Xylem-tapping mistletoes: Water or nutrient parasites, *Science* **227**, 1479-1481.
- [5] F.G. Hawksworth and R.F. Scharpf (1986). Spread of European mistletoe (*Viscum album* L.) in California,U.S.A, *Eur. J. Forest Pathol.* **6**,1-5.
- [6] F.G. Hawksworth and D. Wiens (1996). Dwarf mistletoes (*Viscum album* L.), Biology, pathology and systematics. U.S.A. Department of Agriculture forest service, *Agricultural Handbook* 709.
- [7] D.M. Watson (2001). Mistletoe A keystone resource in forests and woodlands worldwide, *Annual Rev. Ecol. Syst.* **32**, 219-249.
- [8] J. Briggs (2003). Christmas curiosity or medical marvel? A seasonal review of mistletoe, *Biologist* **50**, 249-254.
- [9] D. Zuber (2004). Biological flora of central Europe: *Viscum album* L., *Flora* **199**, 188-203.
- [10] D. Zuber and A. Widmer (2009). Phylogeography and host race differentiation in the European mistletoe (*Viscum album* L.), *Mol. Ecol.* **18**, 1946-1962002E|01.
- [11] C. Türe, H. Böcük and A. Zerrin (2010). Nutritional relationships between hemi-parasitic mistletoe and some of its deciduous hosts in different habitats. Vicas, S., *Biol. Section Bot.* **65**, 859-867.
- [12] H.S. Heide-Jorgensen (2004). About *Viscum* a mistletoe *Viscum* dk.http://www.Viscum.dk/eng-sider/about_Viscum.htm.
- [13] N. Zeybek (1985). *Pharmaceutical botany, closed seeded plants system (Angiospermae) and important components*, Ege University, Pharmacy Faculty. Ege University printing house, İzmir-Turkey, Number, 1.
- [14] T. Üstüner (2003). Identification and density of *Viscum* species in Nigde province, *The J. Turk. Weed Sci.* **6** (2),45-53.
- [15] T. Üstüner, S. Düzenli and Y.E. Kitis (2015). Determination of infection rate of mistletoe (*Viscum album*) on hosts in Nigde province, *The J. Turk. Weed Sci.* **18(1)**, 6-14.
- [16] P. Escher, M. Eiblmeier, I. Hetzger and H. Rennenberg (2004). Seasonal and spatial variation of carbohydrates in mistletoes (*Viscum album*) and the xylem sap of its hosts (*Populus euamericana* and *Abies alba*), *Physiol. Plantar.* **120**, 212-219.
- [17] H. Ziegler, J. Weber and U. E. Lüttge (2009). Thermal dissipation probe measurements of sap flow in the xylem of trees documenting dynamic relations to variable transpiration given by instantaneous weather changes and the activities of a mistletoe xylem parasite, *Trees.* **23**, 441-450.
- [18] O.J. Oyetunji and D.E. Edagbo (2013). Comparative ecophysiological study of *Tapinanthus bangwensis*, [Engl. And R. Krause] Danser (African Mistletoe) On two host plants, *J. of Chem. Bio. Phy. Sci.* **3**, 1933-1941.
- [19] R. Krishnan, A.S. Lubaina and K. Murugan (2014). Interaction between hemiparasitic (*Dendrophthoe falcata* (L.) Etting.) on mangifera Indica linn. Some observations. *World J. Pharm. Pharmaceut. Sci.* **3(6)**, 585-607.

- [20] R.K. Sairam, K.V. Rao and G.C. Srivastava (2002). Differential response of wheat genotypes to long term salinity stress in relation to oxidative stress, antioxidant activity and osmolyte concentration, *Plant Sci.* **163**, 1037-1046.
- [21] D.M. Arnon (1949). Copper enzymes in isolated chloroplasts poly phenoloxidase in *Beta vulgaris*, *Plant Physiol.* **24**, 1-15.
- [22] H.K. Lichtenthaler and A.R. Wellburn (1983). Determinations of total carotenoids and chlorophyll a and b of leaf extracts in different solvents, *Biochem. Soc. Transac.* **11**, 591-592.
- [23] I. Cakmak and H. Marschner (1992). Magnesium deficiency and high light intensity enhance activities of superoxide dismutase, ascorbate peroxidase and glutathione reductase in bean leaves, *Plant Physiol.* **98**, 1222-1227.
- [24] J.R. Spies (1957). Colorimetric procedures for amino acids. In: Methods in enzymology. III. (Ed. by S. P. Colowick and N. O. Kaplan), *Academic press*, New York, 468-471.
- [25] L.S. Bates, R.P. Waldren and I.D. Teare (1973). Rapid determination of free proline for water-stress Studies, *Plant Soil* **39**, 205-207.
- [26] P.J. Kramer and J.S. Boyer (1995). Water relation of plants and soils. *Academic press*, San Diego, 495.
- [27] E.D. Schulze, N.C. Turner and G. Glatzel (1984). Carbon, water and nutrient relationships of two mistletoes and their hosts: A hypothesis, *Plant Cell environ.* **7**, 293-299.
- [28] S.M. Hosseini, D. Kartoolinejad, S.K. Mirnia, Z. Tabibzadeh, M. Akbarinia and F. Shayanmehr (2008). The European mistletoe effects on leaves and nutritional elements of two host species in Hyrcanian forests. *Silva Lusitana*, © EFN, Lisboa, *Portugal* **16** (2), 229-237.
- [29] G. Glatzel and B.W. Geils (2009). Mistletoe ecophysiology: Host-parasite interactions. *Botany.* **87**, 10.
- [30] K.U. Tennakoon and J.S. Pate (1996). Effect of parasitism by a mistletoe on the structure and functioning of branches of its host-plant. *Cell and environment.* **19**, 517-528.
- [31] A. Chatterjee and S.K. Ghosh (2008). Alterations in biochemical components in mesta plants infected with yellow vein mosaic disease, *Braz J. Plant Physiol.* **20** (4), 267-275.
- [32] H. Becker (2000). European mistletoe: Taxonomy, host trees, parts used, physiology. In: Mistletoe: The genus *Viscum*, ed. A. Büssing, Harwood Academic publishers.
- [33] N. Smirnoff and S.V. Colombe (1988). Drought influences the activity of the chloroplast hydrogen peroxide scavenging system, *J. Experiment. Bot.* **39**, 1097-1108.
- [34] H.J. Bohnert and E. Sheveleva (1998). Plant stress adaptations making metabolism move, *Curr. Opinion in Plant Biol.* **1**, 267-277.
- [35] R.K. Sairam, D.S. Shukla and D.C. Saxena (1998). Stress induced injury and antioxidant enzymes in relation to drought tolerance in wheat genotypes, *Biol. Plantar.* **40**, 357-364.
- [36] R.K. Sairam and D.C. Saxena (2000). Oxidative stress and antioxidants in wheat genotypes: Possible mechanism of water stress tolerance, *J. Agronomy Crop Sci.* **184** (1), 55-61. DOI:10.1046/j.1439-037x.2000.00358.xView/save citation.
- [37] L.C. McDowell (1964). Physiological relationships between dwarf mistletoe and ponderosa pine, Oregon state university, Corvallis, OR, Oregon State University, **68**.
- [38] J.M. Hibberd and W.D. Jeschke (2001). Solute flux into parasitic plants, *J. Exp. Botany.* **52** (363), 2043-2049.
- [39] A. Fidan and S. Düzenli (2007). The effects of some pesticides on the physiological and anatomical properties of citrus varieties. Cukurova university, Institute of natural and applied sciences, MSc thesis, Page: 40. Adana-Turkey.
- [40] E. Karacif and N. Boyraz (2012). Determination of levels of some antioxidant and enzymes after infection with *Erwinia amylovora* of apple and pear varieties. Selcuk university, The graduate school of natural and applied science, The degree of master of science, Plant protection, June, Konya-Turkey.
- [41] J. Patykowski and J. Kołodziejek (2013). Comparative analysis of antioxidant activity in leaves of different hosts infected by mistletoe (*V. album* L. subsp. *album*), *Archives Biol. Sci.* **65**, 851.
- [42] J. Patykowski and J. Kołodziejek (2016). Changes in antioxidant enzyme activities of European mistletoe (*V. album* subsp. *album*) leaves as a response to environmental stress caused by pollution of the atmosphere by nitrogen dioxide, *Pol. J. Environ. Stud.* **25** (2), 725-732.
- [43] W.C. Evans (1989). Trease and Evans' pharmacognosy, Bailliere Tindall, London, **13. Edition**, 665.
- [44] R.K. Sairam, G.C. Srivastava, S. Agarwal and R.C. Meena (2005). Differences in antioxidant activity in response to salinity stress in tolerant and susceptible wheat genotypes, *Biol. Plantar.* **49**, 85-91.
- [45] M. Tiryakioglu, S. Eker, F. Ozkutlu, S. Husted and I. Cakmak (2006). Antioxidant defense system and cadmium uptake in barley genotypes differing in cadmium tolerance, *J. Trace Elem. Med. Biol.* **20**, 181-189.
- [46] O.I. Popova and D.A. Murav'eva (1990). Polysaccharides of mistletoe (*Viscum album*) polysaccharide fractions from European mistletoe and firmiana, *Farmatsiya* **39** (5), 41-44.

- [47] O.I. Popova (1991). Phenolic acids of *Viscum album*, *Khim, Prir. Soedin.* **1**, 139-140.
- [48] I.K. Smith, A.C. Kendall, A.J. Keys, J.C. Turner and P.J. Lea (1985). The regulation of the biosynthesis of glutathione in leaves of barley (*Hordeum vulgare* L.), *Plant Sci.* **41**,11-17.
- [49] D.M. Hodges and C.F. Forney (2000). The effects on ethylene, depressed oxygene and elevated carbon dioxide on antioxidant profiles of senescing spinach leaves, *J. Experiment. Bot.* **51** (334), 645- 655.
- [50] I. Meram and S. Aktaran (2002). Effects of free radicals on biomolecules, *Archive* **11**, 299.
- [51] M.G. Daryaei and E.S. Moghadam (2012). Effects of mistletoe (*Viscum album*) on leaves and nutrients content of some host trees in Hyrcanian forests (Iran). *International Journal of Agriculture: Res. Review* **2** (3), 85-90.
- [52] P. Stypiński (1997). Biology and ecology of mistletoe (*Viscum album*, Viscaceae) in Poland, Instytut Botaniki im. W. Szafera PAN, Kraków, 1-117.
- [53] M. Łuczkiwicz, W. Cisowski, P. Kaiser, R. Ochocki and A. Piotrowski (2001). Comparative analysis of phenolic acids in mistletoe plants from various hosts, *Acta Pol. Pharm.* **58**, 373-379.

A C G
publications

© 2017 ACG Publications