

Synthesis of new amino-1,5-benzodiazepine and benzotriazole derivatives from dimedone

Norah Bennamane, Rachedine Kaoua, Lamouri Hammal and Bellara
Nedjar-Kolli*

Laboratory of Applied Organic Chemistry, University of Sciences and Technology
Houari Boumediene, BP 32, El-Alia, Algiers, Algeria.

(Received July 14, 2008; Revised September 29, 2008; Accepted October 14, 2008)

Abstract: High-yielding syntheses of two unpublished series of amino-1,5-benzodiazepines derivatives **3** and 1,2,3-benzotriazoles **4** bearing a dimedone moiety are respectively reported. Derivatives **3** were obtained by intramolecular cyclisation using cyanogenbromide of enaminone **2**. On the other hand, when enaminones **2** were treated with sodium nitrite under acidic conditions, cyclisation led to benzotriazole **4**.

Keywords: Dimedone; benzotriazoles; benzodiazepines, cyanobromide; enaminones.

1. Introduction

The benzodiazepine nucleus is a pharmacophoric scaffold and represent a class of heterocycles with a wide range of biological applications.¹ Many of them are widely used as anticonvulsant, antianxiety, sedative, antidepressive, hypnotic and neuroleptic agents.²⁻⁵ Some heterocycles containing benzodiazepines moiety were reported to possess anti inflammatory⁶, antiviral⁷, anti-HIV-1⁸, antimicrobial⁹ and antitumor¹⁰ activities. Other than their biological importance, benzodiazepines are valuable syntons for the preparation of fused ring compounds, such as triazolo¹¹, thiazolo¹², imidazo¹³ and pyrimido-benzodiazepines.¹⁴ It has been noticed that introduction of an additional ring to the benzodiazepine core tends to exert profound influence in conferring novel biological activities in these molecules.¹¹⁻¹⁵ Although many methods for synthesizing benzodiazepine ring systems have been reported, they continue to receive a great deal attention.^{9,16-18}

Another class of heterocycles used as scaffolds in medicinal chemistry is devoted to benzotriazole derivatives. They exhibit useful pharmacological properties and clinical applications.¹⁹⁻²²

* E-Mail: bellara_kollidz@yahoo.fr


In addition to these considerable biological applications, benzotriazoles are important intermediates, protecting groups and final products in organic synthesis.²³

In the course of our research and as part of our program involving the synthesis of new heterocyclic compounds having potential biological interest, we have already reported the synthesis of many pyranobenzodiazepines²⁴⁻³¹, pyranobenzodiazepin-2-thione and benzotriazoles derivatives³² using enaminolactones as starting material. Recently, reactions of 5-amino-3-methyl-1H-pyrazole with dimedone and aldehydes was reported to give regioselectively tricyclic linear 3,7,7-trimethyl-4,7,8,9-tetrahydro-2H-pyrazolo[3,4-b]quinolin-5(6H)-ones.³³

In pursuance of our interests for investigating the reactivity of enaminones towards electrophile reagents we now extend the scope of these reactions to an other active compound the 5,5-dimethylcyclohexan-1,3-dione (dimedone). They appear in a natural tricyclic heterocycle Benzylhiocrellidone isolated from sponge *Crella spinulata*³⁴⁻³⁵ and is an effective synthon in the synthesis of several polyheterocyclic compounds.³⁶⁻³⁸ As expected, we get access to 1,5-benzodiazepines **3** and Benzotriazoles **4** fused and linked respectively to a dimedone moiety (Scheme 1).

2. Results and discussion


We have previously demonstrated that the enaminolactones intermediates allowed versatile access to different heterocycles²⁴⁻³⁰, depending on the nature of electrophile used. In this work intermediate **2**, obtained by reaction of dimedone with substituted *o*-phenylenediamines under similar conditions to already described procedure²⁴, was used as starting point for the synthesis of benzodiazepines **3** and benzotriazoles **4** according to our synthetic pathways which are outlined in the scheme 1.



Scheme 1: Intramolecular cyclisations of enaminones **2**

Treatment of **2** with 1 equivalent of cyanobromide proceeds smoothly in refluxing, yielding by filtration product **3** in high purity.

As shown in scheme 2, the formation of seven membered ring **3** was due to the nucleophilic attack on the iminium ion by the adjacent electron-rich double bond of the dimedone moiety.



Scheme 2: Postulated mechanism for the formation of compounds **3** and **4**

Table 1: Physico-chemical data of derivatives **3a-c** and **4a-c**

Compounds	X	Yields [%]	Time (h)	mp (°C)
3a	H	45	4	274-276
3b	CH_3	49	4	280-282
3c	Cl	30	6	292-294
4a	H	86	1	180-182
4b	CH_3	77	1	190-192
4c	Cl	65	1	200-202

Similarly the reaction of binucleophilic enaminone intermediates **2** with sodium nitrite under acidic conditions according to a described procedure,³⁶⁻³⁸ allowed in all cases a single product **4** bearing a dimedone moiety in good yield (table-1). The probable mechanism in this case seems to be the diazotization of the primary amine of **2**, thus the formed diazo salt undergo cyclization, by nucleophilic attack of the secondary amino group leading to a triazole nucleus. The formation of benzotriazepine that would result from an intramolecular cyclization through the double bond of dimedone moiety as reported in the case of formation of benzodiazepines **3** was not observed. The purity and structures of the newly synthesized compounds were established by elemental analysis, IR, ^1H , ^{13}C NMR DEPT, and mass spectroscopic analysis.

Compounds **3a-c** exhibit in particular a ^1H NMR spectrum with a broadened singlet around 9.6-9.8 ppm characterizing the presence of NH^+ . In addition the complete lack of the ethylenic proton at the C-2 of the dimedone ring in the ^1H NMR is accompanied by absorption for the quaternary

carbon of junction C-11a and C-11 of amidine bond at ~ 96 ppm and ~ 174 ppm respectively. The ^1H NMR spectra of **4a-c** not only showed the absence of both signals of NH at 8.06-8.13 ppm and NH_2 at 4.75-5.19 ppm, but also indicate the presence of C(2)-H signal at 6.55-6.58 ppm. Moreover ^{13}C NMR spectrum contains a signal around 112.8-114.8 ppm assigned to the tertiary ethylenic carbon C-2 in the dimedone unit thus substantiate the formation of all suggested compounds.

3. Conclusion

In conclusion, enamines **2** resulting from the commercially available dimedone and *o*-phenylenediamines derivatives constitute an interesting and efficient precursor for the synthesis of new 1,5-benzodiazepines and benzotriazoles systems which are condensed and linked respectively to a dimedone moiety. These products are expected to possess biological activities.

4. Experimental

Melting points were determined using a Stuart Scientific SMP1 melting point apparatus and were uncorrected. The IR spectra were recorded in KBr pellets on a Perkin-Elmer FTIR Paragon 1000 spectrometer. The ^1H NMR spectra (400 MHz) and ^{13}C NMR (100.6 MHz) were run in DMSO- d_6 on a Bruker Avance II spectrometer. Chemical shifts are given in ppm relative to TMS as internal standard. The positive-mode electrospray ionisation (ESI) mass spectra were recorded on a Perkin-Elmer SCIEX API 300 spectrometer. The elemental analysis data were performed on a Thermo Electron Flash EA 1112. Chemicals were purchased from Aldrich and Acros Organics and used without further purification.

General procedure for the preparation of compounds **2**:

To a magnetically stirred solution of **1** (20 mmol, 1 eq) in 100 ml of absolute ethanol was added 4- substituted -1,2-phenylenediamine compounds (20 mmol, 1eq), and the mixture was allowed to react at room temperature for 12 h (as monitored by TLC). The reaction mixture was concentrated by rotatory evaporation to a volume of 20 ml and then stored in a refrigerator for 3 h. The precipitate obtained was filtered, washed several times with diethyl ether and recrystallized from ethanol to afford compounds **2**. Yields: 59-65 %.

General procedure for the preparation of compounds **3**:

Cyanobromide 3M soln. EtOH (2.3 mmol, 1.15 eq) was added dropwise to a magnetically stirred solution of **2** (2 mmol, 1 eq) in ethanol 15 ml over a period of 30 min at room temperature. The mixture was then heated under reflux for 4-6 h. The precipitate formed was collected by filtration. Recrystallization from the proper solvent furnished the desired products **3a-c** as yellow solid.

11-amino-3,3-dimethyl-2,3,4,5-tetrahydro-1H-dibenzo[b,e][1,5]diazepin-1-one hydrobromide (3a). Yield: 45%. IR (KBr): 3443-3250 (NH, NH_2); 1652 ($\text{C}=\text{O}$), 1580 ($\text{C}=\text{N}$) cm^{-1} . ^1H NMR (400 MHz, DMSO- d_6) (δ /ppm): 1.01 (s, 6H, 2 CH_3), 2.34 (s, 2H, NC- CH_2), 2.73 (s, 2H, OC- CH_2), 6.89-7.22 (m, 4H, Ar), 8.55 (s, br, 1H, NH), 9.72 (s, br, 1H, NH), 10.65 (s, 1H, NH_2), 10.73 (s, 1H, NH_2). ^{13}C NMR (100.6 MHz, DMSO- d_6) (δ /ppm): 27.08, 29.21, 45.99, 50.65, 96.04, 121.31, 122.57, 126.06, 128.11, 130.62, 131.02, 164.41, 174.28, 195.93. MS (ESI) m/z 256.5 [(M-HBr) $^+$, 16%]. Anal. Calcd. For. $\text{C}_{15}\text{H}_{17}\text{N}_3\text{O}$, HBr: C, 53.58%; H, 5.40%; N, 12.50%. Found: C, 53.37 %; H, 5.20%; N, 12.29%.

11-amino-3,3,7-trimethyl-2,3,4,5-tetrahydro-1H-dibenzo[b,e][1,5]diazepin-1-one hydrobromide (3b). Yield: 49%. IR (KBr): 3431-3237 (NH,NH₂), 1667 (C=O), 1596 (C=N) cm⁻¹. ¹H NMR (400 MHz, DMSO-d₆) (̴/ppm): 1.01 (s, 6H, 2CH₃), 2.23 (s, 3H, Ar-CH₃), 2.34 (s, 2H, NC-CH₂), 2.71 (s, 2H, OC-CH₂), 6.70-7.05 (m, 3H, Ar), 8.53 (s, br, 1H, NH), 9.67 (s, br, 1H, NH), 10.61 (s, 1H, NH₂), 10.81 (s, 1H, NH₂). ¹³C NMR (100.6 MHz, DMSO-d₆) (̴/ppm): 20.13, 27.06, 29.19, 46.00, 50.64, 95.48, 121.45, 122.53, 126.45, 128.22, 130.47, 138.04, 164.41, 173.65, 195.85. MS (ESI) m/z 270.5 [(M-HBr)⁺, 22%]. Anal. Calcd. For. C₁₆H₁₉N₃O, HBr: C, 54.87 %; H, 5.76 %; N, 12.00 %. Found: C, 54.74 %; H, 5.74 %; N, 11.87 %.

11-amino-7-chloro-3,3-dimethyl-2,3,4,5-tetrahydro-1H-dibenzo[b,e][1,5]diazepin-1-one hydrobromide (3c). Yield: 30%. IR (KBr): 3460-3255 (NH,NH₂), 1660 (C=O), 1590 (C=N) cm⁻¹. ¹H NMR (400 MHz, DMSO-d₆) (̴/ppm): 1.02 (s, 6H, 2CH₃), 2.35 (s, 2H, NC-CH₂), 2.73 (s, 2H, OC-CH₂), 6.90-7.22 (m, 3H, Ar), 8.60 (s, br, 1H, NH), 9.81 (s, br, 1H, NH), 10.75 (s, 1H, NH₂), 10.92 (s, 1H, NH₂). ¹³C NMR (100.6 MHz, DMSO-d₆) (̴/ppm): 27.04, 29.20, 46.15, 50.69, 95.72, 121.01, 121.8, 122.50, 126.00, 129.8, 130.44, 164.92, 173.85, 195.78. MS (ESI) m/z 290.5 [(M-HBr)⁺, 10%]. Anal. Calcd. for. C₁₅H₁₆N₃OCl, HBr: C, 48.60 %; H, 4.62 %; N, 11.34 %. Found: C, 48.32 %; H, 4.42 %; N, 11.09 %.

General procedure for the preparation of compounds 4:

To an ice-cooled (0-5°C) and stirred solution of (2 mmol, 1eq) of the appropriate enaminones **2** in 6 ml of an 1M solution of HCl a solution of sodium nitrite 0.165g (2.4 mmol, 1.2 eq) in 10 ml of H₂O was added drop by drop. After 20 min, the ice-bath was removed and the resulting solution was stirred for an additional hour. The precipitate obtained was collected by filtration and washed with H₂O. Recrystallization from ethyl acetate/petroleum ether gave the target compounds 4a-c.

3-(1H-Benzotriazol-1-yl)-5,5-dimethylcyclohex-2-en-1-one (4a). Yield: 86%. IR (KBr): 1669 (C=O), 1629 (C=N) cm⁻¹. ¹H NMR (400 MHz, DMSO-d₆) (̴/ppm): 1.17 (s, 6H, 2CH₃), 2.43 (s, 2H, NC-CH₂), 3.25 (s, 2H, OC-CH₂), 6.59 (s, 1H, NC=CH), 7.54-8.25 (m, 4H, Ar). ¹³C NMR (100.6 MHz, DMSO-d₆) (̴/ppm): 27.63, 32.80, 40.00, 50.12, 112.80, 114.34, 120.13, 125.50, 129.56, 130.87, 146.16, 152.18, 198.05. MS (ESI) m/z 242.0 [(M⁺-H)⁺, 100%], 264.0 [(M⁺-Na)⁺, 65%]. Anal. Calcd. For. C₁₄H₁₅N₃O: C, 69.69 %; H, 6.27 %; N, 17.41 %. Found: C, 69.88 %; H, 6.22 %; N, 17.25 %.

5,5-dimethyl-3-(5-methyl-1H-benzotriazol-1-yl)cyclohex-2-en-1-one (4b). Yield: 77%. IR (KBr): 1667 (C=O), 1626 (C=N) cm⁻¹. ¹H NMR (400 MHz, DMSO-d₆) (̴/ppm): 1.15 (s, 6H, 2CH₃), 2.41 (s, 2H, NC-CH₂), 2.50 (s, 3H, Ar-CH₃), 3.23 (s, 2H, OC-CH₂), 6.55 (s, 1H, NC=CH), 7.53-7.99 (m, 3H, Ar). ¹³C NMR (100.6 MHz, DMSO-d₆) (̴/ppm): 20.68, 27.65, 32.76, 40.06, 50.11, 112.37, 113.82, 119.05, 129.29, 131.38, 135.36, 146.78, 152.21, 198.08. MS (ESI) m/z 256.5 [(M⁺-H)⁺, 100 %], 278 [(M⁺-Na)⁺, 47%]. Anal. Calcd. for. C₁₅H₁₇N₃O: C, 70.56 %; H, 6.71 %; N, 16.46 %. Found: C, 70.74 %; H, 6.55 %; N, 16.28 %.

3-(5-chloro-1H-Benzotriazol-1-yl)-5,5-dimethylcyclohex-2-en-1-one (4c). Yield: 65%. IR (KBr): 1670 (C=O), 1630 (C=N) cm⁻¹. ¹H NMR (400 MHz, DMSO-d₆) (̴/ppm): 1.16 (s, 6H, 2CH₃), 2.43 (s, 2H, NC-CH₂), 3.22 (s, 2H, OC-CH₂), 6.58 (s, 1H, NC=CH), 7.71-8.40 (m, 3H, Ar). ¹³C NMR (100.6 MHz, DMSO-d₆) (̴/ppm): 27.61, 32.81, 39.92, 50.10, 114.38, 115.01, 119.41, 129.74, 129.81, 129.87, 146.83, 151.82, 198.01. MS (ESI) m/z 276.0 [(M⁺-H)⁺, 100%], 298.0 [(M⁺-Na)⁺, 43%]. Anal. Calcd. For. C₁₄H₁₄ClN₃O: C, 60.98 %; H, 5.12 %; N, 15.24 %. Found: C, 60.72 %; H, 4.93 %; N, 14.92 %.

Acknowledgments

The authors sincerely thanks Professor Gerard Coudert (I.C.O.A, University of Orleans, France) for providing laboratory facilities and for his help in obtaining the IR, NMR, and spectral data.

References

- [1] Randall, L. O. *Psychopharmacological Agents*, ed: M. Gordon, New York, Academic Press, **1974**, 3, 175-281.
- [2] Fryer, R. I. *Bicyclic Diazepines*, In: *Comprehensive Heterocyclic Chemistry*, ed: E.C. Taylor, Wiley, New York, **1991**, 50, Chapter II.
- [3] Devi, T.K.; Achaiah, G.; Reddy, V. M. Synthesis of 6-alkyl/aryl-5,11-dihydro-9-nitro [1] benzopyrano [2,3-b] [1,5] benzodiazepine-1,3-ones as possible antipsychotic agents. *J. Indian Chem. Soc.* **1988**, 65, 567-570.
- [4] Narayana, B.; Vijaya Raj, K. K.; Ashalatha, B. V.; Suchetha Kumari, N. Synthesis of some new substituted triazolo [4,3-a] [1,4] benzodiazepine derivatives as potent anticonvulsants. *Eur. J. Med. Chem.* **2006**, 41, 417-422.
- [5] Hussenether, T.; Hübner, H.; Gmeiner P.; Troschutz R. Clozapine derived 2,3-dihydro-1H-1,4 and 1,5-benzodiazepines with D₄ receptor selectivity: synthesis and biological testing. *Bioorg. Med. Chem.* **2004**, 12, 2625-2637.
- [6] Roma, G.; Grossi, G. C.; Di Braccio, M.; Ghia, M.; Mattioli, F. 1,5-Benzodiazepines. IX. A new route to substituted 4H-[1,2,4] triazolo [4,3-a] [1,5] benzodiazepin-5-amines with analgesic and/or anti-inflammatory activity. *Eur. J. Med. Chem.* **1991**, 26, 489-496.
- [7] Kavali, J. R.; Badami, B. V. 1,5-Benzodiazepine derivatives of 3-arylsydnonones: Synthesis and antimicrobial activity of 3-aryl-4-[2'-aryl-2',4',6',7'-tetrahydro-(1'H)-1',5'- benzodiazepine-4'-yl] sydnones. *Il Farmaco*, **2000**, 55, 406-409.
- [8] Di Braccio, M.; Grossi, G. C.; Roma, G.; Vargiu, L.; Mura, M.; Marongiu, M. E. 1,5-Benzodiazepines. Part XII. Synthesis and biological evaluation of tricyclic and tetracyclic 1,5-benzodiazepines derivatives as nevirapine analogues. *Eur. J. Med. Chem.*, **2001**, 36, 935-949.
- [9] Kumar, R.; Joshi, Y. C. Synthesis spectral studies and biological activity of 3H-1,5-benzodiazepine derivatives. *Arkivoc* **2007**, (XIII), 142-149.
- [10] Kamal, A.; Shankaraiah, N.; Prabhakar, S.; Reddy, C. R.; Markandeya, N.; Laxma K.; Devaiah, X. Solid-phase synthesis of new pyrrolobenzodiazepine-chalcone conjugates: DNA-binding affinity and anticancer activity. *Bioorg. Med. Chem. Lett.*, **2008**, 18, 2434-2439.
- [11] Kosychova, L.; Stumbreviciute, Z.; Pleckaitiene, L.; Janciene, R.; Puodziunaite, B.D. Synthesis of substituted 5,6-Dihydro-4H-[1,2,4]triazolo-[4,3-a][1,5] benzodiazepines. *Chem. Heterocycl. Comp.* **2004**, 40, 811-815.
- [12] Janciene, R.; Vektariene, A.; Stumbreviciute, Z.; Kochychova, L.; Klimavicius, A. A. Straightforward synthesis of novel 4H-thiazolo [3,2-d][1,5] benzodiazepine derivatives. *Heteroatom Chem.*, **2004**, 15, 363-368.
- [13] Kosychova, L.; Pleckaitiene, L.; Staniulyte, Z.; Janciene, R.; Palaima, A. and Puodziunaite, B. D. A convenient synthesis of novel substituted imidazo[1,2-a][1,5] benzodiazepine derivatives. *Arkivoc* **2006**, (xiii), 158-164.
- [14] Che, X.; Zheng, L.; Dang, Q.; Bai, X. Synthesis of Novel tricyclic pyrimidine-fused 5,6-dihydrobenzodiazepines via a Pictet-Spengler-like cyclization. *Tetrahedron*, **2006**, 62, 2563-2568.
- [15] Terret, N. K.; Bojanic, D.; Merson, J. R.; Stephenson, P. T. Imidazo [2'-3'/6,5] dipyrindo[3,2-b/2',3'-e]-1,4-diazepines: non-nucleoside HIV-1 reverse transcriptase inhibitors with greater enzyme affinity than nevirapine. *Bioorg. Med. Chem. Lett.* **1992**, 2, 1745-1750.
- [16] (a) Sanghetti, J. N.; Kokare, N. D.; Shinde, D. B. Sulfanic acid catalyzed solvent-free synthesis of 1,5-Benzodiazepine derivatives. *Chinese Chem. Lett.* **2007**, 18, 1305-1308.
- [17] Kusanur Raviraj A.; Ghate, M.; Kulkarni M. V. Synthesis of spiro[indolo-1,5-benzodiazepines] from 3-acetyl coumarins for use as possible antianxiety agents. *J. Chem. Sci.*, **2004**, 116, 265-270.

- [18] Kidwai, M.; Ruby, M. K. and Venkataraman R. A facile synthesis of substituted benzodiazepines using solid support. *Chem. Heterocyclic Comp.* **2004**, 40, 631-634.
- [19] Sanna, P.; Carta, A.; Nikoogar, M. E. Synthesis and Antitubercular activity of 3-aryl substituted-2-(1H(2H) benzotriazol-1(2)-yl) acrylonitriles. *Eur. J. Med. Chem.* **2000**, 35, 535-543.
- [20] Kopanska, K.; Najda, A.; Zebrowska, J.; Chomicz, L.; Piekarczyk, J.; Myjak, P. and Bretner, M. Synthesis and activity of 1H-benzimidazole and 1H-benzotriazole derivatives as inhibitors of *Acanthamoeba castellanii*. *Bioorg. Med. Chem.*, **2004**, 12, 2617-2624.
- [21] Boido, A.; Vazzana, I.; Mattioli, F.; Sparatore, F. Antiinflammatory and antinociceptive activities of some benzotriazolylalkanoic acid. *Il Farmaco*, **2003**, 58, 33-44.
- [22] Biagi, B.; Calderone, V.; Giorgi, I.; Livi, O.; Scartoni, V.; Baragatti, B.; Martinotti, E. Some structural changes on triazolyl-Benzotriazoles and triazolyl-Benzimidezolones as potential potassium channel activators. *Il Farmaco*, **2001**, 56, 841-849.
- [23] Katritzky, A. R.; Lan, X.; Yang, J. Z.; Denisko, O. V. Properties and synthetic utility of N- substituted benzotriazoles. *Chem. Rev.* **1998**, 98, 409-548.
- [24] Nedjar-Kolli, B.; Hamdi, M.; Pecher, J. Synthesis of (3,4-c) Dihydropyrano-1,5-Dihydrobenzodiazepines. *Synth. Comm.* **1990**, 20, 1579-1587.
- [25] Greenhill, John V. Aromatic enamionones: Part 1. Ultraviolet absorption of N-aryl enamionones derived from dimedone. *J. Chem. Soc. Perkin Trans. 1*, **1976**, 2207-2210.
- [26] Rashed, N.; Sayed, M.; Elashry, E. S. H. Dibenzo[b,e][1,4]diazepin-1-ones, synthesis and derivatization, *J. Chinese Chem. Soc.* **1993**, 40, 189-194.
- [27] Cortes, E.C.; Banos, M.A.; De Cortes, O.G.M. Efficient synthesis and spectral determination of 11-[(o-; m-; and p-substituted)-phenyl]-8-chloro-3,3-dimethyl-2,3,4,5,10,11-hexahydro-1H-dibenzo[b,e] [1,4] diazepin-1-ones. *J. Het. Chem.* **2004**, 41, 277-280.
- [28] Kalyanam, N.; Likhate, M. A. Synthesis of dibenzo[b,e][1,4]diazepines. *Indian J. Chem. B* **1988**, 27, 754-755.
- [29] Fodili, M.; Amari, M.; Nedjar-Kolli, B.; Robert, A.; Baudy-Floch, M.; Legrel, P. An efficient Synthesis of New 2-pyronyl-1,5-benzodiazepine derivatives. *Synthesis*, **1999**, 811-814.
- [30] Amari, M.; Nedjar-Kolli, B. Nouvelles voies de synthèses des structures pyranobenzodiazepines et pyranobenzodiazepinones. *J. Soc. Alger. Chim.* **2001**, 11, 77-84.
- [31] Amari, M.; Fodili, M.; Nedjar-Kolli, B.; Hoffmann, P.; Périé, J. Reactivity Studies on 4-Aminopyrones: Access to Benzimidazole and Benzimidazolone derivatives. *J. Heterocyclic. Chem.* **2002**, 39, 811-816.
- [32] Hammal, L.; Bouzroua, S.; André, C.; Nedjar-Kolli, B.; Hoffmann, P. Versatile cyclization of aminoanilino Lactones: Access to benzotriazoles and condensed Benzodiazepine-2-thiones. *Synth. Comm.* **2007**, 37, 501-511.
- [33] Quiroga, J.; Mejia, D.; Insuasty, B.; Abonia, R.; Noguerras, M.; Sanchez, A.; Cobo, J. and Low, J. N. Regioselective synthesis of 4,7,8,9-tetrahydro-2H-pyrazolo [3,4-b]-quinolin-5 (6H)-ones. Mechanism and structural analysis. *Tetrahedron*, **2001**, 57, 6947-6953.
- [34] Martinez, R.; Gustavo A. J.; Ramirez M. T.; Perez A.; Martinez, A. Tetrahydropyrrolo [3,2-c] azepin-4-ones as a new class of cytotoxic compounds. *Bioorg. Med. Chem.*, **2006**, 14, 4007-4016.
- [35] Lam, H. W.; Cooke, P. A.; Pattenden, G.; Bandaranayake, W. M.; Wickramasinghe, W. A. Structure and total synthesis of benzylthiocrellidone, a novel Dimedone-based vinyl sulphide from the sponge *Crella spinulata*. *J. Chem. Soc., Perkin Trans. 1*, **1999**, 847-848.
- [36] Bandaranayake, W. M. and Wickramasinghe, W. A. *AIMS Report*, **1995**, 23, p.27.
- [37] Baraldi, P. G.; El-Kashef, H.; Farghaly, A. R.; Vanelle, P.; Fruttarolo, F. Synthesis of new pyrazolo[4,3-e]-1,2,4-triazolo[1,5-c]pyrimidines and related heterocycles. *Tetrahedron*, **2004**, 60, 5093-5104.
- [38] Ashton, B. W.; Suschitzky, H. The Graebe-Ullmann carbazole synthesis. *J. Chem. Soc.* **1957**, 4559-4562.