

Chemical Composition of Essential Oil of *Lantana camara* L. (Verbenaceae) and Synergistic Effect of the Aminoglycosides Gentamicin and Amikacin

Erlânio O. Sousa¹, Thiago S. Almeida¹, Irwin R. A. Menezes¹, Fabíola F. G. Rodrigues¹, Adriana R. Campos², Sidney G. Lima² and José G. M. da Costa^{1*}

¹Programa de Pós-Graduação em Bioprospecção Molecular, Departamento de Química Biológica, Laboratório de Pesquisa de Produtos Naturais, Universidade Regional do Cariri, Rua Cel. Antônio Luiz 1161, Pimenta, 63105-000 Crato-CE, Brasil.

²Vice-Reitora de Pesquisa e Pós-Graduação, Universidade de Fortaleza, Av. Washington Soares 1321, Edson Queiroz, 60811-905, Fortaleza-CE, Brasil.

³Departamento de Química, Universidade Federal do Piauí, Campus Universitário Ministro Petrônio Portella, 64049-550, Bairro Ininga, Teresina-PI, Brasil

(Received November 20, 2010; Revised July 15, 2011; Accepted July 15, 2011)

Abstract: The leaves of *Lantana camara* L. (Verbenaceae) were subjected to hydrodistillation, and the essential oil extracted was examined with respect to chemical composition, antibacterial and antibiotic modifying activity by gaseous contact. Among the 25 identified components, bicyclogermacrene (26.1%), β -caryophyllene (24.4%), germacrene D (19.2%) and valecene (12.0%) were the main constituents. The essential oil volatile constituents inhibited the growth of *Staphylococcus aureus* and *Pseudomonas aeruginosa* with MIC of 1 and > 1 mg/L, respectively. The activity of the antibiotic amikacin was increased by 65% against *S. aureus* and *P. aeruginosa* after contact with the volatile components.

Keywords: *Lantana camara*; chemical composition; antibacterial and modulatory activities.

1. Introduction

Pseudomonas aeruginosa is an opportunistic affection that usually affects hospitalized or immunocompromised persons. Usually occurs infection of the airways by *P. aeruginosa* occurs commonly in patients with cystic fibrosis but also occurs in patients with other forms of bronchiectasis [1]. In view of the high antibiotic resistance and virulence, the infections associated to *P. aeruginosa* are considered to have difficult management [2].

*Corresponding author: E-Mail: galberto.martins@gmail.com

In Germany, analysis of the sputum from patients with Cystic Fibrosis during a period of 12 months showed the presence of *P. aeruginosa* in 50% of these individuals and *Staphylococcus aureus* in 63.3% are the most prevalent pathogens [3-4]. *S. aureus* has emerged as one of the most important human pathogens, and it can be found in 20-40% of adult superior aerial vias [5].

Lantana is a genus of about 150 species of perennial flowering plants popularly used as antirheumatic, stimulant, antibacterial, biologic control and as ornamental plant [6]. Phytochemical studies of *Lantana* species lead to identification and isolation of terpenoids, flavonoids, phenylethanoid glycosides, furanonaphthoquinones, iridoid glycosides and steroids [6-8].

Lantana camara L. (camará) is a shrub that belongs to Verbenaceae family and it's native from America and Africa. Different parts of the plant, mainly the leaves, have been used in treatment of scratching, stomachache, rheumatism, wound healing, biliary fever, toothache, bronchitis, antiseptic and other affections [9].

L. camara is a rich source of many bioactive molecules and the phytochemical studies have resulted in the isolation of many triterpenes, steroids and flavonoids [10-11]. This plant has been claimed to present activities antiprotozoal [12], antibacterial and antifungal [9,13], antioxidant [14], insecticidal [15], antiviral [16], allelopathic properties [17] among others activities, but there is no previous report regarding to modulatory activity of the essential oil by gaseous contact.

In this work, we report the chemical composition, antibacterial activity and antimicrobial modulatory activity of *L. camara* essential oil from Cariri Cearense, Brazilian Northeast, by the minimal inhibitory concentrations and gaseous contact methods.

2. Materials and Methods

2.1. Plant Material

Leaves of *Lantana camara* L. were collected in April, 2009, from the Small Aromatic and Medicinal Plants Garden of the Natural Products Research Laboratory (LPPN) at University Regional do Cariri (URCA), Crato - Ceara state, Brazil. A voucher specimen (#1662) was deposited in the "Herbário Caririense Dárdaro de Andrade Lima" of Regional University of Cariri, Crato.

2.2 Isolation of the essential oil

Samples of fresh leaves (400 g) were triturated and submitted to hydrodistillation process, in a Clevenger-type apparatus for 2 hours, resulting in essential oil yield of 0.18%. The collected essential oil was subsequently dried by anhydrous sodium sulfate (Na_2SO_4), and stored under refrigeration at $< 4^\circ\text{C}$ until be tested.

2.3 GC and GC/MS analysis

Analysis by GC/MS of the essential oil was carried out on a Shimadzu GC-17 A/ MS QP5050A (GC/MS system) using a DB-5HT fused silica capillary column (30 m x 0.25 mm i.d., 0.25 m film thickness); carrier gas helium, flow rate 1.7 mL/min in split mode. The injection port and detector temperature were 270°C and 290°C , respectively. The column temperature was programmed from 35°C to 180°C at $4^\circ\text{C}/\text{min}$ and then 180°C to 250°C at $10^\circ\text{C}/\text{min}$. Mass spectra were recorded from 30 – 450 *m/z*. injected volume: 1 μL of 5 $\mu\text{g}/\text{mL}$ solution in ethyl acetate. Solvent cut time was 3 min. Individual components were identified by matching their 70 eV mass spectra with those of the spectrometer data base using the Wiley L-built library [18] as well as by visual comparison of the fragmentation pattern with those reported in the literature [19].

The percentage compositions were obtained from electronic integration measurements using flame ionization detection (FID), also set at 250°C . *n*-Alkanes (C_9 - C_{24}) were used as reference points in the calculation of relative retention indices. The concentration of the identified compounds was

computed from the GC peak area without any correction factor. GC analyses were performed on a Hewlett Packard 5890 SERIES II equipped with a flame ionization detector (FID) and a J & W Scientific DB-5 fused silica capillary column (30 m x .25 mm x 0.25 μ m). GC oven temperature and conditions were as described above. Injector and detector temperatures were 270°C and 290°C, respectively. Hydrogen was used as carrier gas, flow rate 1.0 mL/min, split mode (1:10).

2.4 Gaseous contact

The antibacterial activity of *L. camara* essential oil was analyzed by the gaseous contact method (indirect contact) [20]. In this assay, two standard strains (*S. aureus* - ATCC 12692; *P. aeruginosa* - ATCC 15442), were obtained from Fundação Oswaldo Cruz – FIOCRUZ, were used. Petri dishes containing gar Brain Heart Infusion agar (BHI) were inoculated with the strains (24 h; 35 \pm 2 °C). The concentration of each inoculum was confirmed by viable count on Plate Count Agar (PCA). After this, appropriate dilutions (0,5 scale McFarland, 1 x 10⁸ CFU/mL) were plated onto Plate Count Agar (PCA).

An amount of 50 μ g of oil was dissolved in 50 μ L of DMSO (1:1). The assay was performed in triplicate and a dilution series of this essential oil solution was prepared: 50, 25, 12.5 and 6.25 μ g of oil. 100 μ L of each dilution was placed inside the upper part of Petri dishes, in order to promote an interaction between the volatiles constituents of the essential oil and the antibiotics. Petri dishes were incubated at 35 \pm 2 °C (24 h).

The minimal inhibitory concentration (MIC) was defined as the minimal inhibitory dose per unit space required to suppress the growth of microorganism in a closed system. The MIC values were expressed as weight per unit volume (mg/L air), where the solution with 50 μ g equals 1 mg/L air [20].

2.5 Antibiotic modulatory activity evaluation

The antibiotic modifying activity of the gaseous component was determined using the same method and the solution with a total of 50, 25, 12.5 and 6.25 μ g of oil was used. In these plates, antibiotics disks with gentamicin and amikacin were used to determine changes in the inhibition zone diameter of *P. aeruginosa* and *S. aureus*. Plates without the essential oil and with DMSO alone were used as control.

2.6 Statistical analysis

The average inhibition zones obtained were submitted the statistical analysis using Analysis of Variance (ANOVA) followed by the Student-Newman Keuls-test Multiple Comparisons. The results with $p < 0.05$ were considered to be significant.

3. Results and Discussion

Table 1 summarizes the chemical composition and retention indices (RI) found using the Hewlett-Packard Model 5971 GC/MS. The average essential oil yield of the experiments was 0.18 %. Essential oil GC/MS analysis permitted the identification and quantification of twenty-two constituents (100.0 %). Bicyclogermacrene (26.1%), β -caryophyllene (24.4%), germacrene D (19.2%) and valecene (12.0%) were the main constituents identified (Table 1). Previous reports of the essential oil of *L. camara* leaves is constituted mainly by sesquiterpenes, specially, β -caryophyllene, isocaryophyllene, germacrene D and bicyclogermacrene [21-25].

Caryophyllene isomers were present between the main constituents of essential oil *L. camara* from Brazil Northeastern in different daytime [26]. In the evaluation seasonal of the essential oil of *L. camara* collected in Madagascar [27], concentration of caryophyllene has been reported to be consistently high throughout the year, independent of sampling seasons.

Table 1. Chemical constituents of essential oil of the leaves of *L. camara*.

Order	Constituents	RI ^a	RI ^b	(%)
1	α -pinene	937	938	0.1
2	sabinene	969	973	0.2
3	β -pinene	973	978	0.1
4	terpinolene	1082	1084	0.1
5	terpinene-4-ol	1175	1177	0.1
6	α -terpineol	1186	1189	0.1
7	cis-3-hexenyl isovalerate	1216	1217	0.1
8	δ -elemene	1337	1337	0.3
9	α -copaene	1371	1376	1.1
10	β -elemene	1382	1385	1.6
11	β -caryophyllene	1416	1417	24.4
12	γ -elemene	1429	1433	5.4
13	aromadendrene	1431	1439	0.8
14	germacrene-D	1473	1474	19.2
15	bicyclogermacrene	1490	1491	26.1
16	valencene	1497	1496	12.0
17	δ -cadinene	1520	1524	1.2
18	germacrene B	1558	1560	1.2
19	spathulenol	1575	1576	1.3
20	caryophyllene oxide	1582	1581	0.2
21	viridiflorol	1591	1590	4.1
22	δ -cadinol	1635	1636	0.3
Total identified				100

^a relative retention indices experimental: n-alkanes (C₉-C₂₄) were used as reference points in the calculation of relative retention indices. ^b relative retention indices [19].

The results of antibacterial tests by gaseous contact show that *P. aeruginosa* no was susceptible to essential oil volatile constituents, and *S. aureus* was susceptible (MIC 1 mg/L air). Other study showed that *S. aureus* was more susceptible (MIC 0.25 mg/L air) to the volatile constituents of the essential oil of *L. montevidensis* Briq. [28]. In previous reports was verified the antibacterial activity of *L. camara* essential oil against *S. aureus* by direct contact method [23-25], but there is no previous report regarding to antibacterial activity by indirect contact.

In other study essential oil of *L. camara* show antibacterial activity by direct contact against *Arthrobacter protophormiae*, *Micrococcus luteus*, *Rhodococcus rhodochrous* and *S. aureus* with minimal bactericidal concentrations of 50, 25, 12.5 and 200 μ g/mL, respectively [23].

The antibiotic activity of amikacin against *S. aureus* was enhanced in the presence of the essential oil by gaseous contact Table 2. Enhancement of antibacterial activity of amikacin and gentamicin against *P. aeruginosa* by the essential oil was verified too. The amikacin zone inhibition diameter was increased (65%), Table 2.

Table 2 show that more significative synergic effects are associated to an increase of essential oil volatile constituents concentrations, and this is statistically significant ($p < 0.05$) in comparison with controls (antibiotics and DMSO).

One study was observed synergistic effects of gentamicin and amikacin activities against *S. aureus* in the presence of the essential oil constituents of *L. montevidensis* Briq., by gaseous contact method. The amikacin zone inhibition diameter was increased (29%). Enhancement of antibacterial activity of amikacin and gentamicin against *P. aeruginosa* by the essential oil was verified too, it was verified a increasing in 102% of the amikacin activity [28].

In one study oil essential of the leaves of *L. camara* was examined to modulatory activity by microdilution test against two multiresistant strains, *E. coli* from sputum and *S. aureus* from surgical

wound, obtained from clinical material. The synergism of the essential oil on aminoglycosides was verified which showed significant reduction of MICs (1250 to 5 µg/mL) against *E. coli* [22].

In other study was verified the synergistics effects of the extracts etanolic of leaves and roots of *L. montevidensis* Briq. on aminoglycosides activity by microdilution test. The maximum effects were obtained with extract roots on gentamicin activity against multiresistant strains of *E. coli* with MIC reduction (312 to 2 µg/mL) [29].

Table 2. Modification of the antibiotic activity of the volatile constituents of *L. camara* essential oil by gaseous contact on *S. aureus* and *P. aeruginosa*.

Treatment	<i>Staphylococcus aureus</i> (mm ± DP)				<i>Pseudomonas aeruginosa</i> (mm ± DP)			
	Gentamicin	(%)	Amikacin	(%)	Gentamicin	(%)	Amikacin	(%)
Antibiotics	16.3±0.6	-	17.3±0.6	-	14.3±0.6	-	15.0±0.0	-
DMSO	16.7±0.6	-	17.3±0.6	-	14.0±0.0	-	15.3±0.6	-
EOLc 50 µg	16.3±0.6	0	22.3±1.1*	29	17.3±0.6*	21	24.7±0.6*	65
EOLc 25 µg	16.3±0.6	0	21.0±1.1*	21	17.0±0.6*	19	23.3±0.6*	55
EOLc 12.5 µg	16.3±1.1	0	19.3±0.6	11	16.5±0.0	15	20.0±0.0*	33
EOLc 6.25 µg	16.0±0.0	0	18.0±0.0	4	16.0±1.0	12	18.3±0.5*	22

EOLc - Essential Oil of *L. camara*; (%) - Percentages of enhancement on antibiotic activity; *The mean values of inhibition zones (mm ± DP) are statistically significant when compared with controls ($p < 0.05$ – ANOVA followed by the Student-Newman Keuls-test Multiple Comparison). The results are expressed as mean ± DP (n=3).

Essential oils may interact with and affect the plasma membrane, interfering with respiratory chain activity and energy production [30]. The improvement of antibacterial activity against the gram-negative bacteria *P. aeruginosa* demonstrated a significative result, as the gram-positive bacteria are more susceptible to natural products [31].

The mechanism of action of terpenes is not fully understood but is speculated to involve membrane disruption by the lipophilic compounds, with permeability enhancement [32]. This property can facilitate the antimicrobial agents to penetrate into cell, leading to an activity enhancement. That is a plausible explanation for the positive interactions between the sesquiterpenes constituents and the conventional antibiotics [33].

Many plants have shown not only antibacterial properties, but also the ability to interfere with the antibiotic resistance. The sesquiterpenes constituents (guaiazulene, nerolidol (mixture of the *cis* and *trans* isomers) and germacrene D in association with ciprofloxacin, erythromycin, gentamicin and vancomycin demonstrated synergic effect against *E. coli* and *S. aureus* [34].

The data cited in the literature regarding the essential oil interference, by gaseous contact, show relevant and promising results. In one study *Croton zehntneri* essential oil reinforced the gentamicin activity against *P. aeruginosa* increasing the inhibition halo (48.2%) [35].

The results obtained here show that *L. camara* volatile constituents suppress the *S. aureus* growth, pathogenic bacteria of respiratory system and could be a source of metabolites with antibacterial modifying activity to be used as adjuvants to antibiotic therapy against these pathogens. In part, this study can justify the popular use of *L. camara* to treat respiratory affections.

Acknowledgments

The authors would like to acknowledge financial support from CAPES, CNPq and FUNCAP, UFPI for the chromatograms and FIOCRUZ for the microbial strains. CNPq-INCT for excitotoxicity and neuroprotection.

References

- [1] K. A. Miszkiel, A. U. Wells, M. B. Rubens, P. J. Cole and D. M. Hansell (1997). Effects of airway infection by *Pseudomonas aeruginosa*: a computed tomographic study, *Thorax*. **52**, 260-264.
- [2] N. H. Valerius, C. Koch and N. Hoiby (1991). Prevention of chronic *Pseudomonas aeruginosa* colonisation in cystic fibrosis by early treatment, *Lancet*. **338**, 725-726.
- [3] H. D. M. Coutinho, V. S. Falcao-Silva and G. F. Goncalves (2008). Pulmonary bacterial pathogens in cystic fibrosis patients and antibiotic therapy: a tool for the health workers, *Int. Arch. Med*, **1**, 24.
- [4] G. Valenza, D. Tappe, D. Turnwald, M. Frosch, C. Konig and H. Hebestreit (2008). Prevalence and antimicrobial susceptibility of microorganisms isolated from sputa of patients with cystic fibrosis, *J. Cyst. Fibros*, **7**, 123-127.
- [5] E. W. Konemann (2001). Diagnóstico microbiológico. Texto e Atlas. Medsi: Rio de Janeiro, Rio de Janeiro.
- [6] E. L. Ghisalberti (2000). *Lantana camara* L. (Verbenaceae). *Fitoterapia*. **71**, 467-486.
- [7] J. T. Barre, B. F. Bowden, J. C. Coll, J. De Jesus, V. De La Fuente, G. C. Janairo and C. Y. A. Ragasa (1997). Bioactive triterpene from *Lantana camara*, *Phytochemistry*. **45**, 321-324.
- [8] D. K. Verma, S. K. Singh, G. Nath and V. Tripathi (1997). Antimicrobial active triterpenoids from *Lantana* species, *Indian Drugs*. **34**, 390-392.
- [9] M. J. Deena and J. E. Thoppil (2000). Antimicrobial activity of the essential oil of *Lantana*, *Fitoterapia*. **71**, 453-455.
- [10] S. Begum, S. Q. Zehra and B. S. Siddiqui (2008). Two new pentacyclic triterpenoids from *Lantana camara* Linn. *Chem. Pharm. Bull.* **56**, 1317-1320.
- [11] D. K. Verma, S. K. Singh and V. Tripathi (1997). A rare antibacterial flavone glucoside from *Lantana camara*, *Indian Drugs*. **34**, 32-35.
- [12] C. Clarkson, V. J. Maharaj, N. R. Crouch, O. M. Grace, P. Pillay, M. G. Matsabisa, N. Bhagwandin, P. J. Smith and P. I. Folb (2004). *In vitro* antiplasmodial activity of medicinal plants native to or naturalized in South Africa, *J. Ethnopharmacol.* **92**, 177-191.
- [13] O. O. Sonibare and I. Effiong (2008). Antibacterial activity and cytotoxicity of essential oil of *Lantana camara* L. leaves from Nigeria, *African J. Biotechnol.* **7**, 2618-2620.
- [14] J. Benites, C. Moiteiro, G. Miguel, L. Rojo, J. Lopez, F. Venancio, L. Ramalho, S. Feio, S. Dandlen, H. Casanova and I. Torres (2009). Composition and biological activity of the essential oil of Peruvian *Lantana camara*, *J. Chilean Chem. Soc.* **54**, 379-384.
- [15] V. K. Dua, A. C. Pandey and A. P. Dash (2010). Adulticidal activity of essential oil of *Lantana camara* leaves against mosquitoes, *Ind. J. Med. Res.* **131**, 434-439.
- [16] C. Garcia, E. G. Acosta, A. C. Carro, B. M. C. R. Fernandez, R. Bomben, C. B. Duschatzky, M. Perotti, C. Schuff and E. B. Damonte (2010). Virucidal activity and chemical composition of essential oils from aromatic plants of central west Argentina, *Nat. Prod. Commun.* **5**, 1307-1310.
- [17] M. Verdeguer, M. A. Blázquez and H. Boira (2009). Phytotoxic effects of *Lantana camara*, *Eucalyptus camaldulensis* and *Eriocephalus africanus* essential oils in weeds of Mediterranean summer crops, *Biochem. System. Ecol.* **37**, 362-369.
- [18] J. W. Alencar, A. A. Craveiro, F. J. A. Matos and M. I. L. Machado (1900). Kovats indices simulation in essential oils analysis, *Quim. Nova*, **13**, 282-284.
- [19] R. P. Adams (2007). Identification of Essential Oil Components by Gas Chromatography/Mass Spectrometry, 4th Ed. Allured Publishing Corporation. Carol Stream, Illinois.
- [20] S. Inouye, T. Takizawa and H. Yamaguchi (2001). Antibacterial activity of essential oils and their major constituents against respiratory tract pathogens by gaseous contact, *J. Antimicrob. Chemother.* **47**, 565-573.
- [21] R. C. Padalia, R. S. Verma and V. Sundaresan (2010). Volatile constituents of three invasive weeds of Himalayan region, *Rec. Nat. Prod.* **4**, 109-114.
- [22] E. O. Sousa, N. F. Silva, F. F. G. Rodrigues, A. R. Campos, S. G. Lima and J. G. M. Costa (2010). Chemical composition and resistance-modifying effect of the essential oil of *Lantana camara* Linn. *Pharmacog. Mag.* **6**, 79-82.
- [23] N. P. Kurade, V. Jaitak, V. K. Kaul and O. P. Sharma (2010). Chemical composition and antibacterial activity of essential oils of *Lantana camara*, *Ageratum houstonianum* and *Eupatorium adenophorum*, *Pharm. Biol.* **48**, 539-544.

- [24] J. G. Costa, E. O. Sousa, F. F. G. Rodrigues, S. G. Lima and R. Braz-Filho (2009). Chemical composition, evaluation of antibacterial activity and toxicity of the essential oils from *Lantana camara* L. and *Lantana* sp. *Braz. J. Pharmacogn.* **19**, 721-725.
- [25] T. Hernández, M. Canales, J. G. Avila, A. M. García, A. Martínez, J. Caballero, A. Romo de Vivar and R. Lira (2005). Composition and antibacterial activity of essential oil of *Lantana achyranthifolia* Desf. (Verbenaceae), *J. Ethnopharmacol.* **96**, 551-554.
- [26] E. O. Sousa, A. V. Colares, F. F. G. Rodrigues, A. R. Campos, S. G. Lima and J. G. Costa (2010). Effect of collection time on essential oil composition of *Lantana camara* Linn (Verbenaceae) growing in Brazil Northeastern, *Rec. Nat. Prod.* **4**, 31-37.
- [27] J-A. Randrianalijaona, P. A. R. Ramanoelina, J. R. E. Rasoarahona and E. M. Gaydou (2005). Seasonal and chemotype influences on the chemical composition of *Lantana camara* L. Essential oils from Madagascar, *Anal. Chim. Acta.* **545**, 46-52.
- [28] E. O. Sousa, F. F. G. Rodrigues, H. D. M. Coutinho, A. R. Campos, S. G. Lima and J. G. M. Costa (2011). Chemical composition and aminoglycosides synergistic effect of *Lantana montevidensis* Briq. (Verbenaceae) essential oil, *Rec. Nat. Prod.* **5**, 60-64.
- [29] E. O. Sousa, T. S. Almeida, F. F. G. Rodrigues, A. R. Campos, S. G. Lima and J. G. M. Costa, (2011). *Lantana montevidensis* Briq improves the aminoglycoside activity against multiresistant *Escherichia coli* and *Staphylococcus aureus*, *Ind. J. Pharm.* **43**, 180-182.
- [30] K. Nicolson, G. Evans and P. W. Toole (1999). Potentiation of methicillin activity against methicillin-resistant *Staphylococcus aureus* by diterpenes, *FEMS Microbiol. Lett.* **179**, 233-239.
- [31] J. G. Silva, I. A. Souza, J. S. Higinio, J. P. Siqueira-Junior, J. V. Pereira and M. S. V. Pereira (2007). Atividade antimicrobiana do extrato de *Anacardium occidentale* Linn em amostras multiresistentes de *Staphylococcus aureus*, *Braz. J. Pharmacogn.* **17**, 572-577.
- [32] M. M. Cowan (1999). Plant products as antimicrobial agents, *Clin. Microbiol. Rev.* **2**, 564-582.
- [33] J. L. Ríos and M. C. Recio (2005). Medicinal plants and antimicrobial activity, *J. Ethnopharmacol.* **100**, 80-84.
- [34] M. Simões, S. Rocha, M. A. Coimbra and M. J. Vieira (2008). Enhancement of *Escherichia coli* and *Staphylococcus aureus* antibiotic susceptibility using sesquiterpenoids, *Med. Chem.* **4**, 616-623.
- [35] F. F. G. Rodrigues, J. G. M. Costa and H. D. M. Coutinho (2009). Synergy effects of the antibiotics gentamicin and the essential oil of *Croton zehntneri*, *Phytomedicine.* **16**, 1052-1055.