Supporting Information

Rec. Nat. Prod. 9:2 (2015) 184-189

Isolation, Structure Determination, In Vivo/Vitro Assay and Docking Study of a Xanthone with antitumor activity from Fungus *Penicillium oxalicum*.

Gang Chen^{1, 2}, Jiao Bai^{1, 2}, Haifeng Wang^{1, 2}, Shuling Zhang³ and Yuehu Pei^{1, 2}*


Table of Contents	Page
S1: The ¹ H NMR spectrum of compound 1	2
S2: The ¹³ C NMR spectrum of compound 1	2
S3: The ESIMS spectrum of compound 1	3
S4: The HSQC spectrum of compound 1	3
S5: The HMBC spectrum of compound 1	4

¹ School of Traditional Chinese Materia Medica, Shenyang Pharmaceutical University, Shenyang 110016, China


² Key Laboratory of Structure-Based Drug Design & Discovery Ministry of Education, Shenyang Pharmaceutical University, Shenyang 110016, China

³ Division of Oncology, Shengjing Hospital of China Medical University, Shenyang, 110004, China

^{*} Corresponding author: E-Mail: peiyuehu@vip.163.com; Phone:086-024-23986485




 $\boldsymbol{S1}$: The 1H NMR spectrum of compound $\boldsymbol{1}$




 $\mathbf{S2}$: The 13 C NMR spectrum of compound $\mathbf{1}$


Direct Mass Spectrometry Analysis



S3: The ESIMS spectrum of compound 1



S4: The HSQC spectrum of compound **1**



S5: The HMBC spectrum of compound **1**