

Supporting Information

Rec. Nat. Prod. 10:6 (2016) 659-700

Determination of the Volatile Organic Compounds in Fourty Five *Salvia* Species by Thermal Desorption-GC-MS Technique

Seda Damla Hatipoglu^{1,2}, Nihal Zorlu¹, Tuncay Dirmenci³,
Ahmet C. Goren¹, Turan Ozturk² and Gulacti Topcu*⁴

¹TÜBİTAK UME, Chemistry Group Laboratories, 41470, Kocaeli, Türkiye,

²Department of Chemistry, Faculty of Science and Letters, Istanbul Technical University,
34469, Istanbul, Türkiye,

³Department of Biology, Balıkesir University, 10145, Balıkesir, Türkiye,

⁴Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Bezmialem Vakıf
University, Istanbul, Türkiye.

Table of Contents	Page
S1: Unidentified compounds	2
S2: 3,5,6,7,8,8a-Hexahydro-4,8a-dimethyl-6-(1-methylethenyl)- 2(1H)naphthalenone	2

S1: Unidentified compounds

RT*	m/z
43.270	127.2 (5), 113.1 (10), 99.1 (20), 85.1 (50), 71.1 (65), 57.0 (100), 42.9 (75), 40.9 (30), 38.9 (5)
44.265	206.9 (10), 127.2 (8), 109.1 (12), 99.1 (20), 85.1 (50), 71.1 (70), 57.0 (100), 42.9 (90), 40.9 (45), 38.9 (10)
44.730	206.9 (20), 177.1 (15), 145.1 (20), 133.1 (30), 110.0 (35), 90.9 (45), 55.0 (50), 42.9 (100), 40.9 (50)
44.818	234.2 (20), 193.2 (20), 164.1 (25), 133.1 (70), 42.9 (100)
45.213	234.2 (20), 216.1 (60), 192.1 (70), 173.2 (50), 149.1 (55), 135.1 (40), 107.06 (40), 90.9 (80), 77.0 (50), 55.0 (60), 42.9 (100), 40.9 (70)

* Retention time

S2: 3,5,6,7,8,8a-hexahydro-4,8a-dimethyl-6-(1-methylethenyl)-2(1H)naphthalenone