

A New Dibenzofuran from the Barks of *Sorbus commixta*

Seong Yeon Choi¹, Birang Jeong¹, Hyeon Seok Jang¹, Jiho Lee¹
Kiwon Ko¹, Hyunha Kim¹, Jua Hong¹, Young Soo Bae² and
Heejung Yang^{*1}

¹College of Pharmacy; Kangwon National University, Chuncheon 24341, Republic of Korea

²College of Forest and Environmental Sciences, Kangwon National University, Chuncheon 24341, Republic of Korea

(Received June 20, 2017; Revised September 13, 2017; Accepted Month September 14, 2017)

Abstract: A new dibenzofuran derivative, 1,2,4-trimethoxydibenzofuran-3,9-diol (**7**), was isolated from the ethyl acetate fraction of *Sorbus commixta* barks, along with six known compounds, lupeol (**1**), betulin (**2**), betulinic acid (**3**), ursolic acid (**4**), β -sitosterol (**5**) and β -pyrufuran (**6**). Their structures were determined by NMR spectroscopic analysis, including of ¹H and ¹³C NMR, ¹H-¹H COSY, HSQC and HMBC spectra data. Cytotoxic activities of seven compounds were evaluated in five cancer cell lines, HEP-2, A549, MCF-7, PC-3 and SKOV-3 at the concentrations ranging from 10 to 100 μ M.

Keywords: *Sorbus commixta*; triterpenes; dibenzofuran; 1,2,4-trimethoxydibenzofuran-3,9-diol, cytotoxicity. © 2017 ACG Publications. All rights reserved.

1. Plant Source

The barks of *S. commixta* was collected from Gamjeong-ri, Dong-Myeon, Chuncheon, Kangwon Province, Korea, in August 2015. It was identified by Prof. Young Soo Bae, College of Forest and Environmental Sciences, Kangwon National University, and has been stored in the Herbarium of College of Pharmacy, Kangwon National University (KNUP-SC-1).

2. Previous Studies

Sorbus commixta Hedl (Rosaceae) is mainly distributed in Korea and China. It has used for the treatment of neuralgia, cough and expectorant in folk remedies [1-3]. Until now, the study on the *S. commixta* has been mainly focused on the berries of *S. commixta*, and their biological activities, anti-inflammatory and antioxidant effect, have been studied [4-6]. The anticancer activity of the bars of *S. commixta* has been previously reported, [7] but whether single compounds have the anti-proliferation activity against cancer cells have not reported yet. Previous research on the plant had reported the isolation of sorbic acid, chalcone glucoside, prunasin, amygdalin and triterpenoids such as lupenone

and lupeol[4,8-9]. Recently, the methanol extract from the barks of *S. commixta* was shown to have antioxidant, anti-inflammatory and the potent radical scavenging activity [4,5].

3. Present Study

In the present study, the ethyl acetate fraction of the barks of *S. commixta* showed the cytotoxic activity against the cancer cells. Further, we isolated seven compounds (1-7) from the barks of *S. commixta* and evaluated their cytotoxic activities against five cancer cell lines, HEp-2, A549, MCF-7, PC-3 and SKOV-3.

Extraction and isolation: The dried barks of *S. commixta* (2.3 kg) were extracted three times with 80% methanol for 3h. The extract was concentrated under reduced pressure to obtained total extract (183.5g), which was then suspended in water and sequentially partitioned with *n*-hexane, EtOAc and *n*-butanol to give four fractions. The EtOAc fraction (46.6 g) was subjected to silica gel C.C eluting with gradient of *n*-hexane/EtOAc and CHCl₃/MeOH to afford ten fractions (Fr. 1 - 10). Among these, Fr. 4, 5 and 6 were used to obtain compounds 1-7.

Fr. 4 and 5 were subjected to normal phase silica gel C.C eluting with *n*-hexane/EtOAc (5:1 to 0:100) to give four and six sub-fraction, respectively (Fr. 4.1 - 4.4 and Fr 5.1 – 5.6). The sub-fraction Fr. 4.3 was re-crystallized with MeOH to yield compound 1 (5.8 mg). The sub-fraction Fr.5.6 was purified by C₁₈ HPLC (MeOH/H₂O, 3:2) to yield compound 5 (4.4 mg). Fr. 6 was subjected to reverse phase silica gel C.C eluting with MeOH/H₂O (9.5:0.5, isocratic) to give four sub-fractions (Fr. 6.1 - 6.4). The sub-fraction Fr. 6.1 was purified by C₁₈ HPLC (MeOH/H₂O, 3.75:1.25) to yield compound 6 (3.1 mg) and compound 7 (5.4 mg). Compound 3 (4.0mg) was isolated from the sub-fraction Fr. 6.2 by recrystallization in MeOH. The residual solution of the sub-fraction Fr. 6.2 was purified by C₁₈ HPLC (MeOH/H₂O, 3:2) to yield compound 2 (1.9 mg). Compound 4 (1.6 mg) was isolated the sub-fraction Fr. 6.4 by C₁₈ HPLC (MeOH/H₂O, 3.5:1.5).

Seven compounds were isolated from the EtOAc fraction using various chromatographic techniques. Compounds were identified as lupeol (1)[10], betulin (2)[11], betulinic acid (3)[11], ursolic acid (4) [12], β -sitosterol (5)[10], β -pyrufuran (6)[13] and 1,2,4-trimethoxydibenzofuran-3,9-diol (7), respectively. The compound 7 isolated for the first time from nature. Their structures were determined by NMR spectroscopic analysis, including ¹H and ¹³C NMR, ¹H-¹H COSY, HSQC, and HMBC spectra data, which were acquired in the Central Laboratory of Kangwon National University (Figure 1).

1,2,4-trimethoxydibenzofuran-3,9-diol (7) : yellowish syrup; UV (MeOH): λ_{max} (log ϵ), 301 (3.97), 280 (3.88), 285 (3.83); HRESIMS m/z 289.0708 [M-H]⁻ (calcd. for m/z 289.0712 for C₁₅H₁₄O₆); ¹H and ¹³C NMR data see Table 1 and supporting information.

Cytotoxicity: HEp-2 cells were cultured in Dulbecco's modified Eagle's media (DMEM) and MCF-7, A549, PC-3 and SKOV3 cells were cultured in Roswell Park Memorial Institute medium (RPMI) supplemented with 10% fetal bovine serum (FBS), 100 IU/mL penicillin and 100 μ g/mL streptomycin respectively. The cells were maintained at 37°C in a humidified atmosphere at air (95%) and CO₂ (5%). Cell cytotoxicity was measured by MTT assay.

The ¹H NMR spectrum of 7 exhibited three methoxyl signals at δ 4.18 (3H, *s*, OCH₃-1), 4.16 (3H, *s*, OCH₃-4) and 3.97 (3H, *s*, OCH₃-2), and two hydroxyl signals at δ 8.61 (1H, *s*, OH-9) and 5.91 (1H, *s*, OH-3). The ¹³C NMR spectrum displayed twelve down-fielded carbons at around δ 109.4 – 150.9 deduced the presence of two aromatic groups, and three up-fielded carbons at δ 62.4 (OCH₃-4), 61.4 (OCH₃-2) and 61.2 (OCH₃-1). The HSQC spectrum confirmed that three aromatic proton existed through the correlation between δ^H 7.24 (1H, *d*, *J* = 8.1 Hz, H-7) and δ^C 127.8 (C-7), δ^H 7.00 (1H, *d*, *J* = 8.1 Hz, H-6) and δ^C 109.3 (C-8), and δ^H 6.79 (1H, *d*, *J* = 8.1 Hz, H-8) and δ^C 102.9 (C-6). (Table 1).




Figure 1. Structures of compounds **1-7** isolated from the barks of *S. commixta*

Table 1. ¹H NMR (400 MHz) and ¹³C NMR (100 MHz) spectra data of compounds **6** and **7** in CDCl₃ (δ in ppm, *J* in Hz).

Compound	6		7		
	Position	δ_{H}	δ_{C}	δ_{H}	δ_{C}
1					129.4
2					135.8
3					141.1
4					142.7
4a					143.0
5a					155.9
6	7.50 (1H, m)	111.2	7.00 (1H, <i>d</i> , <i>J</i> =8.2)	102.9	111.2
7	7.35 (1H, m)	125.7	7.24 (1H, <i>m</i>)	127.8	125.7
8	7.29 (1H, m)	123.0	6.79 (1H, <i>d</i> , <i>J</i> =8.1)	109.3	123.0
9		121.9		150.9	121.9
9a		123.5		111.7	123.5
9b		114.6		109.4	114.6
OCH ₃ -1	4.05 (3H, <i>s</i>)	60.9	4.18 (3H, <i>s</i>)	61.2	60.9
OCH ₃ -2	4.18 (3H, <i>s</i>)	61.2	3.97 (3H, <i>s</i>)	61.4	61.2
OCH ₃ -4	3.99 (3H, <i>s</i>)	61.5	4.16 (3H, <i>s</i>)	62.4	61.5

The ¹H and ¹³C NMR spectroscopic data of compound **7** showed a quite similar pattern with **6**, β -pyrofuran, except for the signal corresponding to a hydroxyl group. In the HMBC spectra, δ 8.61 (1H, *s*, OH-9) showing the strong hydrogen signal of the hydroxyl moiety was strongly correlated with δ 111.7 (C-9a), which described that the hydroxyl moiety was located at δ 150.9 (C-9), not 102.9 (C-6) (Figure 2 and See Supporting file). Also, the correlation signals of δ 8.61 (1H, *s*, OH-9) with δ 150.9 (C-9) and 109.3 (C-8) proved the presence of the hydroxyl moiety of δ 150.9 (C-9).




Figure 2. Chemical structure and HMBC correlations of compound **7**

Three concentrations (1, 50 and 100 μ M) of seven compounds and the positive control, docetaxel, were measured to confirm cytotoxicity in five cancer cell lines (HEp-2, MCF-7, A549, PC-3 and SKOV-3). Compounds **2**, **3** and **4** showed significant cytotoxicity in HEp-2 and A549 at a concentration of 100 μ M, compared to the positive control, docetaxel (Table 2). But, compounds **1**, **5**, **6** and **7** showed no or weak activities against five cancer cell (data not shown).

Table 2. The cytotoxic effects of compounds **2-4** against five cancer cell lines.

	HEp-2	A549	MCF-7	PC-3	SKOV-3
2	43.5 \pm 2.32 ^{***}	38.0 \pm 3.26 ^{***}	-12.4 \pm 4.74	19.2 \pm 7.68	6.1 \pm 4.92
3	37.2 \pm 3.08 ^{***}	41.0 \pm 4.75 ^{***}	23.3 \pm 22.76	20.4 \pm 5.91	15.8 \pm 16.86
4	84.8 \pm 0.23 ^{***}	80.8 \pm 0.31 ^{***}	80.8 \pm 0.22 ^{***}	76.6 \pm 3.89 ^{***}	77.1 \pm 0.36 ^{***}
docetaxel	83.6 \pm 1.69 ^{***}	64.8 \pm 3.52 ^{***}	-	88.4 \pm 3.73 ^{***}	77.6 \pm 0.77 ^{***}

Analysis of cell cytotoxicity of 100 μ M Compounds 2-4 using the MTT assay in cancer cell lines at 4 hours post-UV at 590 nm irradiation. Columns, mean of three experiments; SD. *, p<0.05; **, p<0.01; ***, p<0.001 for control (untreated) versus treated.

Acknowledgments

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Science, ICT & Future Planning (No. NRF-2015R1C1A1A01053892)

Supporting Information

Supporting Information accompanies this paper on <http://www.acgpubs.org/RNP>

ORCID

Seong Yeon Choi: [0000-0002-3425-6683](https://orcid.org/0000-0002-3425-6683)

Birang Jeong: [0000-0002-1971-7779](https://orcid.org/0000-0002-1971-7779)

Hyeon Seok Jang: [0000-0001-5538-0328](https://orcid.org/0000-0001-5538-0328)

Jiho Lee: [0000-0003-0073-5763](https://orcid.org/0000-0003-0073-5763)

Kiwon Ko: [0000-0001-7986-6357](https://orcid.org/0000-0001-7986-6357)

Hyunha Kim: [0000-0001-7710-0657](https://orcid.org/0000-0001-7710-0657)

Jua Hong: [0000-0002-6816-2482](https://orcid.org/0000-0002-6816-2482)

Young Soo Bae: [0000-0003-1108-9269](https://orcid.org/0000-0003-1108-9269)

Heejung Yang: [0000-0001-5986-9024](https://orcid.org/0000-0001-5986-9024)

References

- [1] L.R. Bhatt, M.S. Bae, B.M. Kim, G. Oh and K.Y. Chai (2009). A chalcone glycoside from the fruits of *Sorbus commixta* Hedl., *Molecules* **14**, 5323-5327.
- [2] J. Bae, G. Sim, J. Kim, H. Pyo, J. Yun and B. Lee (2007). Antioxidative activity of the hydrolytic enzyme treated *Sorbus commixta* Hedl. and its inhibitory effect on matrix metalloproteinase-1 in UV irradiated human dermal fibroblasts, *Arch. Pharm. Res.* **30**, 1116-1123.
- [3] S.C. Hong, J.H. Yoo, M.H. Oh, H. Lee, Y.S. Park M.K. Pyo (2015). Effect of the mixture of *Pueraria lobata* and *Sorbus commixta* extract on the alcohol-induced hangover in rats, *Nat. Prod. Sci.* **21**, 98-103.
- [4] M. Kim, H. Seong and H. Sohn (2016). In-vitro antithrombosis activity of different parts of *Sorbus commixta* from Ulleung island, *J. Life Sci.* **26**, 289-295.
- [5] M.K. Na, S.M. Lee, Y.H. Kim and S.S. Kang (2002). Antioxidant compounds from the stem bark of *Sorbus commixta*, *Nat. Prod. Sci.* **8**, 26-29.

- [6] D.G. Kang, E.J. Sohn, A.S. Lee, J.S. Kim, D.H. Lee and H.S. Lee (2007). Methanol extract of *Sorbus commixta* cortex prevents vascular inflammation in rats with a high fructose-induced metabolic syndrome, *Am. J. Chin. Med.* **35**, 265-277.
- [7] M. Lee, H. Lee, J. Lee, J. Oh, G. Choi and J. Kim (2002). Anticancer effect of *Sorbus commixta* Hedl extracts, *Korean J. Med. Crop Sci.* **10**, 403-408.
- [8] K. Takaishi and H. Kuwajima (1976). Prunasin and amygdalin from *Sorbus commixta*, *Phytochemistry* **15**, 1984.
- [9] S. Lee and C. Lee (1999). Isolation and gas chromatographic analysis of lupenone and lupeol from *Sorbus cortex*, *Anal. Sci. Technol.* **12**, 136-140.
- [10] S. Mouffok, H. Haba, C. Lavaud, C. Long and M. Benkhaled (2012). Chemical constituents of *Centaurea omphalotricha* Coss. & Durieu ex Batt. & Trab. *Rec. Nat. Prod.* **6**, 292-295.
- [11] G. Uddin, B. Siddiqui, M. Alam, A. Sadat, A. Ahmad and A. Uddin (2011). Chemical constituents and phytotoxicity of solvent extracted fractions of stem bark of *Grewia optiva* Drummond ex Burret, *Middle-East. J. Sci. Res.* **8**, 85-91.
- [12] R. A. El-shiekh, D.A. Al-Mahdy, M. S. Hifnawy, T. Tzanova, E. Evain-Bana, S. Philippot, D. Bagrel, E. A. Abdelsattar (2017). Chemical and biological investigation of *Ochrosia elliptica* Labill. cultivated in Egypt, *Rec. Nat. Prod.* **11**, 552-557.
- [13] M.S. Kemp, R.S. Burden and R.T. Loeffler (1983). Isolation, structure determination, and total synthesis of the dibenzofurans α - and β -pyrufuran, new phytoalexins from the wood of *Pyrus communis* L. *J. Chem. Soc. Perkin Trans. 1*, **0**, 2267-2272.

ACG
publications

© 2017 ACG Publications