

Photooxygenation of 4-(5-methylfuran-2-yl)aminocyanopyridine derivatives

Aliye Altundas^{✉*} and Berna Gül Aslan[✉]

Department of Chemistry, Faculty of Science, Gazi University, 06500 Teknikokullar, Ankara-Türkiye

(Received September 02, 2018; Revised September 27, 2018 ; Accepted September 28, 2018)

Abstract: One pot syntheses of six-, seven-membered cycloalkane and pyrane junctured aminocyanopyridines bearing furan unit were successfully prepared from the ketones, 2-((5-methylfuran-2-yl)methylene) malononitrile, and ammonium acetate. Photooxygenation of 2-(4-aminocyanopyridine) substituted 5-methylfurans gave 1,4-dicarbonyl functionalized aminocyanopyridine derivatives in good yield in one single operation.

Keywords: 4-Furyl-2-amino-3-cyanopyridines; singlet oxygen; 4-oxopent-2-enoylnicotinonitrile. ©2018 ACG Publication. All right reserved.

1. Introduction

Furans are useful precursors for a wide variety of building blocks and intermediates in organic synthesis.¹ The oxidation of furan generates the 1,4-enedione unit which can be found in several bioactive natural products.² The 1,4-enedione units are important intermediates in synthetic organic chemistry.³ The oxidation of furan and its derivatives have been explored to access many biologically active compounds.⁴ Usually, the oxidation of furan can be achieved by either chemical oxidation⁵ or singlet oxygen.⁶ Generally the chemical oxidation reagents are usually not compatible with the functional group which are commonly present in substrate. On the other hand, singlet oxygen is not only to be more functional group tolerant but also it is considered as higher yielding and nontoxic oxidizer as well.^{7,8}

Pyridines are important heterocyclic compounds that they serve as the main component to form the back bone of many natural products and bioactive molecules⁹ with the several biological activities such as anti-microbial, cardiotoxic, anti-Parkinson, anti HIV, antitumor, and anti-inflammatory. Multi-functionalized pyridines with amino- and cyano substituents (AmCyPys) also exhibit biological activity. In addition, further functionalization of the cyano and amino groups allow new biologically active compounds.^{10,11}

Aryl ketone moieties are widely found in drugs and medicinal intermediates as they have extraordinary biological and pharmaceutical properties.^{12,13} In addition, 4-acetylpyridine is an anticonvulsant and protects against hypothermic restraint stress-induced gastric ulceration in mice. 3-Acetylpyridine function as a neurotoxin and shows niacin-like activity.¹⁴⁻¹⁷ The syntheses of aminocyanopyridine derivatives have been interest to unearth their potentials such as carbonic anhydrase inhibition^{18,19} and antimicrobial effects.²⁰

Herein, we report the syntheses of furan-fused aminocyanopyridines; 2-amino-4-(5-methylfuran-2-yl)-5,6,7,8-tetrahydroquinoline-3-carbonitrile **3a**, 2-amino-4-(5-methylfuran-2-yl)-6,7,8,9-tetrahydro-5H-cyclohepta[b]pyridine-3-carbonitrile (**3b**) and 2-amino-4-(5-methylfuran-2-yl)-7,8-

* Corresponding author: E-Mail: aaltundas@gazi.edu.tr

dihydro-5H-pyrano[4,3-b]pyridine-3-carbonitrile (**3c**) and their photooxygenation to construct the α,β -unsaturated-1,4-enedione unit as the tunable functional group in addition to the present aminocyano functionality from simple starting material.

2. Experimental

2.1. Materials and apparatus

In the study all of commercial reagents and chromatography solvents were used as obtained if otherwise was not stated. Analytical thin layer chromatography (TLC) was carried out on Merck silica gel 60 F254. Fluka Silica gel 60 (0.063-0.2 mm) was employed for column chromatography. NMR spectra were recorded by means of a Varian 400 MHz NMR instrument (^1H NMR at 400 MHz, ^{13}C NMR at 100 MHz) or a Bruker 400 MHz instrument (^1H NMR at 400 MHz, ^{13}C NMR at 100 MHz). ^1H NMR data are stated as follows: chemical shift (δ ppm), multiplicity (s=singlet, d=doublet, dd=doublet of doublets, ddd=doublet of doublets of doublets, brs=broad singlet, t=triplet, q=quartet, quint=quintet, m=multiplet or otherwise stated), integration, coupling constant (J, Hz). ^{13}C NMR data are presented with reference to chemical shift (δ ppm).

2.2. Methods

2.2.1. General method for preparation of **2**

α -Cyano- β -aryl crotonitriles **2** were prepared starting from 5-methylfuran-2-carbaldehyde in accordance with the procedure described in the literature.^{11, 18}

2.2.2. General method for preparation of **3a-c**

Compound **2** (5 mmol) was suspended in toluene (10 mL), and then ammonium acetate (7.5 mmol) and cycloalkanone derivatives (5 mmol) (cyclohexanone, cycloheptanone, tetrahydro-4H-pyran-4-one) were added to the mixture. The flask was fitted with a reflux condenser and a water separator, and the mixture was refluxed for 6 h. Then, the solvent was evaporated, and the mixture was re-dissolved in chloroform (150 mL) and washed with water (2x50 mL). The organic phase was dried with MgSO_4 , filtered off and recrystallized from ethyl acetate. All aminocyanopyridines (**3a-c**) were prepared by using this procedure¹⁸⁻²⁰ (Scheme 1).

2-Amino-4-(5-methylfuran-2-yl)-7,8-dihydro-5H-pyrano[4,3-b]pyridine-3-carbonitrile (3c): Yield 72%; m.p.: 183-185 °C; IR (cm^{-1}) 3433-3307 (NH_2), 3161 (CH Aryl), 2965 (CH Alkyl), 2203 (CN). ^1H -NMR (DMSO-d_6 , δ , ppm), 2.39 (s, 3H, CH_3), 2.75 (t, $J= 5.8$ Hz, 2H, $-\text{CH}_2-$), 3.93 (t, $J= 5.8$ Hz, 2H, CH_2), 4.61 (s, 2H, CH_2), 6.66 (s, 2H, NH_2), 6.36 (d, $J= 3.6$ Hz, 1H, CH_{furan}), 6.93 (d, $J= 3.6$ Hz, 1H, CH_{furan}). ^{13}C (DMSO-d_6 , δ , ppm), 15.1, 33.7, 65.8, 67.5, 85.2, 110.2, 117.5, 118.8, 139.8, 146.7, 155.7, 156.1, 159.9, 161.0.

2.2.3. General procedure for photooxygenation of 2-amino-3-cyanopyridine derivatives **3a-c** with TPP

The solution of **3a-c** (1 equiv) in DCM (10 mL) was illuminated at 0 °C by using a halogen lamp (500 Watt) in the presence of TPP (0.4 mol%) while oxygen was being bubbled in the solution. The reaction was kept at the same temperature during the process. The reaction was monitored with TLC. When the starting material consumed, the reaction was treated with Me_2S (10 equiv) at 0 °C and allowed to warm to RT. Solvent was removed in vacuo. The residue was purified by radial chromatography or column chromatography (EtOAc/hexanes).

(E,Z)-2-amino-4-(4-oxopent-2-enoyl)-5,6,7,8-tetrahydroquinoline-3-carbonitrile (4a): Yield 88%; IR (cm^{-1}) 3412-3314 (NH_2), 3184 (CH Aryl), 2823 (CH Alkyl), 2218 (CN), 1766 and 1714 (C=O). ^1H -NMR (CDCl_3 , δ , ppm) 1.71-1.80 (m, 4H), 2.39 (s, 3H, CH_3), 2.72 (t, $J= 10.9$ Hz, 2H, $-\text{CH}_2-$), 2.86 (t, $J= 10.9$

Hz, 2H, CH₂), 5.27 (s, 2H, NH₂), 6.55; 6.56 (d, $J_E = 5.5$ Hz, 1H, CH_{olefin}), 6.57; 6.60 (d, $J_Z = 12.0$ Hz, 1H, CH_{olefin}), 6.63; 6.66 (d, $J_Z = 12.0$ Hz, 1H, CH_{olefin}), 6.70; 6.72 (d, $J_E = 5.5$ Hz, 1H, CH_{olefin}).


(*Z*)-2-amino-4-(4-oxopent-2-enoyl)-6,7,8,9-tetrahydro-5H-cyclohepta[b]pyridine-3-carbonitrile (**4b**): Yield: 85%; IR (cm⁻¹) 3472-3359 (NH₂), 3184 (CH Aryl), 2859 (CH Alkyl), 2218 (CN), 1766 and 1709 (C=O). ¹H-NMR (CDCl₃, δ, ppm) 1.60-1.86 (m, 6H), 2.37 (s, 3H, CH₃), 2.71 (t, $J = 10.9$ Hz, 2H, -CH₂-), 2.91 (t, $J = 10.9$ Hz, 2H, CH₂), 5.12 (s, 2H, NH₂), 6.55; 6.56 (d, $J_E = 5.5$ Hz, 1H, CH_{olefin}), 6.57; 6.60 (d, $J_Z = 12.0$ Hz, 1H, CH_{olefin}), 6.55 (d, $J_Z = 5.49$ Hz, 1H, CH_{olefin}), 6.60 (d, $J_E = 5.49$ Hz, 1H, CH_{olefin}). ¹³C-NMR (CDCl₃, ppm) 13.7, 26.2, 28.0, 29.2, 31.7, 39.5, 112.6, 113.1, 126.8, 127.2, 134.7, 135.0, 135.1, 138.2, 158.9, 169.3.

(*E,Z*)-2-amino-4-(4-oxopent-2-enoyl)-5,8-dihydro-6H-pyrano[3,4-b]pyridine-3-carbonitrile (**4c**): Yield: 89%; IR (cm⁻¹) 3415-3340 (NH₂), 3189 (CH Aryl), 2854 (CH Alkyl), 2218 (CN), 1728 and 1709 (C=O). ¹H-NMR (CDCl₃, δ, ppm) 2.39 (s, 3H, CH₃), 2.64 (s, 3H, CH₃), 2.91 (m, 4H, CH₂), 4.01 (t, $J = 6.0$ Hz, 2H, CH₂), 4.05 (t, $J = 6.0$ Hz, 2H, CH₂), 4.78 (b, 2H, -CH₂-), 4.90 (b, 2H, -CH₂-), 5.20 and 5.29 (s, 2H, NH₂), 6.57 (d, $J_Z = 5.49$ Hz, 1H, CH_{olefin}), 6.67; (d, $J_E = 11.8$ Hz, 1H, CH_{olefin}), 6.76 (d, $J_E = 11.8$ Hz, 1H, CH_{olefin}), 6.78 (d, $J_Z = 5.49$ Hz, 1H, CH_{olefin}).

3. Result and Discussion


We commenced with the methylfurfural **1** and malononitrile to furnish **2**. In our previous study^{11,18-20}, three aminocyanopyridines; 2-amino-4-(5-methylfuran-2-yl)-5,6,7,8-tetrahydroquinoline-3-carbonitrile, 2-amino-4-(5-methylfuran-2-yl)-6,7,8,9-tetrahydro-5H-cyclohepta[b]pyridine-3-carbonitrile and 2-amino-4-(5-methylfuran-2-yl)-7,8-dihydro-5H-pyrano[4,3-b]pyridine-3-carbonitrile (**3a-c**) were obtained in 75, 73 and 72% yields respectively as seen Scheme 1.

When starting materials in hand, we began to establish the best photooxygenation condition for every substrate by using different sensitizer, solvent and reaction temperature. For this purposes, **3a** was subjected to the RB (Rose Bengal) sensitized photooxygenation by carrying out in MeOH at -78 °C using a 500 W halogen lamp while oxygen was passed through the solution. After the consumption of aminocyanopyridines judged by TLC, the reaction was quenched with excess Me₂S to cleavage the peroxide bond. ¹H NMR of crude mixtures showed unpromising result. For this reason, the photooxygenation of **3a** was repeated under the TPP-sensitized condition in DCM at 0 °C by monitoring the reaction with TLC. After the reaction was completed, the crude material was purified on silica gel column to obtain **4a** 88% conversion in yield. Encouraging this pleasing result, **3b** and **3c** were also oxidized under the similar photooxidation condition to give **4b** and **4c**, respectively (Scheme 2).



Scheme 1. Synthesis of 4-furyl-2-amino-3-cyanopyridine derivatives (**3** cyclic ketones: cyclohexanone; cycloheptanone; tetrahydro-4H-pyran-4-one).

Asta et al.²¹ synthesized of (*E*)/(*Z*) 3-alkene-2,4-diones from 2,5-dialkylfurans via laccase-catalyzed ring opening. They reported ³ $J_{3-H,4-H} = 11.8$ Hz for (*Z*)-3-octene-2,5-dione and ³ $J_{3-H,4-H} = 11.9$ Hz for (*Z*)-3-nonene-2,5-dione. In our study, the coupling constant of vinylic protons in enedione units of **4b** was ³ $J = 5.5$ Hz, which is also indicating its *Z* configuration.



Scheme 2. Photooxygenation of 4-furyl-2-amino-3-cyanopyridine derivatives

4. Conclusion

Photooxygenation of aminocyanopyridines **3a-c** gave the corresponding 2-amino-3-cyanopyridine bearing α,β -unsaturated-1,4-dicarbonyl unit that otherwise can not be constructed in one single operation. To the best of our knowledge, these compounds are not known in the literature and can be taken for further structural manipulation to prepare a new series of structurally polyfunctional compounds for drug discovery research.

Acknowledgements

The authors would like to thank R. Altundas (Gebze Technical University) for helpful discussions and his critical linguistic editing of the manuscript.

ORCID

Aliye Altundas: [0000-0003-4616-4513](https://orcid.org/0000-0003-4616-4513)

Berna Gül Aslan: [0000-0002-7510-9540](https://orcid.org/0000-0002-7510-9540)

References

- [1] Dean F. M. Recent advances in furan chemistry. Part I. *Adv. Heterocycl. Chem.* **1982**, *30*,167-238.
- [2] Yu J.Q.; Corey E. J. A mild catalytic, and highly selective method for the oxidation of α,β -enones to 1,4-enediones. *J. Am. Chem. Soc.* **2003**, *125*, 3232-3233.

- [3] Kobayashi Y.; Nakano M.; Kumar G. B.; Kishihira K. Efficient conditions for conversion of 2-substituted furans into 4-oxygenated 2-enoic acids and its application to synthesis of (+)-aspicilin, (+)-patulolide A, and (-)-pyrenophorin. *J. Org. Chem.* **1998**, *63*, 7505-7515.
- [4] Kobayashi, Y.; Kumar, G. B.; Kurachi, T.; Acharya, H. P.; Yamazaki, T.; Kitazume, T. Furan ring oxidation strategy for the synthesis of macrosphelides A and B. *J. Org. Chem.* **2011**, *66*, 2001-2018.
- [5] Clive, D. L. J.; Ou, L. Conversion of furans into γ -hydroxybutenolides: Use of sodium chlorite. *J. Org. Chem.* **2005**, *70*, 3318-3320.
- [6] Graziano, M. L.; Lesce, M. R.; Scarpati, R. Photosensitized oxidation of furans. Part 1. Synthesis and properties of furan endo-peroxides. *J. Chem. Soc. Perkin Trans I*, **1980**, 1955-1959.
- [7] Montagnon, T.; Noutsias, D.; Alexopoulou, I.; Tofi, M.; Vassilikogiannakis, G. Green oxidations of furans-initiated by molecular oxygen-that give key natural product motifs. *Org. Biomol. Chem.* **2011**, *9*, 2031-2039.
- [8] Kazancioglu, E. A.; Kazancioglu, M. Z.; Fistikçi, M.; Secen, H.; Altundas, R. Photooxygenation of azidoalkyl furans: Catalyst-free triazole and new endoperoxide rearrangement. *Org. Lett.* **2013**, *15*, 4790-4793.
- [9] Onitsuka, S.; Nishino, H.; Kurosawa, K. Photooxygenation of 3-acetyl-5-aryl-2-methylfurans via endoperoxide intermediate and the following reactions. *Tetrahedron* **2001**, *57*, 6003-6009.
- [10] Baldwin, J. J.; Engelhardt, E. L.; Hirschmann, R.; Ponticello, G. S.; Atkinson, J. G.; Wasson, B. K.; Sweet, C. S.; Scriabine, A. Heterocyclic analogs of the antihypertensive. β -adrenergic blocking agent (S)-2-[3-(tert-butylamino)-2-hydroxypropoxy]-3-cyanopyridine. *J. Med. Chem.* **1980**, *23*, 65-70.
- [11] Manna, F.; Chimenti, F.; Bolasco, A.; Bizzarri, B.; Filippelli, W.; Filippelli, A.; Gagliardi, L. Anti-inflammatory, analgesic and antipyretic 4,6-disubstituted 3-cyano-2-aminopyridines. *Eur. J. Med. Chem.* **1999**, *34*, 245-254.
- [12] Kambe, S.; Saito, K.; Sakurai, A.; Midorikawa, H. A simple method for the preparation of 2-amino-4-aryl-3-cyanopyridines by the condensation of malononitrile with aromatic aldehydes and alkyl ketones in the presence of ammonium acetate. *Synthesis* **1980**, *5*, 366-368.
- [13] Bandarugattu, V. B.; Naveen, K. K.; Waro, N.; Satoko, H.; Naseem, A.; Petar, M. M. Experimental and theoretical advances in functional understanding of flavonoids as anti-tumor agents. *Anticancer Agents Med. Chem.*, **2013**, *13*, 307-332.
- [14] Yang, Y.; Wang, L.; Chen, Y.; Dai, Y.; Sun, Z. One-pot synthesis of α,α -disubstituted aryl-1-ethanones via the Wittig-Horner reaction. *Phosphorus, Sulfur and Silicon Relat. Elem.* **2018**, *193*, 121-126.
- [15] Uwai, K.; Konno, N.; Kitamura, S.; Ohta, S.; Takeshita, M. Purification and characterization of rat liver enzyme catalyzing stereoselective reduction of acetylpyridines. *Chirality*, **2005**, *17*, 494-500.
- [16] Hall, A.; Bose, R.; Glavin, G.; Pinsky, C. Effects of 4-acetylpyridine on stress ulcer formation in mice. *Neurosci. Lett.* **1988**, *85*, 110-112.
- [17] Wullner, U.; Weller, M.; Groscurth, P.; Loschmann, P. A.; Schulz, J. B.; Muller, I.; Klockgether, T. Evidence for an active type of cell death with ultrastructural features distinct from apoptosis: The effects of 3-acetylpyridine neurotoxicity. *Neuroscience* **1997**, *81*, 721-734.
- [18] Altundas, A.; Gul, B.; Cankaya, M.; Atasever, A.; Gulcin, I. Synthesis of 2-amino-3-cyanopyridine derivatives and investigation of their carbonic anhydrase inhibition effects. *J. of Biochem. Mol. Toxicol.* **2017**, *12*, 1-8.
- [19] Ayvaz, S.; Cankaya, M.; Atasever, A.; Altuntas, A. 2-Amino-3-cyanopyridine derivatives as carbonic anhydrase inhibitors. *J. Enzyme Inhibit. Med. Chem.* **2013**, *2*, 305-310.
- [20] Altundas, A.; Ayvaz, S.; Logoglu, E. Synthesis and evaluation of a series of aminocyanopyridines as antimicrobial agents. *Med. Chem. Res.* **2011**, *1*, 1-8.
- [21] Asta, C.; Conrad, J.; Mika, S.; Beifuss U. Laccase-catalyzed stereoselective oxidative ring opening of 2,5-dialkylfurans into 2-ene-1,4-diones using air as an oxidant. *Green Chem.*, **2011**, *13*, 3066-3069.