Supporting Information

Org. Commun. 17:1 (2024) 1-7

Protection of aldehydes, hydroxy compounds with acetic anhydride in presence of sulfonic acid functionalized ionic liquid

Reddi Mohan Naidu Kalla^{1,2}, Raju Chakali ³ and Chamarthi Naga Raju⁴

¹School of Chemical Engineering, Yeungnam University, Geyongsan 38541, Republic of South Korea ²Department of Science and Humanities, Sri Venkateswara Engineering College, Karakambadi Road, Tirupati, Andhra Pradesh 517507, India

³Department of Electronics and Communication Engineering, Sri Venkateswara Engineering College, Karakambadi Road, Tirupati, Andhra Pradesh 517507, India

⁴Department of Chemistry, S. V. University, Tirupati-517501, Andhra Pradesh, India

| Table of Contents | Page |
|---|------|
| Figure S1: ¹ H NMR spectrum of compound 1 | 2 |
| Figure S2: ¹³ C NMR spectrum of compound 1 | 2 |
| Figure S3: ¹ H NMR spectrum of compound 5 | 3 |
| Figure S4: ¹³ C NMR spectrum of compound 5 | 3 |
| Figure S5: ¹ H NMR spectrum of compound 6 | 4 |
| Figure S6: ¹³ C NMR spectrum of compound 6 | 4 |
| Figure S7: ¹ H NMR spectrum of compound 15 | 5 |
| Figure S8: ¹³ C NMR spectrum of compound 15 | 5 |
| Figure S9: ¹ H NMR spectrum of compound 17 | 6 |
| Figure S10: ¹³ C NMR spectrum of compound 17 | 6 |





Figure S2 : ¹³C NMR spectrum of compound 1








Figure S4 : ¹³C NMR spectrum of compound 5


© 2024 ACG Publications. All rights reserved.



© 2024 ACG Publications. All rights reserved.



© 2024 ACG Publications. All rights reserved.



© 2024 ACG Publications. All rights reserved.