

Org. Commun. 11:3 (2018) 168-172

organic communications

# Synthesis of (E)-10-hydroxy-2-decenoic acid ethyl ester via a one-pot tandem oxidation-Wittig process

Jacob T. Heppell <sup>10\*1</sup>, Wah Chin Boon <sup>10\*1</sup> and Jasim M. A. Al-Rawi <sup>10\*1</sup>

(Received June 29, 2018; Revised August 20, 2018; Accepted September 01, 2018)

**Abstract:** The synthesis of (E)-10-hydroxy-2-decenoic acid ethyl ester (10H2DA-EE) (3) was achieved via a one-pot, tandem oxidation-Wittig process from commercially available 1,8-octanediol (1), Wittig reagent (Ph<sub>3</sub>P=CHCO<sub>2</sub>Et) (2) and activated MnO<sub>2</sub>. Subsequent hydrolysis with sodium hydroxide gave (E)-10-hydroxy-2-decenoic acid (10H2DA) (4).

**Keywords:** One-pot tandem oxidation-Wittig process; (E)-10-hydroxy-2-decenoic acid ethyl ester; (E)-10-hydroxy-2-decenoic. ©2018 ACG Publication. All right reserved.

#### 1. Introduction

(E)-10-hydroxy-2-decenoic acid (10H2DA) is a biologically active fatty acid found in the royal jelly of honey bees that possesses a range of noteworthy pharmacological effects including anti-tumour [1], anti-bacterial [2], neurogenesis [3-4], angiogenesis inhibition [5], and oestrogen receptor modulation [6]. The synthesis of 10H2DA has been reported by several synthetic routes [7-13].

## 2. Background

In many cases, the  $\alpha$ , $\beta$ -unsaturated carbonyl component is formed through the Wittig or Horner-Wittig olefination reactions with 8-(hydroxy or acetyloxy)octanal to give the ester form of 10H2DA, which after hydrolysis yields the free acid (10H2DA). Synthesis of the 8-(hydroxy or acetyloxy)octanal precursor is a vital component of the whole process and involves a multistep procedure starting from compounds including 1,8-octanediol [8], cyclooctanone [7, 13], oleic acid [10], and 6-chlorohexanol [12]. In this work, the one-pot synthesis of (E)-10-hydroxy-2-decenoic acid ethyl ester (10H2DA-EE) and (E)-10-hydroxy-2-decenoic acid (10H2DA) were achieved.

## 3. Experimental

## 3.1. Chemistry

All chemicals and solvents were purchased from commercial suppliers and were used without further purification. Flash chromatography was performed using commercially available Grace-Resolv or Grace Reveleris HP Silica Flash Cartridges (40 g). Thin layer chromatography was performed using Grace-

<sup>&</sup>lt;sup>1</sup>School of Pharmacy and Applied Science, La Trobe University Bendigo, P.O. Box 199 Bendigo 3550, Australia

<sup>&</sup>lt;sup>2</sup> Steroid Neurobiology Laboratory, Florey Institute of Neuroscience and Mental Health, Parkville, VIC, Australia

<sup>\*</sup> Corresponding author: E-mail: j.al-rawi@latrobe.edu.au Tel: +61354447364; Fax: +61354447476

Reveleris Al-backed TLC Plates (UV254) and were visualised with UV 254 nm light and potassium permanganate stain (0.75 g KMnO<sub>4</sub>, 5.0 g K<sub>2</sub>CO<sub>3</sub>, 0.6 mLof 10% (w/w) NaOH, 100 mL water). Infrared spectra were recorded on an Agilent Technologies Cary 630 FT-IR spectrometer. <sup>1</sup>H NMR, <sup>13</sup>C NMR, <sup>1</sup>H, <sup>1</sup>H COSY and <sup>1</sup>H-<sup>13</sup>C HSQC spectra were recorded on a Bruker Avance 300 NMR spectrometer at 300 MHz for <sup>1</sup>H (T=300K) and 75 MHz for <sup>13</sup>C respectively. All <sup>1</sup>H and <sup>13</sup>C NMR spectral results were recorded as chemical shifts (δ). Chemical shifts recorded in CDCl<sub>3</sub> are relative to the internal TMS (0 ppm) for <sup>1</sup>H spectra and solvent peak (77.1 ppm) for <sup>13</sup>C spectra. <sup>1</sup>H NMR multiplicities are expressed as singlet (s), broad singlet (bs) doublet (d), doublet of doublets (dd), triplet of doublets (td), triplet (t), doublet of triplets (dt), triplet of triplets (tt), quartet (q), doublet of quartets (dq), quintet(quin), and multiplet (m). Microanalysis was performed by Chemical and Microanalytical Services Pty Ltd in Belmont, Victoria.

## 3.1.1. Synthesis of (E)-10-hydroxy-2-decenoic acid ethyl ester (10H2DA-EE) (3)

A 250 mL two-neck round bottom flask was attached with a condenser and purged under nitrogen for 10 minutes. 1,8-octandiol (2.00 g, 13.68 mmol), (Ethoxycarbonylmethylene)triphenylphosphorane 2 (5.96 g, 17.10 mmol), MnO<sub>2</sub> (11.89 g, 136.77 mmol)(Sigma Aldrich, technical, activated, ≥90%) and dichloromethane (100 mL) were added and allowed to stir under nitrogen bubbling for 10 minutes before refluxing under nitrogen for 48 hours. The MnO<sub>2</sub> was then filtered and washed with dichloromethane (3 x 20 mL). The filtrate evaporated to give an oily material, which was triturated with 1:9 (v:v) ethyl acetate – nhexane (30 mL). The resulting triphenylphosphine oxide precipitate was removed by filtration then washed with additional 1:9 (v:v) ethyl acetate – n-hexane (3 x 10 mL). The filtrate was then evaporated to give an oil, which was subjected to flash chromatography (0:100 to 30:70 ethyl acetate:n-hexane) to give (E:Z-90:10)-10-hydroxy-2-decenoic acid ethyl ester (3) (2.25 g, 77%) as a pale yellow oil. IR (ATR): 3358 (broad)(w, OH st), 2928 (m, sp<sup>3</sup> C-H st), 2856 (m, sp<sup>3</sup> C-H st), 1718 (s, C=O st), 1653 (m, C=C st), 1264 (m C-O st), 1179 (s, C-O st), 1037 (s, C-O st), 977 (m, C=C bd) cm<sup>-1</sup>. H NMR (CDCl<sub>3</sub>):  $\delta = 1.29$  (t, 3H, H-12, J = 7.2 Hz, 1.33-1.48 (m, 8H, H-5, H-6, H-7, H-8), 1.56 (quin, 2H, H-9, J = 6.6 Hz), 2.05 (bs, 1H, OH), 2.20  $(dq, 2H, H-4, J=6.9 Hz, J_{H4-H2}=1.5 Hz), 3.62 (t, 2H, H-10, J=6.6 Hz), 4.18 (q, 2H, H-11, J=7.2 Hz), 5.81$ (td, 1H, H-2, J = 15.6 Hz, J = 1.5 Hz), 6.96 (td, 1H, H-3, J = 15.6 Hz, J = 6.9 Hz) ppm. <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 14.3 \text{ (C-3)}, 25.7 \text{ (C-8)}, 28.0 \text{ (C-5)}, 29.1 \text{ (C-6/C-7)}, 29.2 \text{ (C-6/C-7)}, 32.2 \text{ (C-4)}, 32.7 \text{ (C-9)}, 60.2 \text{ (C-11)},$ 62.8 (C-10), 121.3 (C-2), 149.4 (C-3), 166.9 (C-1) ppm. C<sub>12</sub>H<sub>22</sub>O<sub>3</sub> (214.30): calcd. C 67.26, H 10.35; found. C 67.24, H 10.47.

#### 3.1. 2. Synthesis of (E)-10-hydroxy-2-decenoic acid (10H2DA) (4)

(E:Z-90:10)-10-hydroxy-2-decenoic acid ethyl ester **3** (2.00 g, 9.33 mmol), 1 M NaOH (15 mL) and ethanol (5 mL) were heated in a 50 mLround bottom flask at 100 °C for 20 minutes. The reaction mixture was then acidified with conc. HCl (pH ~2) and extracted with diethyl ether (3 x 20 mL). The combined organic layers were washed with Brian water (20 ml), dried over MgSO<sub>4</sub>, filtered and evaporated under reduced pressure. The resulting oil solidified upon cooling to give (E:Z-90:10)-10-hydroxy-2-decenoic acid **4** (1.65 g, 95%) as a white solid. Two recrystallizations from a co-solvent of diethyl ether and n-hexane gave (E:Z >99:1)-10-hydroxy-2-decenoic acid (**4**) (1.23 g, 71%) as a white solid. Mp = 64-66 °C (lit. mp = 64-65 °C [<sup>14</sup>]). IR (ATR): 3423 (m, OH st), 2927 (m, sp³ C-H st), 1693 (s, C=O st), 1653 (s, C=C st), 980 (s, C=C bd) cm⁻¹¹. H NMR (CDCl₃):  $\delta$  = 1.33-1.50 (m, 8H, H-5, H-6, H-7, H-8), 1.57 (quin, 2H, H-9,  $J_{\text{H9-H10}} = J_{\text{H9-H8}} = 6.6 \text{ Hz}$ ), 2.22 (dq, 2H, H-4, J = 6.9 Hz,  $J_{\text{H4-H2}} = 1.2 \text{ Hz}$ ), 3.64 (t, 2H, H-10, J = 6.6 Hz), 5.82 (td, 1H, H-2, J = 15.6 Hz, J = 1.5 Hz), 6.88 (bs, 2H, OH), 7.06 (td, 1H, H-3, J = 15.6 Hz, J = 6.9Hz) ppm. <sup>13</sup>C NMR (75MHz, CDCl₃, T=300K):  $\delta$  = 25.6 (C-5), 27.8 (C-8), 29.1 (C-6/C-7), 29.2 (C-6/C-7), 32.3 (C-4), 32.6 (C-9), 62.9 (C-10), 120.9 (C-2), 152.0 (C-3), 171.5 (C-1) ppm.

#### 4. Present Study

The trans 10H2DA was involved in the treatment of prophylaxis of neurological diseases [4]. In this communication the trans10H2DA was synthesised and used in a previously reported patent by one of the authors (Wah Chin Boon)[4], which was found to be useful in the treatment of prophylaxis Therefore it was important to develope a simple, scalable and high yielding synthetic procedure for 10H2DA to support the in vivo and preclinical future treatment of prophylaxis of neurodevelopmental disorders in mammals.

#### 4. 1. Synthesis of 10H2DA-EE

The need to synthesize the 8-(hydroxy or acetyloxy)octanal intermediate is a limiting step in the preparation of 10H2DA (4). However, the ability to carry out a number of synthetic steps in a one-pot process would circumvent the need to prepare and isolate intermediates. From literature review, an efficient tandem process for the synthesis of terminal-hydroxy-2-alkenoic acid ethyl esters has indeed been reported by via a tandem oxidation-Wittig process involving terminal alkanediols, Wittig reagent (Ph<sub>3</sub>P=CHCO<sub>2</sub>Et, 2) and MnO<sub>2</sub> [15]. However, the synthesis of 10H2DA-EE (3) was not included in the study as the longest chain reported was 8-hydroxy-2-octenoic acid ethyl ester. In this publication, we disclose the extension of this procedure to the synthesis of 10H2DA-EE (3) followed by base-catalysed hydrolysis to give 10H2DA (4). Careful choice of the MnO<sub>2</sub> grade was crucial owing to the varied oxidative capacity among the various grades that are commercially available [15]. We tested three grades of MnO<sub>2</sub> (Table 1) which revealed that the yield of 10H2DA-EE (3) was highest when activated MnO<sub>2</sub> (Sigma Aldrich, technical, ≥90%) was employed (Entries 3, Table 1).

HO

OH

$$OH$$
 $OH$ 
 $OH$ 

**Scheme 1**. Synthesis of 10H2DA-EE (3) by a tandem oxidation-Wittig process.

Table 1. Determining the most suitable grade of MnO<sub>2</sub> for the synthesis of 10H2DA-EE (3).

Entry*	MnO <sub>2</sub> grade <sup>#</sup>	Yield (%)	E:Z ratio
1	ReagentPlus, ≥99%	trace	90:10
2	10μm, ≥90%	7	90:10
3	activated, ≥90%	39	90:10

<sup>\*20</sup>x MnO<sub>2</sub>, 2.4x Wittig (2), DCM, 25°C, 24 hours. \*Purchased from Sigma Aldrich.

**Table 2.** Refining the conditions to optimise the yield of 10H2DA-EE (3).

Entry	Solvent	Wittig (2) Eq	MnO <sub>2</sub> Eq.	Time (h)	Yield (%)	E:Z ratio
4	DCM	1.5	10	24	42	90:10
5	DCM	1.5	10	48	62	90:10
6	DCM	1.25	10	48	58	90:10
7	DCM	1.25	10	72	58	90:10
8	DCM	1.25	6	48	43	90:10
9*	DCM	1.25	10	48	45	95:5
10**	DCM	1.25	10	48	57	92:8
11	THF	1.25	10	72	trace	90:10
12#	Toluene	1.25	10	72	37	90:10
13##	DCM	1.25	10	48	77	90:10

<sup>\*0.3</sup>x Benzoic acid added; \*\*0.1x Benzoic acid added;

The original method regarding terminal-hydroxy-2-alkenoic acid ethyl ester synthesis utilised a very large excess of Wittig reagent (2) (2.4 equ) and MnO<sub>2</sub> (20 x equ) relative to the terminal-alkanediol [15]. Through our investigation, we found such an excess to be superfluous, and often complicated the workup and purification. Reducing the excess of Wittig reagent (2) and MnO<sub>2</sub> (Table 2) was generally found to have only minimal impact on the isolated yield while also facilitating workup and purification. Interesting, a notable yield increase was observed when performing the reaction under nitrogen (Entry 13). However the utilising of THF as the solvent was extremely deleterious as only a trace amount of compound 3 was isolated

<sup>\*</sup>Reaction performed at 60°C; \*\*Reaction mixture purged with nitrogen.

(Entry 11). Similarly, toluene was inferior to DCM (Entry 12), although the effect was not as profound as THF, and with no improvement in the E:Z ratio.

10H2DA-EE (3) synthesized in this investigation (Scheme 1) was a mixture of the trans (E) and cis (Z) isomers with a ratio of 90:10 respectively as shown by <sup>1</sup>H NMR spectroscopy. Since it is (E)-10H2DA (4) that occurs naturally in royal jelly, we sought to increase the proportion of the E isomer. The addition of reagents such as salts and benzoic acid have a profound effect on the reaction rate and/or the E:Z ratio of alkenes formed through Wittig or Wittig-Horner reactions [16-17]. We observed that the addition of a 0.3 x stoichiometric equivalent of benzoic acid (Entry 9) improved the E:Z ratio from 90:10 to 95:5, but also unacceptably reduced the yield from 58% (Entry 6) to 45%. A lower 0.1 x stoichiometric equivalent (Entry 10) gave only a modest improvement in the E:Z ratio from 90:10 to 92:8. Overall, the observed increase in the E:Z ratio did not recompense the reduction in yield.

Additionally, complete separation of the trans (E) and cis (Z) isomers by flash column chromatography was generally cumbersome as multiple runs were required (typically three) to achieve a E:Z ratio of  $\sim$ 99:1. Although employing high performance 20  $\mu$ M-silica flash cartridges in our procedure (Grace Reveleris HP) did provide a noticeable improvement over standard 60  $\mu$ M silica.

## 4.1.2. Hydrolysis of 10H2DA-EE to give the free acid 10H2DA.

The hydrolysis of the (E:Z-90:10)-10H2DA-EE (3) (Scheme 3) with aqueous NaOH followed by acidification with HCl gave (E:Z-90:10)-10H2DA (4) in a 95% yield as a white, waxy solid.

Scheme 2. Synthesis of 10H2DA (4) by hydrolysis of 10H2DA-EE (3) with sodium hydroxide.

Interestingly, flash chromatography was not required for the separation of the E and Z isomers because a single recrystallization from a co-solvent of diethyl ether and hexane yielded (E:Z~96:4)-10H2DA (4) as seen by <sup>1</sup>H NMR; a second recrystallization gave (E:Z>99:1)-10H2DA (4). Therefore, it is much easier and less cumbersome to enrich the E-isomer after formation of the 10H2DA free acid.

The structures of the new products **3** and **4** were confirmed by the analysis of their IR, <sup>1</sup>H NMR and <sup>13</sup>C NMR spectra in addition to the C and H microanalysis. The assignments of the <sup>1</sup>H NMR spectra of compounds **3** and **4** were achieved using <sup>1</sup>H-<sup>1</sup>H correlation spectroscopy (COSY), while the protonated signals of the <sup>13</sup>C NMR spectra were assigned using heteronuclear single quantum coherence spectroscopy (HSQC) and using the assigned proton NMR spectra

#### 5. Conclusion

The one pot synthesis of (E)-10-hydroxy-2-decenoic acid ethyl ester was described. Then the E:Z-90:10)-10H2DA-EE was hydrolysed and purified and gave E:Z-99:1)-10H2DA in 71% yield. It is worth noting that this method has the potential to be a scalable process for the synthesise to produce the biologically important (E)-10-hydroxy-2-decenoic acid.

#### **Supporting Information**

Supporting information accompanies this paper on <a href="http://www.acgpubs.org/organic-communications">htttp://www.acgpubs.org/organic-communications</a>

## ORCID ©

Jacob T. Heppell: <u>0000-0002-0842-5615</u> Wah Chin Boon: <u>0000-0002-8351-9961</u> Jasim M. A. Al-Rawi: <u>0000-0002-1813-0634</u>

#### References

- [1] Townsend, G. F.; Brown, W. H.; Felauer, E. E.; Hazlett, B. Studies on the *in vitro* antitumor activity of fatty acids. *Can. J. Biochem. Physiol.* **1961**, *39*, 1765-1770.
- [2] Garcia, M. C.; Finola, M. S.; Marioli, J. M. Bioassay directed identification of Royal Jelly's active compounds against the growth of bacteria capable of infecting cutaneous wounds. *Adv. Microbiol.* **2013**, *3*, 138-144.
- [3] Hattori, N.; Nomoto, H.; Fukumitsu, H.; Mishima, S.; Furukwa, S. Royal jelly and its unique fatty acid, 10-hydroxy-trans-2-decenoic acid, promote neurogenesis by neural stem/progenitor cells in vitro. *Biomed. Res.* **2007**, *28* (5), 261-266.
- [4] Cheung, S.; Boon, W. C. Methods of treating neurodevelopmental diseases and disorders using trans-10-hydroxy-2-decenoic acid. WO2015184509A1, **2015**.
- [5] Izuta, H.; Chikaraishi, Y.; Shimazawa, M.; Mishima, S.; Hara, H. 10-Hydroxy-2-decenoic acid, a major fatty acid from Royal Jelly, inhibits VEGF-induced angiogenesis in human umbilical vein endothelial cells. *Adv. Access Pub. eCAM* **2009**, *6* (4), 489-494.
- [6] Moutsatsou, P.; Papoutsi, Z.; Kassi, E.; Heldring, N.; Zhao, C.; Tsiapara, A.; Melliou, E.; Chrousos, G. P.; Chinou, I.; Karshikoff, A.; Nilsson, L.; Dahlman-Wright, K. Fatty acids derived from Royal Jelly are modulators of estrogen receptor functions. *Plos One* **2010**, *5* (*12*), e15594.
- [7] Brayer, J.-L.; Frison, N.; Folleas, B. Novel ceramide analogues, processes for preparing same and uses therof. WO/2013/014354.
- [8] Potier, P.; Picot, F.; Brayer, J.-L. Procede de preparation des hydroxy-acides gras insatures et de leurs esters, leur utilisation comme agent anti-collagenase CA2459760A1, 2003.
- [9] Quan, Z.; Zhan, L.; Piao, H.; Song, Z., Synthesis of 10-hydroxy-2-decenoic acid. *Yanb. Yixuey. Xueb.* **1992**, 15 (1), 19-20.
- [10] Kharisov, R. Y.; Botsman, O. V.; Botsman, L. P.; Ishmuratova, N. M.; Ishmuratov, G. Y.; Tolstikov, G. A. Synthesis of 10-hydroxy- and 9-oxo-2E-decenoic acids from oleic acid. *Chem.Nat. Compd.* 2002, 38 (2), 145-148
- [11] Tsuji, J.; Masaoka, K.; Takahashi, T.; Suzuki, A.; Miyaura, N. Synthesis of 10-hydroxy-2-decenoic acid (Royal Jelly Acid) from the butadiene telomer. *Bull. Chem. Soc. Japan* **1977**, *50* (*9*), 2507-2508.
- [12] Fray, G. I.; Jaeger, R. H.; Morgan, E. D.; Robinson, R.; Sloan, A. D. B. Synthesis of trans-10-hydroxydec-2-enoic acid and related compounds. *Tetrahedron* **1961**, *15*, 18-25.
- [13] Frison, N.; Folleas, B.; Brayer, J.-L. Novel method for preparing unsaturated fatty hydroxyacids. WO 2007/045741 A2, 2007.
- [14] Fray, G. I.; Jaeger, R. H.; Robinson, R. Synthesis of trans-10-hydroxydec-2-enoic acid and related compounds. *Tetrahedron* **1961**, *15* (*1-4*), 18-25.
- [15] Phillips, D. J.; Pillinger, K. S.; Li, W.; Taylor, A. E.; Graham, A. E. Diol desymmetrization as an approach to the synthesis of unsymmetrical dienyl diesters. *Tetrahedron* **2007**, *63*, 10528-10533.
- [16] Thompson, S. K.; Heathcock, C. H. Effect of cation, temperature, and solvent on the stereoselectivity of the Horner-Emmons reaction of trimethyl phosphonoacetate with aldehydes. *J. Org. Chem.* **1990**, *55* (*10*), 3386-3388
- [17] Harcken, C.; Martin, S. F. Improved E-selectivity in the Wittig reaction of stabilized ylides with α-alkoxyaldehydes and sugar lactols. *Org. Lett.* **2001**, *3* (22), 3591-3593.

