

Oxidation of some benzyl substituted fused quinazoline derivatives

Dmytro Kravtsov *

*Department of Organic and Bioorganic Chemistry, Zaporizhzhia State Medical University,
Mayakovsky ave., 26, 69035, Zaporizhzhia, Ukraine*

(Received November 04, 2019; Revised December 11, 2019; Accepted December 12, 2019)

Abstract: In this study, syntheses of some benzoyl substituted fused quinazoline derivatives **4a-d**, using Fieser's reagent, are reported. An unexpected product **5a** and **6** were isolated from the reaction mixture. Based on the experimental data, a possible oxidation mechanism of ketone **4a** with chromium trioxide is described. The synthesized compounds were characterized by ^1H , ^{13}C , ^{19}F NMR and LC-MS data.

Keywords: Oxidation; fused quinazoline derivatives; Fieser's reagent; benzyl to benzoyl; mechanism. © 2019 ACG Publications. All rights reserved.

1. Introduction

Oxidation reactions are important tools in organic chemistry. This type of processes is particularly attractive for its versatile possibilities. It allows both to modify the molecule by introducing a new hydrophilic group and to carry out its degradation. Today, practically for each reaction the exact or approximate mechanism is known. Based on this, we can easily predict the reaction product.

Herein, an unusual oxidation of benzyl group is described and tested on other samples.

2. Background

Fused quinazoline derivatives were frequently described as a potential class of biologically active agents.¹⁻⁷ Thus, the chemical modifications of such compound are highly desirable.

Commonly used method for chemical modification is an oxidation reaction. Fieser's reagent⁸ (a mixture of chromium trioxide in acetic acid) is one of the most accessible and effective oxidation system.^{9,10} The aim of this research is to convert benzyl substituted fused quinazolines to the corresponding benzoyl-containing derivatives using chromium trioxide, during which an unexpected reaction took place.

3. Present Study

Chemical Material and Apparatus: Substances **1a-d** were synthesized according to the reported procedures.¹¹⁻¹³ Other chemicals and solvents were obtained from commercially available sources and

* Corresponding author: E-mail: kravtsovssynthesis@gmail.com

were used without additional purification. Melting points were determined in open capillary tubes and uncorrected. The reactions were monitored *via* silica gel coated aluminum TLC plate, with fluorescent indicator F₂₅₄ (Merck). Column chromatography was carried out using Kieselgel Merck 60 (230-400 mesh) as the stationary phase. ¹H, ¹³C and ¹⁹F NMR spectra were recorded at 500 or 400 MHz, 125 MHz and 376 MHz, respectively. Chemical shifts were reported in ppm downfield from tetramethylsilane (¹H, ¹³C) or CFCl₃ (¹⁹F), which were used as internal standards. Mass spectra were recorded with an LC-MS instrument using chemical ionization (CI). LC-MS data were acquired with an Agilent 1200 HPLS system equipped with a DAD/ELSD/LCMS-6120 diode matrix and mass-selective detector, column: Poroshell 120 SBC18, 4.6 mm × 30 mm; eluent A: acetonitrile/water, 99:1, with 0.1% of FA; eluent B: water with 0.1% of FA.

General Procedure: A mixture of the phenylacetic acid (4.4 mmol), CDI (4.8 mmol) and dioxane (20 mL) was stirred at 80 °C for 1 h. Then, the appropriate amine **1a-d** (4.4 mmol) was added and the mixture was refluxed for 3 h., after which H₂O (75 mL) was added. The precipitate was filtered,^a washed thoroughly with H₂O and dried at 60 °C. The crude product was dissolved in AcOH (50 mL) and refluxed for 8 h. After that, the solvent was removed *in vacuo* and MeOH (10 mL) was added. The resulting precipitate was filtered and washed with cold MeOH and dried at 60 °C.^b A solution of CrO₃ (4.2 mmol (~ three-fold excess)) in AcOH (40 mL) was added to a stirred solution of the crude product in AcOH (20 mL) over 30 min at 60-63 °C. The mixture was stirred for 3.5 h. at 60-63 °C, then, poured into a solution of Na₂SO₃ (2.8 mmol) in ice-water (100 mL). The residue was filtered,^c washed thoroughly with H₂O and dried at 60 °C. The resulting material was purified to obtain the products (silica gel column chromatography EtOAc-CHCl₃-PE (6:2:2) **4a^d** and **4b**. Crystallization from DMF-H₂O afforded **4c** and **4d**). (Please see the supporting information file to see the information of ^{a-d})

N-(2-(1*H*-Benzo[d]imidazol-2-yl)phenyl)-2-phenylacetamide (**2a**): Gray solid. M.p. 211 °C. Yield: 88.9%. ¹H NMR (δ, ppm, 400 MHz, DMSO-*d*₆): δ 13.24 (s, 1H), 12.92 (s, 1H), 8.68 (d, *J* = 8.3 Hz, 1H), 8.06 (d, *J* = 7.8 Hz, 1H), 7.71 (d, *J* = 5.6 Hz, 1H), 7.54 (d, *J* = 6.0 Hz, 1H), 7.47 (d, *J* = 7.1 Hz, 2H), 7.39 (t, *J* = 7.8 Hz, 1H), 7.31 (t, *J* = 7.3 Hz, 2H), 7.27 – 7.20 (m, 3H), 7.17 (t, *J* = 7.6 Hz, 1H), 3.81 (s, 2H). ¹³C NMR (δ, ppm, 125 MHz, DMSO-*d*₆): 170.09, 153.03, 151.15, 148.73, 138.76, 135.88, 131.08, 130.87, 129.87, 129.00, 127.74, 127.32, 123.36, 120.37, 116.60, 115.89, 115.45, 110.56, 45.90. MS (CI): *m/z* 328.0 [*M* + H]⁺.

6-Benzylbenzo[4,5]imidazo[1,2-*c*]quinazoline (**3a**): White solid. M.p. 199 °C (lit. 200-202 °C).¹⁴ Yield: 70.0%. ¹H NMR (δ, ppm, 400 MHz, DMSO-*d*₆): 8.58 (d, *J* = 7.9 Hz, 1H), 8.10 (d, *J* = 8.4 Hz, 1H), 7.94 (d, *J* = 8.1 Hz, 1H), 7.87 – 7.77 (m, 2H), 7.71 (t, *J* = 7.3 Hz, 1H), 7.52 (t, *J* = 7.7 Hz, 1H), 7.47 – 7.28 (m, 5H), 7.25 (t, *J* = 6.7 Hz, 1H), 4.92 (s, 2H). ¹³C NMR (δ, ppm, 125 MHz, DMSO-*d*₆): 149.96, 147.74, 144.31, 142.13, 135.54, 132.34, 129.30, 129.13, 129.04, 128.40, 127.93, 127.29, 125.79, 124.18, 123.36, 119.96, 118.26, 115.87, 41.53. MS (CI): *m/z* 310.2 [*M* + H]⁺.

Benzo[4,5]imidazo[1,2-*c*]quinazolin-6-yl(phenyl)methanone (**4a**): Yellow solid. M.p. 183-185 °C (lit. 222-224 °C).¹⁵ Yield: 22.8%. ¹H NMR (δ, ppm, 400 MHz, DMSO-*d*₆): 8.67 (d, *J* = 7.6 Hz, 1H), 8.28 (d, *J* = 7.2 Hz, 2H), 7.99 (d, *J* = 7.9 Hz, 2H), 7.89 (t, *J* = 7.3 Hz, 1H), 7.86-7.79 (m, 2H), 7.65 (t, *J* = 7.6 Hz, 2H), 7.54 (t, *J* = 7.5 Hz, 1H), 7.44 (d, *J* = 8.1 Hz, 1H), 7.36 (t, *J* = 7.3 Hz, 1H). ¹³C NMR (δ, ppm, 125 MHz, DMSO-*d*₆): 188.22, 147.22, 144.30, 143.88, 141.12, 136.28, 134.00, 132.65, 131.49, 129.99, 129.79, 128.77, 128.00, 126.32, 124.44, 123.74, 120.40, 119.57, 113.83. MS (CI): *m/z* 324.0 [*M* + H]⁺.

(2-(4-Fluorophenyl)-[1,2,4]triazolo[1,5-*c*]quinazolin-5-yl)(phenyl)methanone (**4b**): White solid. M.p. 233-236 °C. Yield: 5.8%. ¹H NMR (δ, ppm, 400 MHz, DMSO-*d*₆): 8.63 (d, *J* = 8.2 Hz, 1H), 8.24 (dd, 2H), 8.17 – 8.09 (m, 3H), 8.01 (t, *J* = 7.8 Hz, 1H), 7.95 (t, *J* = 7.5 Hz, 1H), 7.81 (t, *J* = 7.5 Hz, 1H), 7.62 (t, *J* = 7.7 Hz, 2H), 7.36 (t, *J* = 8.7 Hz, 2H). ¹³C NMR (δ, ppm, 125 MHz, DMSO-*d*₆): 186.41, 163.42, 152.72, 143.24, 142.01, 135.07 (d, *J* = 242.9 Hz), 133.25, 131.07, 130.47, 130.06 (d, *J* = 9.3 Hz), 129.69, 129.33, 126.70, 126.45, 124.09, 120.29, 118.54, 116.63 (d, *J* = 22.1 Hz). ¹⁹F NMR (δ, ppm, 376 MHz, DMSO-*d*₆): -109.80. MS (CI): *m/z* 369.2 [*M* + H]⁺.

6-Benzoyl-3-phenyl-2H-[1,2,4]triazino[2,3-*c*]quinazolin-2-one (4c): White needle crystals. M.p. 239–245 °C. Yield: 16.5%. ¹H NMR (δ, ppm, 400 MHz, DMSO-*d*₆): 8.66 (d, *J* = 8.1 Hz, 1H), 8.16 (d, *J* = 7.9 Hz, 2H), 8.07 (t, *J* = 7.6 Hz, 1H), 7.98 (d, *J* = 8.1 Hz, 1H), 7.89 (t, *J* = 7.5 Hz, 1H), 7.84 – 7.73 (m, 3H), 7.62 (t, *J* = 7.7 Hz, 2H), 7.50 (t, *J* = 6.9 Hz, 1H), 7.39 (t, *J* = 7.6 Hz, 2H). ¹³C NMR (δ, ppm, 125 MHz, DMSO-*d*₆): 187.48, 159.88, 151.78, 150.28, 148.04, 143.34, 136.19, 135.82, 134.44, 131.9, 131.76, 130.40, 130.31, 129.76, 129.46, 128.78, 128.56, 126.10, 120.62. MS (CI): *m/z* 379.2 [*M* + *H*]⁺.

6-Benzoyl-3-(4-fluorophenyl)-2H-[1,2,4]triazino[2,3-*c*]quinazolin-2-one (4d): White needle crystals. M.p. 218–223 °C. Yield: 13.7%. ¹H NMR (δ, ppm, 400 MHz, DMSO-*d*₆): 8.66 (d, *J* = 8.0 Hz, 1H), 8.14 (d, *J* = 7.7 Hz, 2H), 8.08 (t, *J* = 7.9 Hz, 1H), 7.98 (d, *J* = 8.1 Hz, 1H), 7.96 – 7.85 (m, 3H), 7.79 (t, *J* = 7.5 Hz, 1H), 7.61 (t, *J* = 7.8 Hz, 2H), 7.24 (t, *J* = 8.7 Hz, 2H). ¹³C NMR (δ, ppm, 125 MHz, DMSO-*d*₆): 187.42, 164.43 (d, *J* = 250.5 Hz), 159.84, 151.78, 149.20, 147.95, 143.30, 136.23, 135.85, 134.40, 132.05 (d, *J* = 9.0 Hz), 130.41, 130.36, 129.77, 128.57, 128.22 (d, *J* = 2.4 Hz), 126.10, 120.56, 115.99 (d, *J* = 21.7 Hz). ¹⁹F NMR (δ, ppm, 376 MHz, DMSO-*d*₆): -107.99. MS (CI): *m/z* 397.0 [*M* + *H*]⁺.

Benzo[4,5]imidazo[1,2-*c*]quinazolin-6(5*H*)-one (5a): Beige solid. M.p. >300 °C. Yield: 33.2%, ¹H NMR (δ, ppm, 500 MHz, DMSO-*d*₆): 11.95 (s, 1H), 8.36 (d, *J* = 7.9 Hz, 1H), 8.31 (d, *J* = 8.0 Hz, 1H), 7.86 (d, *J* = 7.9 Hz, 1H), 7.65 (t, *J* = 7.7 Hz, 1H), 7.55 – 7.29 (m, 4H). ¹³C NMR (δ, ppm, 125 MHz, DMSO-*d*₆): 148.13, 146.84, 143.98, 137.61, 132.73, 131.09, 125.46, 124.89, 124.10, 123.83, 119.58, 116.36, 115.20, 112.25. MS (CI): *m/z* 236.1 [*M* + *H*]⁺.

Benzoic acid (6): White crystals. M.p. 118–120 °C (lit. 122.4 °C).¹⁶ Yield: 8.1%. ¹H NMR (δ, ppm, 400 MHz, DMSO-*d*₆): 12.82 (s, 1H), 7.95 (d, *J* = 7.2 Hz, 2H), 7.61 (t, *J* = 7.4 Hz, 1H), 7.49 (t, *J* = 7.6 Hz, 2H). ¹³C NMR (δ, ppm, 125 MHz, DMSO-*d*₆): 167.77, 133.32, 131.23, 129.72, 129.02. MS (CI): *m/z* 121.2 [*M* – *H*][–].




Figure 1. Synthesis and oxidation of some benzyl substituted fused quinazoline derivatives

Oxidation of benzyl substituted fused quinazoline derivatives produced solely one product. Namely, expected corresponding ketones **4b–d**, so the chemistry described herein is completely in accordance with the literature.^{17,18} However, in the case of oxidation of **3a**, three products were isolated (Figure 1), which were identified as quinazolinone **5a** (main product), ketone **4a** and benzoic acid **6**. Thus, a novel oxidation process has been uncovered.

Moreover, a combination of tertiary amine with vicinal carbonyl group was found to be resistant to the oxidation,¹⁹ assuming the reaction is a Malaprade-like oxidation.²⁰ On the other hand, some tetra-substituted cyclobutanones can undergo a Baeyer-Villiger-like rearrangement to yield the

corresponding γ -lactones by chromic acid in a sulfuric acid-acetic acid mixture.²¹ Nevertheless, it does not explain why ketones **4b-d** were not oxidized to the corresponding quinazolinones.

A plausible explanation could be that quinazolinone **5a** and benzoic acid **6** formations might be due to the less electron-withdrawing effect of imidazole in comparison with 1,2,4-triazole and 1,2,4-triazine heterocycles. It led to the supposedly favorable conditions for coordination of the lone pair of the sp^2 hybridized quinazoline nitrogen atom on the Cr^{VI} atom, resulting in the following degradation of ketone **4a** (Figure 2).




Figure 2. Possible oxidation mechanism of ketone **4a** by CrO_3

5. Conclusion

In this study, an unexpected interaction mechanism between ketone **4a** and chromium trioxide is disclosed. Besides, three new benzoyl substituted fused quinazoline derivatives **4b-d** were synthesized. The study showed that the level of electron-withdrawing effect of the fused heterocycle played a key role in the reaction. While the stronger one terminated the oxidation at the step of ketone formation, the weaker one yielded quinazolinone as a main product.

Acknowledgements

The author expresses his gratitude to Prof. Kovalenko S.I. and Dr. Voskoboinik O.Y. for providing some starting compounds and “Enamine Ltd.” (Kiev, Ukraine) for financial support.

Supporting Information

Supporting information accompanies this paper on <http://www.acgpubs.org/journal/organic-communications>



Dmytro Kravtsov: [0000-0001-6880-2518](https://orcid.org/0000-0001-6880-2518)

References

- [1] Kuarm, B. S.; Reddy, Y. T.; Madhav, J. V.; Crooks, P. A.; Rajitha, B. 3-[Benzimidazo- and 3-[benzothiadiazoleimidazo-(1,2-*c*)quinazolin-5-yl]-2*H*-chromene-2-ones as potent antimicrobial agents. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 524-527.
- [2] Habib, O. M. O.; Hassan H. M.; El-Mekabaty, A. Novel quinazolinone derivatives: synthesis and antimicrobial activity. *Med. Chem. Res.* **2013**, *22*, 507-519.
- [3] El-Sayed R.; Wasfy, A. A.F.; Aly, A. A. Synthesis of novel heterocycles with antimicrobial and surface activity. *J. Het. Chem.* **2005**, *42*, 125-130.
- [4] Rohini, R.; Shanker, K.; Reddy, P. M.; Ho, Y.-P. Ravinder, V. Mono and bis-6-arylbenzimidazo[1,2-*c*]quinazolines: A new class of antimicrobial agents. *Eur. J. Med. Chem.* **2009**, *44*, 3330-3339.
- [5] Rohini, R.; Shanker, K.; Reddy, P. M.; Ravinder, V. Synthesis and antimicrobial activities of a new class of 6-arylbenzimidazo[1,2-*c*]quinazolines. *J. Braz. Chem. Soc.* **2010**, *21*(1), 49-57.
- [6] Vostrova, L. N.; Voronina, T. A.; Karaseva, T. L.; Gernega, S. A.; Ivanov, E. I.; Kirichenko, A. M.; Totrova, M. Yu. Synthesis and anticonvulsant activity of benzimidazo[1,2-*c*]quinazolines. *Pharm. Chem. J.* **1986**, *6*, 404-406.

- [7] Bahekar, R. H.; Rao, A. R. R. Bronchodilation and structure-activity relationship studies on new 6-substituted benzimidazo[1,2-*c*]quinazolines. *Arzneim.-Forsch./Drug Res.* **2000**, *50* (2), 712-716.
- [8] Fieser, L. F.; Szmuszkowicz, J. Synthesis of keto acids by chromic anhydride oxidation of tertiary alcohols. *J. Am. Chem. Soc.* **1948**, *70*, 3352-3355.
- [9] Harmsla, W. M.; Eisenbraun, E. J. Chromic acid oxidation of indans and hydrindacenes. Synthesis of 1,1-dimethylindene. *Org. Prep. Proced. Int.* **1972**, *4* (2), 67-72.
- [10] Smith, L. I.; Webster, I. M. The reaction between quinones and sodium enolates. V. 2,3-Dimethylnaphthoquinone and sodium malonic ester. *J. Am. Chem. Soc.* **1937**, *59*, 662-667.
- [11] Hein, D. W.; Alheim, R. J.; Leavitt, J. J. The use of polyphosphoric acid in the synthesis of 2-aryl- and 2-alkyl-substituted benzimidazoles, benzoxazoles and benzothiazoles. *J. Am. Chem. Soc.* **1957**, *79* (2), 427-429.
- [12] Kholodnyak, S. V.; Schabelnyk, K. P.; Zhernova, G. O.; Sergeieva, T. Yu.; Ivchuk, V. V.; Voskoboynik, O. Yu.; Kovalenko, S. I.; Trzhetsinskii, S. D.; Okovytyy, S. I.; Shishkina, S. V. Hydrolytic cleavage of the pyrimidine ring in 2-aryl-(1,2,4)triazole(1,5-*c*)quinazolines: physico-chemical properties and the hypoglycemic activity of the compounds synthesized. *News of Pharmacy* **2015**, *3*, 9-17.
- [13] Sergeieva, T. Yu.; Voskoboynik, O. Yu.; Okovytyy, S. I.; Kovalenko, S. I.; Shishkina, S. V.; Shishkin, O. V.; Leszczynski, J. Hydrazinolysis of 3-R-[1,2,4]triazino[2,3-*c*]quinazolin-2-ones. Synthetic and theoretical aspects. *J. Phys. Chem. A* **2014**, *118*, 1895-1905.
- [14] Linhua, X.; Tingfang, L.; Lianhui, W.; Xiuling, C. Rh(III)-catalyzed one-pot synthesis of benzimidazoquinazolines via C-H amidation-cyclization of *N*-LG-2-phenylbenzoimidazoles. *J. Org. Chem.* **2019**, *84* (2), 560-567.
- [15] Ambethkar, S.; Kalaiselvi, M.; Ramamoorthy, J.; Padmini, V. I₂-Catalyzed oxidative cross-coupling reaction of methyl ketones and 2-(2-aminophenyl) benzimidazole: facile access to benzimidazo[1,2-*c*]quinazoline. *ACS Omega* **2018**, *3*, 5021-5028.
- [16] O'Neil, M. J.; Smith, A.; Heckelman, P. E. *The Merck index : an encyclopedia of chemicals, drugs, and biological*, Merck, 1996.
- [17] Takashi Kishimoto; Yasuo Yamada; Takao Iwasa; Michihiko Matsuda; Renpei Hatano. Triazole compounds and pest control agent. United States Patent 1999, US 005892048A, Example 7.
- [18] Christoph Braun; Haruko Sawada; Helene Lachaise; Stephane Brunet; Philippe Rinolfi; Christophe Dubost; Pierre-Yves Coqueron; Simon Maechling; Anne-Sophie Rebstock; Ulrike Wachendorff-Neumann. Fungicidal 4-substituted-3-{phenyl[(heterocyclyl-methoxy)imino]methyl}-1,2,4-oxadiazol-5(4*H*)-one derivatives. World Intellectual Property Organization | International Application Published Under the Patent Cooperation Treaty 2013, WO 2013/037717 A1, 140.
- [19] Nicolet, B. H.; Shinn, L. A. The action of periodic acid on α -amino alcohols. *J. Am. Chem. Soc.* **1939**, *61*, 1615.
- [20] Wang, Z. Malaprade Reaction *Comprehensive Organic Name Reactions and Reagents*, 2010.
- [21] Jeanne-Carlier, R.; Bourelle-Wargnier, F. Synthese et oxydation chromique de cyclobutanones tetrasubstituees. *Bull. Chim. Soc. Fr.* **1976**, 1-2 (2), 297-300.