

## Cytotoxic Activity and Phytochemical Constituents of *Macrosolen bidouensis* Tangane & V.S. Dang

Kieu Hung Le<sup>1,2</sup>, Tran Canh Nguyen<sup>1</sup>, The Anh Nguyen<sup>3</sup>, Nu Dan Phuong Nguyen<sup>4</sup>, Thi Hong Tuoi Do<sup>5</sup>, Le Thanh Tuyen Nguyen<sup>6</sup>, Van Son Dang<sup>7</sup>, Trong Duc Tran<sup>8</sup>, Nhat Minh Phan<sup>9</sup>, Trong Dat Bui<sup>9</sup>, Dinh Tri Mai<sup>1,9</sup> and Tan Phat Nguyen<sup>\*1,9</sup>

<sup>1</sup>Graduate University of Science and Technology, Viet Nam Academy of Science and Technology, 18 Hoang Quoc Viet, Cau Giay, Hanoi, Viet Nam

<sup>2</sup>Le Quy Don High School for the Gifted, Ba Ria-Vung Tau Province, Viet Nam

<sup>3</sup>Bui Thi Xuan High School, Ho Chi Minh City, Viet Nam

<sup>4</sup>Tran Phu High School, Binh Phuoc Province, Viet Nam

<sup>5</sup>Department of Pharmacology, Faculty of Pharmacy, University of Medicine and Pharmacy at Ho Chi Minh city, Viet Nam

<sup>6</sup>Saigon Pharmaceutical Sciences & Technologies Center, University of Medicine and Pharmacy at Ho Chi Minh city, Viet Nam

<sup>7</sup>Institute of Tropical Biology, Vietnam Academy of Science and Technology, 85 Tran Quoc Toan, Ho Chi Minh city, Viet Nam

<sup>8</sup>GeneCology Research Centre, University of the Sunshine Coast, 90 Sippy Downs Drive, Sippy Downs, Queensland 4556, Australia

<sup>9</sup>Institute of Chemical Technology, Vietnam Academy of Science and Technology, 01 Mac Dinh Chi, Ho Chi Minh City, Viet Nam

(Received February 08, 2020; Revised September 01, 2020; Accepted September 02, 2020)

**Abstract:** Phytochemical investigations of the whole plants of *Macrosolen bidouensis* Tangane & V.S. Dang, a new species discovered in Viet Nam were conducted and led to the purification of ten compounds, comprising three lupane-type (**1-3**), three friedelane-type (**4-6**), and one glutinane-type (**7**) triterpenoids and three cholestane-type steroids (**8-10**) using various chromatographic methods. Their structures were characterized by HR-ESI-MS, NMR experiments and comparison with previous literature. Compounds **1**, **3** and **5** expressed moderate cytotoxicity against two tested cell lines - MDA-MB-231 and RD (IC<sub>50</sub> ranged from 34.19 to 74.25 μM), whereas compounds **4** and **6** exhibited selective cytotoxic activity (IC<sub>50</sub> ranged from 29.07 to 45.20 μM).

**Keywords:** *Macrosolen bidouensis*; Loranthaceae; cytotoxic activity; triterpenoid; steroid. © 2020 ACG Publications. All rights reserved.

\* Corresponding author: E- Mail: [ntphat@ict.vast.vn](mailto:ntphat@ict.vast.vn) Phone +84 91936-0751.

## 1. Plant Source

The whole plants of *Macrosolen bidoupensis* were collected in Bidoup Nui Ba National Park, Lam Dong Province, Viet Nam in December 2018 and identified by Dr. Van Son Dang, Institute of Tropical Biology, Vietnam Academy of Science and Technology. A voucher specimen (No.VH/PHAT-MB1218) was deposited in Bioactive Compounds Laboratory, Institute of Chemical Technology, Vietnam Academy of Science and Technology.

## 2. Previous Studies

*Macrosolen bidoupensis* is a new species that was discovered in Bidoup, Nui Ba National Park, Lam Dong Province, Viet Nam in 2017 [1]. There is no report on the phytochemical components and biological activities of this species.

## 3. Present Study

The whole-plant powder of *M. bidoupensis* (8 kg) was extracted with 96% EtOH (3 x 30 L) at room temperature. After filtering the residue, the solvents were removed under low pressure and the crude extract was obtained. The crude extract (1000 g) was applied to liquid-liquid extraction procedures and successively partitioned into *n*-hexane, ethyl acetate (EtOAc), and aqueous partition. The *n*-hexane extract (98 g) was subjected to silica gel column chromatography with mobile phase *n*-hexane-EtOAc (100-0 to 0-100, v/v) to yield five fractions (MBH.I-MBH.V). As the same manner, the EtOAc extract (33 g) was separated on silica gel column chromatography with solvent *n*-hexane-EtOAc gradient to obtain five fractions (MBE.I-MBE.V).

MTT assay was performed using the MTT test, as previously described [2]. For the cytotoxicity, the percentage of control (%) was calculated =  $\text{OD}_{570 \text{ sample}} / \text{OD}_{570 \text{ control}} \times 100\%$  measured at the different concentrations (12.5, 25.0, 50.0 and 100.0  $\mu\text{g/mL}$ ) by MTT assay.

The cytotoxic effects of all fractions from the *n*-hexane and ethyl acetate extracts of *Macrosolen bidoupensis* whole plants were evaluated against two human cancer cell lines (MDA-MB-231 and RD) by MTT assay (Table 1). All of the fractions except fraction MBE.V displayed cytotoxicity ( $\text{IC}_{50}$  ranged from  $4.52 \pm 0.33$  to  $68.66 \pm 1.81 \mu\text{g/mL}$ ). Among them, fractions MBH.III and MBE.II revealed strong cytotoxic properties towards breast cancer cell lines (MDA-MB-231) with  $\text{IC}_{50}$  value of  $4.52 \pm 0.33$  and  $4.75 \pm 0.42 \mu\text{g/mL}$ , respectively, while fractions MBH.IV and MBE.III exhibited significant cytotoxic activity against Rhabdomyosarcoma cell lines (RD) with  $\text{IC}_{50}$  value of  $20.37 \pm 1.01$  and  $19.78 \pm 3.12 \mu\text{g/mL}$ , respectively. Therefore, all positive fractions were further investigated for their phytochemical constituents.

Fraction MBH.I (45 g) was eluted with solvent *n*-hexane-EtOAc gradient (100-0 to 0-100, v/v) on silica gel column to give six sub-fractions (MBH.I.1-MBH.I.6). Subfraction MBH.I.2 (7 g) was chromatographed with solvent system *n*-hexane-EtOAc (99:1, v/v) to obtain compounds **4** (900 mg) and **7** (5 mg). Fraction MBH.II (39 g) was eluted with solvent *n*-hexane-EtOAc gradient (100-0 to 80-20, v/v) on silica gel column to give six sub-fractions (MBH.II.1-MBH.II.6). Subfraction MBH.II.2 (10 g) was chromatographed with solvent system *n*-hexane-EtOAc (99:1 to 95:5, v/v) to obtain compounds **1** (1000 mg), **2** (6 mg), **8** (4 mg), **9** and **10** (50 mg). Fraction MBH.III (3 g) was separated using solvent *n*-hexane-EtOAc (98-2, v/v) to give **1** (50 mg) and **3** (6 mg). Fraction MBE.II (8 g) was rechromatographed on silica gel with solvent *n*-hexane-EtOAc gradient (99-1 to 80-20, v/v) to yield **5** (20 mg) and **6** (7 mg).

The HR-ESI-MS and NMR data of the three others were consistent with those reports in the literature for lupeol (**1**) [3,4], betulin (**2**) [5], betulinic acid (**3**) [6], friedelin (**4**), 3 $\alpha$ -friedelanol (**5**), 3 $\beta$ -hydroxyfriedelane-28-oic acid (**6**) [7], 3 $\beta$ -hydroxyglutin-5-ene (**7**) [8], 3 $\beta$ -cholesterol acetate (**8**) [9], and a mixtures of 24 $\beta$ -ethylcholesterol (**9**) and 24 $\beta$ -ethylcholesta-5,22-diene-3 $\beta$ -ol (**10**) [10] (Figure 1). All of those compounds, except **1**, were reported from the genus *Macrosolen* for the first time.

**Table 1.** Cytotoxic activity of all fractions

Cell line	Fraction	IC <sub>50</sub> (µg/mL)
MDA-MB-231	H.I <sup>a</sup>	-
	H.II	31.55 ± 1.51
	H.III	4.52 ± 0.33
	H.IV	15.05 ± 0.99
	H.V	68.66 ± 1.81
	E.I	27.24 ± 1.00
	E.II	4.75 ± 0.42
	E.III	18.39 ± 2.15
	E.IV	42.96 ± 1.46
	E.V	> 100.00
	Paclitaxel	8.38 ± 0.75 (µM)
RD	H.I <sup>a</sup>	-
	H.II	32.30 ± 0.34
	H.III	48.40 ± 3.05
	H.IV	20.37 ± 1.01
	H.V	41.36 ± 0.62
	E.I	41.94 ± 3.56
	E.II	30.12 ± 2.96
	E.III	19.78 ± 3.12
	E.IV	50.43 ± 1.11
	E.V	> 100.00
	Paclitaxel	5.73 ± 0.27(µM)


<sup>a</sup>insoluble

The cytotoxic activities of five triterpenoids (**1**, **3-6**) were examined against two human cancer cell lines (MDA-MB-231 and RD) by MTT assay (Table 2) and compounds **1**, **3** and **5** showed meaningful cytotoxicity against all tested cell lines (IC<sub>50</sub> ranged from 34.19 ± 0.53 to 74.25 ± 0.46 µM), whereas, compound **4** displayed selective cytotoxic property against MDA-MB-231 cells (IC<sub>50</sub> value of 45.20 ± 1.37 µM) and compound **6** exhibited cytotoxic effect against RD cells (IC<sub>50</sub> value of 29.07 ± 3.47 µM). According to those results, we conclude that the friedelane and lupane triterpenoids evinced as the essential components that accounted for the cytotoxic effect of *M. bidoupensis* against two tested human cancer cell lines. However, further clinical examinations are required to determine the molecular mechanisms of cytotoxicity as well as qualitative and quantitative identification of main biological triterpenoid markers (**1** and **4**) from this species.

**Table 2.** Cytotoxic activity of five triterpenoids **1**, **4-6**

Cell line	Compounds	IC <sub>50</sub> (µM)
MDA-MB-231	<b>1</b>	57.78 ± 1.95
	<b>3</b>	34.19 ± 0.53
	<b>4<sup>b</sup></b>	45.20 ± 1.37
	<b>5</b>	51.78 ± 0.73
	<b>6<sup>b</sup></b>	> 50
	Paclitaxel	8.38 ± 0.75
RD	<b>1</b>	74.25 ± 0.46
	<b>3</b>	34.41 ± 1.57
	<b>4<sup>b</sup></b>	> 50
	<b>5</b>	70.95 ± 1.94
	<b>6<sup>b</sup></b>	29.07 ± 3.47
	Paclitaxel	5.73 ± 0.27

<sup>b</sup>the maximal soluble concentration is 50 µM



**Figure 1.** Chemical structures of compounds (1-10)

In conclusion, a systematic chemical investigation of *Macrosolen bidoupensis* Tangane & V.S. Dang, was reported for the first time herein. Phytochemical investigation of the active extracts indicated that the triterpenoids present in this species accounted for their cytotoxic activity.

### Acknowledgments

This research is funded by Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 104.01-2018.48.

### Supporting Information

Supporting Information accompanies this paper on <http://www.acgpubs.org/journal/records-of-natural-products>

**ORCID** Kieu Hung Le: [0000-0003-0964-7751](https://orcid.org/0000-0003-0964-7751)Tran Canh Nguyen: [0000-0002-3711-172X](https://orcid.org/0000-0002-3711-172X)The Anh Nguyen: [0000-0002-4405-9413](https://orcid.org/0000-0002-4405-9413)Nu Dan Phuong Nguyen: [0000-0003-0529-0169](https://orcid.org/0000-0003-0529-0169)Thi Hong Tuoi Do: [0000-0002-5445-0694](https://orcid.org/0000-0002-5445-0694)Nguyen Le Thanh Tuyen: [0000-0002-0073-7171](https://orcid.org/0000-0002-0073-7171)Van Son Dang: [0000-0001-8681-4141](https://orcid.org/0000-0001-8681-4141)Trong Duc Tran: [0000-0003-0095-6259](https://orcid.org/0000-0003-0095-6259)Nhat Minh Phan: [0000-0002-7719-9658](https://orcid.org/0000-0002-7719-9658)Trong Dat Bui: [0000-0001-8892-8396](https://orcid.org/0000-0001-8892-8396)Dinh Tri Mai: [0000-0001-8906-6003](https://orcid.org/0000-0001-8906-6003)Tan Phat Nguyen: [0000-0001-9672-2785](https://orcid.org/0000-0001-9672-2785)**References**

- [1] S. Tagane, V. S. Dang, V. N. Nguyen, T. B. Hoang, N. Komada, J. S. Wai, A. Naiki, H. Nagamasu, H. Toyama and T. Yahara (2017). *Macrosolen bidoupensis* (Loranthaceae), a new species from Bidoup Nui Ba National Park, southern Vietnam, *PhytoKeys*. **80**, 113-120.
- [2] T. P. Nguyen, D. T. Mai, T. H. T. Do and N. M. Phan (2017). Flavonoids with hepatoprotective activity from the leaves of *Cleome chelidonii*, *Nat. Prod. Commun*, **12**, 1061-1063.
- [3] F. P. Beserra, M. Xue, M. G. L. de Azevedo, A. L. Rozza, C. H. Pellizzon and C. J. Jackson (2018). Lupeol, a pentacyclic triterpene, promotes migration, wound closure, and contractile effect *in vitro*: possible involvement of PI3K/Akt and p38/ERK/MAPK Pathways, *Molecules* **23**, 2819.
- [4] W.A. Matebie, W.C. Zhang, S. Zhang and G.B Xie (2019). Triterpenoids from *Acokanthera schimperi* in Ethiopia, *Rec. Nat.Prod.* **13(3)**, 182-188.
- [5] A. Tijjani, I. G. Ndukwe and R. G. Ayo (2012). Isolation and characterization of lup-20(29)-ene-3,28-diol (betulin) from the stem-bark of *Adenium obesum* (Apocynaceae), *Trop. J. Pharm. Res.* **11**, 259-262.
- [6] S. Salimuzzaman, H. Farrukh, B. Sabira and S. S. Bina (1988). Oleanderol, A new pentacyclic triterpene from the leaves of *Nerlum Oleander*, *J. Nat. Prod.* **51**, 229-233.
- [7] G. C. M. Salazar, G. D. F. Silva, L. P. Duarte, S. A. Vieira Filho and I. S. Lula (2000). Two epimeric friedelane triterpenes isolated from *Maytenus truncata* Reiss: <sup>1</sup>H and <sup>13</sup>C chemical shift assignments, *Magn. Reson. Chem.* **38**, 977-980.
- [8] M. M. Fragua, M. Reinaa, J. G. Luis and M. L. Rodríguezb (2003). Rhoiptelenol and rhoiptelenone, two pentacyclic triterpenes from *Sideritis macrostachya*, *Z. Naturforsch.* **58c**, 621-625.
- [9] W. K. Wilson, R. M. Sumpter, J. J. Warren, P. S. Rogers, B. Ruan and G. J. Schroepfer (1996). Analysis of unsaturated C<sub>27</sub> sterols by nuclear magnetic resonance spectroscopy, *J. Lipid Res.* **37**, 1529-1555.
- [10] C. Y. Ragasa, R. F. Galian and C. C. Shen (2014). Chemical constituents of *Annona muricata*, *Pharm. Chem.* **6**, 382-387.

**A C G**  
**publications**

© 2020 ACG Publications