

## Genus *Melaleuca* - A Review on the Phytochemistry and Pharmacological Activities of the Non-Volatile Components

Fatma M. Abdel Bar <sup>1,2\*</sup>

<sup>1</sup>Department of Pharmacognosy, Faculty of Pharmacy, Prince Sattam Bin Abdulaziz University, Al-Kharj 11942, Saudi Arabia

<sup>2</sup>Department of Pharmacognosy, Faculty of Pharmacy, Mansoura University, Mansoura 35516, Egypt

(Received September 09, 2020; Revised October 30, 2020; Accepted November 02, 2020)

**Abstract:** Plants of the genus *Melaleuca* which belong to family Myrtaceae, commonly named "tea trees", are economically important plants. When talking about tea trees, the essential oils are the center of attention, leaving all other phytoconstituents in their shade. Many reviews addressed the composition and pharmacological activities of *Melaleuca alternifolia* L. essential oil as the most common one. To date, there are no detailed reviews summarizing the phytochemical and pharmacological properties of the non-volatile components of members of the genus *Melaleuca*. After distillation of the volatile oil, large amounts of these plants' waste remain untapped. This review indicates that this genus is a rich source of diverse groups of bioactive phytochemicals, including flavonoids, triterpenoids, benzylic phloroglucinol-terpene derivatives, polyphenols, hydrolysable tannins, and other compounds. It also discusses the diverse pharmacological activities reported for plants of this genus.

**Keywords:** *Melaleuca*; Myrtaceae; tea tree; paperbark tree; phytochemistry ; pharmacological activities. © 2021 All rights reserved.

### 1. Introduction

Genus *Melaleuca*, as a member of family Myrtaceae, comprises 230 species. *Melaleuca alternifolia* L. is the most famous species and as other members of this genus, is native to Australia. In Australia, species in the genus *Melaleuca* are commonly named "tea trees" and "paperbark trees". The name "tea tree" originated when the sailors of Captain James Cook, during his journey to Australia in 1770, used this herb as a substitute for tea. Today, a variety of myrtaceous shrubs derived from the genera *Leptospermum*, *Melaleuca*, and *Kunzea* are collectively termed "tea trees", whereas the term "paperbark" arose from the bark that can be pulled off as papery strips [1]. One of these species is *M. leucadendron*, in which the bark peels off in papery layers [2]. The economic significance of plants which belong to *Melaleuca* spp. is due to their essential oils. The essential oil of *M. alternifolia* is the most common one. In addition, honey produced from the blossoms of the genus *Melaleuca* is of great value [1]. Research interest in the genus *Melaleuca* has been focused on the chemical composition and antimicrobial activity of the essential oils of these plants. Meanwhile, phytochemical investigation of

\* Corresponding author: E-Mail: [f.abdelbar@psau.edu.sa](mailto:f.abdelbar@psau.edu.sa); [fatma\\_maar@yahoo.com](mailto:fatma_maar@yahoo.com); Phone:0966-54540-3617

the non-volatile content of the plants of this genus revealed the presence of mainly phenolics and terpenoidal compounds. The plant phenolics include flavonoids, phenylpropanoids, polyphenols, and free phenolic acids [1,3-6]. *Melaleuca* trees are also a rich source of triterpenoids, including lupane, ursane, and oleanane derivatives [7-11]. Another unique group namely benzylic phloroglucinol-terpene hybrids or meroterpenoids has been also reported from different plant parts of *M. leucadendron* [12,13]. Remarkably, the genus *Melaleuca* has shown several pharmacological activities, such as antimicrobial, antiparasitic, antioxidant, anti-inflammatory, acetylcholinesterase inhibition, neuroprotective, molluscicidal, antischistosomal, immunostimulant, anticancer, anti-histaminic, cardiovascular-, hepato-protective, antisecretory, and antiulcerogenic activities. Despite the various reviews that detailed the chemical composition and the pharmacological activities of the essential oils of the genus *Melaleuca*, no comprehensive review addressed the chemistry or the biological properties of the non-volatile components of this economically important genus. Therefore, this review summarized the different phytochemical groups and reported the pharmacological activities of this genus. Also, this review could direct the future research on genus *Melaleuca* towards the discovery of valuable medicinally active natural products especially from the plant waste left after distillation of the essential oil.

## 2. The Botany of Genus *Melaleuca*

Genus *Melaleuca* is classified according to Jones and Luchsinger (1986) as follows [14]:

Kingdom: Plantae (Plants)

Division: Angiosperms (Magnoliophyta, flowering plants)

Class: Magnoliopsida (Dicotyledons)

Subclass: Rosidae

Order: Myrtales

Family: Myrtaceae (Myrtle family)

Subfamily: Leptospermoidae

Genus: *Melaleuca*

### 2.1. Order Myrtales

They are woody and herbaceous plants. The leaves are mostly opposite and often glandular or dotted. The flowers are usually regular, tetra- or pentamerous, with valvate calyx. Stamens tend to be in groups or in bundles, sometimes dimorphous and open by terminal pores. The ovary is inferior, and the ovules are reduced in numbers in higher types. This order is spread over seven families, including Myrtaceae [2].

### 2.2. Family Myrtaceae (Myrtle Family)

The family comprises about 140 genera and 3000 species. It is largely distributed in tropical and subtropical regions and abundant in Australia. They are shrubs to large trees. The leaves are usually opposite, sometimes alternate, simple, leathery, evergreen, entire, exstipulate, and characteristically glandular-dotted. Inflorescences are mostly cymose, sometimes racemose, and rarely solitary. Flowers are epigenous, sometimes perigynous, bisexual, and actinomorphic. The calyx is formed of 4-5 sepals, usually free, sometimes more or less united to form a cap which drops off as the flower opens. The corolla is formed of 4-5 petals. Stamens are indefinite. The ovary is formed of 2-3 united carpels, with one to many (often 2-5) locules, with 2-many ovules, the placentation is axile (rarely parietal), the ovary is commonly inferior, and the style is long and simple with a capitate stigma. Fruits are fleshy berries, rarely drupes, or dry capsules or nuts. Seeds have little or no endosperm. The family is classified into two subfamilies: Myrtoideae and Leptospermoideae [14].

### 2.3. Subfamily Leptospermoidae

The Leptospermoidae subfamily is characterized by spiral to opposite leaves, perigynous or epigynous flowers, and dry dehiscent fruits (capsules) [14,15]. This can be distinct from the Myrtoideae subfamily which has opposite leaves and indehiscent fruits [15].

### 2.4. Genus *Melaleuca*

Plants of the genus *Melaleuca* are shrubs or trees that possess spiral leaves. However, taxonomical studies should not be restricted to morphological characteristics, but molecular systematics based on their DNA comparison are very important in establishing the relationship between species [1].

## 3. Chemistry of the Family Myrtaceae

The Myrtaceae family contains various phytoconstituents, mainly phenolics and terpenoids [16,17]. Phenolics include flavonoids, phenylpropanoids, and polyphenols, such as phloroglucinol derivatives and free phenolic acids. Flavonoids are characterized by the presence of *O*- or *C*-methyl flavonoid, isoflavone, anthocyanin, and chalcone derivatives [18]. Additionally, polyhydroxy alkaloids (e.g. casuarine and fagomine) are of great chemotaxonomic importance to Myrtaceae [19]. The volatile oil of myrtaceous plants comprises various components, including monoterpenes, sesquiterpenes, and phenyl propanoids. Some genera produce very economically important volatile oil, such as clove oil, eucalyptus oil, tea tree oil, cajuput oil, niaouli oil, and pimento oil. Other constituents, such as bitter principles, free amino acids, hydrocarbons, and fatty acids, are also present [1,20].

## 4. Chemistry of the Non-volatile Components of gGenus *Melaleuca*

### 4.1. Triterpenoids


Triterpenoids are important C<sub>30</sub> naturally occurring isoprenoid plant metabolites with important chemistry and pharmacology [21, 22]. In particular, pentacyclic triterpenes have attracted many researchers in different areas of research, mainly as anticancer [7, 23], anti-inflammatory [24, 25], antiviral against HIV [26] and *Chlamydia trachomatis* [27]. The plants of the genus *Melaleuca*, reported with triterpenoidal contents, include *M. ericifolia* SM., *M. alternifolia* L., *M. leucadendron* L., *M. quinquenervia* Cav., *M. cajuputi* Powell., *M. cuticularis* Labill., *M. raphniphylla* Schauer. and *M. viminea* Lindl. [7-10, 28, 29] [4, 11, 30-32].

Triterpenoids from the genus *Melaleuca* possess different skeletons, including ursane (**1.1-1.12**, Table 1 and Figure 1), oleanane (**1.13-1.22**, Table 1 and Figure 2), and lupane (**1.23-1.34**, Table 1 and Figure 3). In addition, a group of miscellaneous triterpenoids has been also reported (**1.35-1.40**, Table 2 and Figure 4). Several triterpenoids from different plant parts, including leaves, heartwood, and fruits, exhibited the presence of esterifying C<sub>6</sub>-C<sub>3</sub> acids at the hydroxyl group at C-3 namely *trans*-*p*-coumaric and *cis/trans*-caffeic acids. Another unique group of triterpenoids from this genus is 28-norlupane which is particularly identified from the leaves of *M. leucadendron* and *M. ericifolia* with a hydroxyl or peroxy substituents at C-17, respectively. The presence of 27,28-dioic acids represented by melaleucic acid **1.29** (3 $\beta$ -hydroxylup-20(29)-en-27,28-dioic acid) and **1.19** (3 $\alpha$ -hydroxy-13(18)-oleanene-27,28-dioic acid), is also another characteristic chemical feature of the triterpenoids of the genus *Melaleuca*.


Although the triterpene nucleus was identified from many plants of the genus *Melaleuca*, there are no sufficient reports describing other non-volatile isoprenoid structures, such as diterpenes and oxygenated sesquiterpenes. Meanwhile, acyclic isoprenoids of the corresponding biogenetic origin of sesquiterpene, diterpene and triterpenes have been detected in this genus, which include (2*E*,6*E*)-farnesol, **2.1** (C15) [11], phytol, **2.2** [11] (C20), and squalene, **2.3** [8, 11] (C30), respectively, Table 3 and Figure 5.

**Table 1.** Pentacyclic triterpenoids from the genus *Melaleuca*.


Compound	Name	Plant/ part	Ref.
<b>I- URSANE-TYPE PENTACYCLIC TRITERPENOIDS</b>			
1.1	Asiatic acid; 2 $\alpha$ ,23-Dihydroxyursolic acid	Leaves of <i>M. ericifolia</i>	[7]
1.2	3 $\beta$ -O-Acetylurs-12-en-28-oic acid	Bark and stem of <i>M. alternifolia</i>	[28]
1.3	3 $\beta$ - <i>cis</i> -Caffeoyloxy-2 $\alpha$ -hydroxyurs-12-en-28-oate	Leaves of <i>M. leucadendron</i>	[10]
1.4	3 $\beta$ - <i>trans</i> -Caffeoyloxy-2 $\alpha$ -hydroxyurs-12-en-28-oate	Leaves of <i>M. leucadendron</i>	[10]
1.5	3- $\beta$ - <i>cis</i> - <i>p</i> -Coumaroyloxy-2 $\alpha$ -hydroxyurs-12-en-28-oic acid	Leaves of <i>M. leucadendron</i>	[10]
1.6	3- $\beta$ - <i>trans</i> - <i>p</i> -Coumaroyloxy-2 $\alpha$ -hydroxyurs-12-en-28-oic acid	Leaves of <i>M. leucadendron</i>	[10]
1.7	3- $\beta$ - <i>cis</i> - <i>p</i> -Coumaroyloxy-2 $\alpha$ -hydroxyursa-12,20(30)-dien-28-oate	Leaves of <i>M. leucadendron</i>	[10]
1.8	2 $\alpha$ ,3 $\beta$ -Dihydroxyurs-12-en-28-oic acid	Heartwood of <i>M. leucadendron</i>	[8]
1.9	2 $\alpha$ -Hydroxyursolic acid	Leaves of <i>M. leucadendron</i>	[10]
1.10	Ursolaldehyde	Leaves of <i>M. leucadendron</i> Fruits of <i>M. leucadendron</i> Leaves of <i>M. ericifolia</i>	[10] [9] [7]
1.11	Ursolic acid	Leaves of <i>M. leucadendron</i> Fruits of <i>M. leucadendron</i> Leaves of <i>M. quinquenervia</i> Seeds of <i>M. cajuputi</i> Leaves of <i>M. ericifolia</i>	[10] [9] [33] [29] [7]
1.12	Uvaol	Stems and leaves of <i>M. leucadendron</i>	[30]
<b>II- OLEANANE-TYPE PENTACYCLIC TRITERPENOIDS</b>			
1.13	3 $\beta$ - <i>trans</i> - <i>p</i> -Coumaroyloxy-2 $\alpha$ ,23-dihydroxyolean-12-en-28-oic acid	Heartwood of <i>M. leucadendron</i>	[8]
1.14	23- <i>trans</i> - <i>p</i> -Coumaroyloxy-2 $\alpha$ ,3 $\beta$ - dihydroxyolean-12-en-28-oic acid	Heartwood of <i>M. leucadendron</i>	[8]
1.15	3 $\beta$ - <i>trans</i> -Caffeoyloxy-2 $\alpha$ ,23-dihydroxyolean-12-en-28-oic acid	Heartwood of <i>M. leucadendron</i>	[8]
1.16	3 $\beta$ - <i>cis</i> -Caffeoyloxy-2 $\alpha$ ,23-dihydroxyolean-12-en-28-oic acid	Heartwood of <i>M. leucadendron</i>	[8]
1.17	2 $\alpha$ ,3 $\beta$ -Dihydroxyolean-12-en-28-oic acid	Heartwood of <i>M. leucadendron</i>	[8]
1.18	3 $\beta$ ,23-Dihydroxyolean-12-en-28-oic acid	Heartwood of <i>M. leucadendron</i>	[8]
1.19	3 $\alpha$ -Hydroxy-13(18)-oleanene-27,28-dioic acid	Heartwood of <i>M. leucadendron</i>	[31]
1.20	Oleanolic acid	Leaf and stem of <i>M. quinquenervia</i>	[32]
1.21	2 $\alpha$ ,3 $\beta$ ,23-Trihydroxyolean-12-en-28-oic acid	Heartwood of <i>M. leucadendron</i>	[8]
1.22	2 $\alpha$ ,3 $\beta$ ,23-Trihydroxyolean-12-en-28-oic acid (arjunolic acid)	Bark and stem of <i>M. alternifolia</i>	[28]
<b>III- LUPANE-TYPE PENTACYCLIC TRITERPENOIDS</b>			
1.23	3 $\beta$ -Acetylup-20(29)-en-28-oic acid	Leaves of <i>M. leucadendron</i>	[11]
1.24	Betulin	Stems and leaves of <i>M. leucadendron</i> Bark and stem of <i>M. alternifolia</i> Leaves of <i>M. ericifolia</i>	[30] [28] [7]
1.25	Betulinolaldehyde	Leaves of <i>M. leucadendron</i> Fruits of <i>M. leucadendron</i> Leaves of <i>M. ericifolia</i>	[11] [9] [7]
1.26	Betulinic acid	Leaves of <i>M. leucadendron</i> Fruits of <i>M. leucadendron</i> Bark and stem of <i>M. alternifolia</i> Fruits of <i>M. leucadendron</i> Seeds of <i>M. cajuputi</i> Leaves of <i>M. ericifolia</i>	[11] [9] [28] [4] [29] [7]
1.27	Betulonic acid; 3-Oxo-lup-20(29)-en-28-oic acid	Leaves of <i>M. leucadendron</i>	[11]
1.28	20 $S$ -17 $\beta$ ,29-Epoxy-28-norlupan-3 $\beta$ -ol	Leaves of <i>M. ericifolia</i>	[7]
1.29	Melaleucic acid; (3 $\beta$ -hydroxylup-20(29)-en-27,28-dioic acid)	Bark of ( <i>M. cuticularis</i> , <i>M. raphniphylla</i> and <i>M. viminea</i> ) Bark and stem of <i>M. alternifolia</i> Heartwood of <i>M. leucadendron</i> Leaves of <i>M. ericifolia</i>	[34] [28] [8] [35]
1.30	28-Norlup-20(29)-ene-3 $\beta$ ,17 $\beta$ -diol	Leaves of <i>M. leucadendron</i> Leaves of <i>M. ericifolia</i>	[11] [7]
1.31	28-Norlup-20(29)-en-3 $\beta$ -hydroxy-17 $\alpha$ -hydroperoxide	Leaves of <i>M. ericifolia</i>	[7]
1.32	28-Norlup-20(29)-en-3 $\beta$ -hydroxy-17 $\beta$ -hydroperoxide	Leaves of <i>M. ericifolia</i>	[7]
1.33	Platanic acid	Leaves of <i>M. leucadendron</i> Leaves of <i>M. ericifolia</i>	[11] [7]
1.34	Pyracrenic acid	Fruits of <i>M. leucadendron</i>	[9],[4]



**Figure 1.** Structures of ursane-type pentacyclic triterpenoids (1.1-1.12) from the genus *Melaleuca*




**Figure 2.** Structures of oleanane-type pentacyclic triterpenoids (1.13-1.22) from the genus *Melaleuca*



**Figure 3.** Structures of lupane-type pentacyclic triterpenoids (1.23-1.34) from the genus *Melaleuca*


**Table 2.** Miscellaneous group of pentacyclic triterpenoids from the genus *Melaleuca*

Compound	Name	Plant/ part	Ref.
1.35	3 $\alpha$ ,27-Dihydroxy-28,20 $\beta$ -taraxastanolide	Heartwood of <i>M. leucadendron</i> .	[31]
1.36	Eupha-7,24-diene-3 $\beta$ ,22 $\beta$ -diol	Heartwood of <i>M. leucadendron</i> .	[31]
1.37	Friedelin	Stems and leaves of <i>M. leucadendron</i>	[30]
1.38	<i>Epi</i> -taraxeryl acetate	Stems of <i>M. leucadendron</i>	[30]
1.39	20-Taraxastene-3 $\alpha$ ,28-diol	Heartwood of <i>M. leucadendron</i>	[31]
1.40	Taraxastenone	Stems of <i>M. leucadendron</i>	[30]

**Figure 4.** Structures of a miscellaneous group of pentacyclic triterpenoids (1.35-1.40) from the genus *Melaleuca*


#### 4.2. Acyclic terpenoids

The reported acyclic terpenoids from the genus *Melaleuca* are few. Only three acyclic terpenoids have been reported, including squalene **2.3** (C<sub>30</sub>), isolated from the heartwood and leaves of *M. leucadendron* [8, 11], phytol **2.2** (C<sub>20</sub>) and (2*E*,6*E*)-farnesol **2.1** (C<sub>15</sub>), isolated from the leaves of *M. leucadendron* Table 3 and Figure 5.

**Figure 5.** Structures of acyclic terpenoids (2.1-2.3) from the genus *Melaleuca*

### 4.3. Phytosterols

Phytosterols are essential structural components in the plasma membrane of plant cells which are found free or conjugated through an ester or a glycosidic linkage. They are important precursors in the synthesis of many valuable bioactive compounds, such as steroidal alkaloids, steroidal saponins, plant-derived ecdysteroids, and brassinosteroids [36]. Four phytosterols have been identified from *Melaleuca* spp., including cholesterol C<sub>27</sub> (**3.1**),  $\beta$ -sitosterol C<sub>29</sub> (**3.2**),  $\beta$ -sitostenone C<sub>29</sub> (**3.3**), and stigmasterol C<sub>29</sub> (**3.4**), in Table 3 and Figure 6.



**Figure 6.** Structures of phytosterols (**3.1-3.4**) from the genus *Melaleuca*

**Table 3.** Acyclic terpenoids and phytosterols from the genus *Melaleuca*

Compound	Name	Plant/ part	Ref.
<b>ACYCLIC TERPENOIDS</b>			
<b>2.1</b>	(2 <i>E</i> ,6 <i>E</i> )-Farnesol (C <sub>15</sub> )	Leaves of <i>M. leucadendron</i>	[11]
<b>2.2</b>	Phytol (C <sub>20</sub> )	Leaves of <i>M. leucadendron</i>	[11]
<b>2.3</b>	Squalene (C <sub>30</sub> )	Heartwood of <i>M. leucadendron</i>	[8]
		Leaves of <i>M. leucadendron</i>	[11]
<b>PYTOSTEROLS</b>			
<b>3.1</b>	Cholesterol	Leaf and stem of <i>M. quinquenervia</i>	[32]
<b>3.2</b>	$\beta$ - Sitosterol	Stems and leaves of <i>M. leucadendron</i>	[30]
<b>3.3</b>	$\beta$ - Sitostenone	Leaf and stem of <i>M. quinquenervia</i>	[32]
		Heartwood of <i>M. leucadendron</i>	[8]
		Leaves of <i>M. cajuputi</i>	[37]
		Heartwood of <i>M. leucadendron</i>	[8]
<b>3.4</b>	Stigmasterol	Leaf and stem of <i>M. quinquenervia</i>	[32]


### 4.4. Flavonoids

Flavonoids are very essential and widely distributed plant C<sub>6</sub>-C<sub>3</sub>-C<sub>6</sub> metabolites. Due to their abundancy, flavonoids have been extensively investigated in literature. They possess diverse pharmacological actions, such as antioxidant, anticancer, enzyme inhibitory, neuroprotective, anti-atherosclerosis, anti-inflammatory, cardiovascular protective, and hepatoprotective effects [38-40]. *Melaleuca* is reported to contain different classes of flavonoids, including flavones, flavonols, and flavans (**4.1-4.47**), Table 4 and Figure 7. The predominant group is flavonols and the presence of methyl ethers was a characteristic feature of the reported flavonoids. Several flavonoid glycosides have been reported with several sugars, mainly as glucose, rhamnose, xylose, and glucuronic acid. However, it is worth noted that unlike other members of Myrtaceae, there are only few reports on C-methylated flavones and flavonols in tea trees represented by 5-hydroxy-7,4'-dimethoxy-6,8-dimethylflavone (**4.5**), isolated from the leaves of *M. cajuputi* [37]. Unlike flavones and flavonols, a class of methylated flavanones has been reported, Figure 8 and Table 4. This class was represented by strobopinin (**4.47**) and kryptostrobin (**4.41**), isolated from the flowers of *M. quinquenervia* [41]. Additionally, Seligmann and Wagner (1981) determined the structure of melanervin (**4.46**), the first naturally occurring flavonoid of the triphenylmethane family. Another class of unusual prenylated flavanones has been also reported from the leaves of *M. leucadenderon* represented by leucadenone A-D (**4.42-4.45**) [42], Table 4 and Figure 8.


**Table 4.** Flavonoids from the genus *Melaleuca*

Compound	Name	Plant/ part	Ref.
<b>I- FLAVONES</b>			
4.1	Apigenin	Leaves of <i>M. huegelii</i>	[3]
4.2	Apigenin-7-methyl ether	Leaves of <i>M. huegelii</i>	[3]
4.3	Apigenin-7,4'-dimethyl ether	Leaves of <i>M. huegelii</i>	[3]
4.4	Herbacetin-3- <i>O</i> -glucuronide	Leaves of <i>M. squarrosa</i>	[6]
4.5	5-Hydroxy-7,4'-dimethoxy-6,8-dimethylflavone	Leaves of <i>M. cajuputi</i>	[37]
4.6	Kaempferol	Leaves of <i>M. huegelii</i>	[3]
		Leaves of <i>M. leucadendron</i>	[43]
		Leaf and stem of <i>M. quinquenervia</i>	[32]
		Leaves of <i>M. ericifolia</i>	[35]
		Leaves of <i>M. ericifolia</i>	[5]
4.7	Kaempferol-3-methyl ether	Leaves of <i>M. huegelii</i>	[3]
4.8	Kaempferol-3,7-dimethyl ether	Leaves of <i>M. huegelii</i>	[3]
4.9	Kaempferol-3,7,4'-trimethyl ether	Leaves of <i>M. huegelii</i>	[3]
4.10	Kaempferol-3- <i>O</i> -glucoside	Leaves of <i>M. leucadendra</i>	[44]
4.11	Kaempferol-3- <i>O</i> -xylosyl-(1 <sup>'''</sup> →2 <sup>''</sup> )-glucoside	Leaves of <i>M. ericifolia</i>	[5]
4.12	Kaempferol -3- <i>O</i> -rhamnoside	Leaves of <i>M. ericifolia</i>	[5]
		Leaves of <i>M. styphelioides</i>	[45]
4.13	Kaempferol-3- <i>O</i> -(2 <sup>''</sup> - <i>O</i> -galloyl)-glucuronide	Leaves of <i>M. squarrosa</i>	[6]
4.14	Luteolin	Leaves of <i>M. huegelii</i>	[3]
		Honey of <i>M. quinquenervia</i>	[46]
4.15	Luteolin-7,3 <sup>''</sup> -dimethyl ether	Leaves of <i>M. huegelii</i>	[3]
4.16	Miquelianin	Leaves of <i>M. leucadendron</i>	[43]
4.17	Myricetin	Leaves of <i>M. leucadendron</i>	[43]
4.18	Myricetin 3- <i>O</i> -(2 <sup>''</sup> - <i>O</i> -galloyl)- $\alpha$ -rhamnopyranoside	Leaves of <i>M. quinquenervia</i>	[47]
4.19	Myricitrin	Leaves of <i>M. leucadendron</i>	[43]
		Leaves of <i>M. ericifolia</i>	[5]
	Myricetin 3- <i>O</i> - $\beta$ -D-glucoside	Leaves of <i>M. quinquenervia</i>	[47]
4.20		Leaves of <i>M. leucadendra</i>	[44]
4.21	Myricetin 3- <i>O</i> - $\beta$ -D-glucuronide	Leaves of <i>M. quinquenervia</i>	[47]
4.22	Myricetin 3- <i>O</i> -rhamnoside	Leaves of <i>M. leucadendra</i>	[44]
4.23	Myricetin 3-rutinoside	Leaves of <i>M. leucadendra</i>	[44]
4.24	Myricetin-3- <i>O</i> - $\beta$ -4C1-galactopyranuronoid	Leaves of <i>M. leucadendra</i>	[44]
4.25	Myricetin 3- <i>O</i> -xylosyl-(1 <sup>'''</sup> →2 <sup>''</sup> )-glucoside	Leaves of <i>M. ericifolia</i>	[5]
	Quercetin	Leaves of <i>M. huegelii</i>	[3]
4.26		Fruits of <i>M. leucadendron</i>	[9]
		Leaves of <i>M. leucadendron</i>	[43]
		Leaf and stem of <i>M. quinquenervia</i>	[32]
		Leaves of <i>M. ericifolia</i>	[5]
		Honey of <i>M. quinquenervia</i>	[46]
		Leaves of <i>M. leucadendra</i>	[44]
4.27	Quercetin-3-methyl ether	Leaves of <i>M. huegelii</i>	[3]
4.28	Quercetin-3,7-dimethyl ether	Leaves of <i>M. huegelii</i>	[3]
4.29	Quercetin-3,7,3 <sup>''</sup> -trimethyl ether	Leaves of <i>M. huegelii</i>	[3]
4.30	Quercetin-3,7,4'-trimethyl ether	Leaves of <i>M. huegelii</i>	[3]
4.31	Quercetin-3- <i>O</i> -xyloglucoside	Leaves of <i>M. leucadendron</i>	[43]
4.32	Quercitrin	Leaves of <i>M. leucadendron</i>	[43]
		Leaf and stem of <i>M. quinquenervia</i>	[32]
		Leaves of <i>M. quinquenervia</i>	[47]
		Leaves of <i>M. ericifolia</i>	[5]
		Leaves of <i>M. cajuputi</i>	[37]
		Leaves of <i>M. leucadendra</i>	[44]
4.33	Quercetin-7- <i>O</i> -(6 <sup>''</sup> -galloyl)- $\beta$ -D-glucopyranoside	Leaves of <i>M. quinquenervia</i>	[47]
4.34	Quercetin-3- <i>O</i> -xylosyl-(1 <sup>'''</sup> →2 <sup>''</sup> )-glucoside	Leaves of <i>M. ericifolia</i>	[5]
4.35	Quercetin-3- <i>O</i> -galactoside	Leaves of <i>M. ericifolia</i>	[5]
4.36	Quercetin-3- <i>O</i> -glucoside	Leaves of <i>M. ericifolia</i>	[5]
4.37	Quercetin-3-rutinoside, rutin	Leaves of <i>M. leucadendra</i>	[44]
		Leaves of <i>M. leucadendra</i>	[44]
4.38	5,7,3 <sup>''</sup> ,4'-Tetrahydroxyflavone 2 <sup>''</sup> - <i>O</i> - $\beta$ -D-glucuronide	Leaves of <i>M. quinquenervia</i>	[47]
4.39	Tricetin	Honey of <i>M. quinquenervia</i>	[46]
		Honey of <i>M. quinquenervia</i>	[46]
		Leaves of <i>M. ericifolia</i>	[5]
		Leaves of <i>M. leucadendra</i>	[44]
<b>II-FLAVANS</b>			
4.40	Catechin	Leaves of <i>M. leucadendra</i>	[44]
4.41	Kryptostrobin	Flowers of <i>M. quinquenervia</i>	[41]
4.42	Leucadenone A	Leaves of <i>M. leucadenderon</i>	[42]
4.43	Leucadenone B	Leaves of <i>M. leucadenderon</i>	[42]
4.44	Leucadenone C	Leaves of <i>M. leucadenderon</i>	[42]
4.45	Leucadenone D	Leaves of <i>M. leucadenderon</i>	[42]
4.46	Melanervin	Flowers of <i>M. quinquenervia</i>	[41]
4.47	Strobopinin	Flowers of <i>M. quinquenervia</i>	[41]





**Figure 7.** Chemical structures of flavones (**4.1-4.39**) from the genus *Melaleuca*




**Figure 8.** Chemical structures of flavans (**4.40-4.47**) from the genus *Melaleuca*

#### 4.5. Simple Phenolics and Phenylpropanoids


Simple phenols are available in abundance in genus *Melaleuca* as represented by structures **5.1-5.37** (Table 5 and Figure 9 and 10). Phenylpropanoids are also present in this genus, which are simple secondary metabolites biosynthesized from shikimic acid pathway with the skeleton of C<sub>6</sub>-C<sub>3</sub>, such as the cinnamic acids, free (**5.26, 5.27 and 5.33**) or conjugated as in case of chlorogenic acid (**5.28**). Additionally, various stilbenes and phenylpropanoids derivatives, including lignans and neolignans have been reported from this genus, Table 5 and Figure 10. A unique group of neolignans, known as melaleucins A-C (**5.35-5.37**), have been discovered from the leaves of *M. bracteata* F.Muell. [48]. Another reported group of simple phenols is 1,4-benzopyrone ( $\gamma$ -benzopyrone group), commonly known as chromones (**5.1, 5.4-5.6, 5.14, 5.16, 5.21 and 5.22**), have been identified from the leaves of *M. cajuputi* and *M. leucadendron*. Several members of family Myrtaceae have been reported to contain anthraquinones [49]. However, only one anthraquinone derivative (parietin, **5.23**) has been identified from the leaves of *M. cajuputi*. In addition, ellagic acid in its free, methylated, or glycosylated forms represent a major class of phenols found in this genus (**5.7-5.13**, Figure 9).

**Table 5.** Simple phenolics and phenylpropanoids from the genus *Melaleuca*.

Compound	Name	Plant/ part	Ref.
5.1	Angustiforionol, 5-hydroxy-7-methoxy-2,6,8-trimethylchromone	Leaves of <i>M. cajuputi</i> Leaf of <i>M. leucaderdron</i>	[37] [50]
5.2	Brevifolincarboxylic acid	Fruits of <i>M. leucadendron</i>	[4]
5.3	Dimethylxanthoxylin	Leaf of <i>M. leucaderdron</i>	[50]
5.4	2,5-Dihydroxy-7-methoxy-2,6-dimethylchromanone	Leaves of <i>M. cajuputi</i>	[37]
5.5	2,5-Dihydroxy-7-methoxy-2,8-dimethylchromanone	Leaves of <i>M. cajuputi</i>	[37]
5.6	5,7-Dihydroxy-2,6,8-trimethylchromone	Leaves of <i>M. cajuputi</i> Honey of <i>M. quiquenervia</i>	[37] [51]
5.7	Ellagic acid	Leaves of <i>M. quinquenervia</i> Fruits of <i>M. leucadendron</i> Honey of <i>M. quiquenervia</i>	[52] [4] [51]
5.8	Ellagic acid 3- <i>O</i> -methylether	Leaves of <i>M. quinquenervia</i>	[52]
5.9	Ellagic acid 3-monomethylether-4- <i>O</i> -glucoside	Leaves of <i>M. leucadendra</i>	[44]
5.10	Ellagic acid 3,3'- <i>O</i> -dimethylether	Bark and Stem of <i>M. alternifolia</i>	[28]
5.11	Ellagic acid 3-methoxy 4- <i>O</i> - $\alpha$ -rhamnopyranoside	Leaves of <i>M. ericifolia</i>	[5] [35]
5.12	Ellagic acid 3,4,3'-tri- <i>O</i> -methylether	Leaves of <i>M. quinquenervia</i> Leaves of <i>M. cajuputi</i>	[52] [37]
5.13	Ellagic acid 3,3',4'-trimethoxy- 4- <i>O</i> - $\beta$ -glucoside	Leaves of <i>M. ericifolia</i>	[35]
5.14	Eugenitin	Leaves of <i>M. cajuputi</i> Leaf of <i>M. leucaderdron</i>	[37] [50]
5.15	Gallic acid	Leaves of <i>M. quinquenervia</i> Fruits of <i>M. leucadendron</i> Honey of <i>M. quiquenervia</i> Leaves of <i>M. ericifolia</i> leaves of <i>M. styphelioides</i> Leaves of <i>M. ericifolia</i> Leaves of <i>M. leucadendra</i>	[52] [9] [51] [35] [5] [45] [44]
5.16	Melachromone	Leaves of <i>M. cajuputi</i>	[37]
5.17	Methyl 2-acetyl-3,5-dihydroxyphenylacetate	Leaves of <i>M. cajuputi</i>	[37]
5.18	Methyl gallate	Leaves of <i>M. leucadendra</i>	[44]
5.19	Methyl tri- <i>O</i> -methyl gallate	Leaves of <i>M. ericifolia</i>	[35]
5.20	Methylxanthoxylin	Leaf of <i>M. leucaderdron</i>	[50]
5.21	Noreugenin	Leaves of <i>M. cajuputi</i>	[37]
5.22	Noreungenin 8- <i>C</i> -glucosyl	Leaf of <i>M. leucadendra</i>	[44]
5.23	Parietin	Leaves of <i>M. cajuputi</i>	[37]
5.24	<i>p</i> -Hydroxybenzoic acid	Leaves of <i>M. ericifolia</i>	[5]
5.25	Protocatechuic acid	Fruits of <i>M. leucadendron</i>	[4]
<b>PHENYLPROPANOIDS and STILBENES</b>			
5.26	3',4'-Dimethoxy cinnamaldehyde	Leaves of <i>M. ericifolia</i>	[35]
5.27	Caffeic acid	Leaves of <i>M. leucadendra</i>	[44]
5.28	Chlorogenic acid	Honey of <i>M. quiquenervia</i>	[51]
5.29	Melaleucin A	Leaves of <i>M. bracteata</i>	[48]
5.30	Melaleucin B	Leaves of <i>M. bracteata</i>	[48]
5.31	Melaleucin C	Leaves of <i>M. bracteata</i>	[48]
5.32	Oxyresveratrol	Fruits of <i>M. leucadendron</i>	[9]
5.33	<i>p</i> -Coumaric acid	Honey of <i>M. quiquenervia</i>	[51]
5.34	Piceatannol	Fruits of <i>M. leucadendron</i>	[9],[4]
5.35	Piceatannol-4'- <i>O</i> - $\beta$ -D-gucopyranoside	Fruits of <i>M. leucadendron</i>	[4]
5.36	Piceatannol-4'- <i>O</i> - $\beta$ -D-gucopyranose-6''- <i>O</i> -gallate	Fruits of <i>M. leucadendron</i>	[4]
5.37	Scirpusin B	Fruits of <i>M. leucadendron</i>	[9]



**Figure 9.** Structures of simple and polyphenolics (**5.1-5.25**) from the genus *Melaleuca*




**Figure 10.** Structures of phenylpropanoids and stilbenes (**5.26-5.37**) from the genus *Melaleuca*.

#### 4.6. Hydrolysable Tannins


Hydrolysable tannins are known for their antioxidants effects and are widely distributed in the plant kingdom. Myrtaceae is considered as one of the common plant families known for its tannin contents, especially oligomeric ellagitannins [53]. In *Melaleuca* genus, hydrolysable tannins represent a major class of biomolecules with a fewer number of gallitannins (**6.5-6.12**) compared to ellagitannins (**6.1-6.4** and **6.13-6.30**), Table 6 and Figures 11-13. A great variation in the molecular weight of these compounds is observed. *Melaleuca squarrosa* Donn ex Sm. is characterized by the presence of C-glycosidic ellagitannins oligomers, such as melasquanins A and C (**6.18** and **6.19**) and squarrosanin A-C (**6.26-6.28**), respectively. Also, the leaves of *M. styphelioides* have been reported to contain several ellagitannins, such as pedunculagin (**6.21**), pterocarinin A (**6.23**), stachyurin (**6.25**), tellimagrandin I (**6.29**), and tellimagrandin II (**6.30**), in addition to the galloyl esters 1,2,3,6-tetra-*O*-galloyl- $\beta$ -D-glucose (**6.11**), and 1,2,3,4,6-penta-*O*-galloyl- $\beta$ -D-glucose (**6.12**).

**Table 6.** Hydrolysable tannins from the genus *Melaleuca*


Compound	Name	Plant/ part	Ref.
6.1	4-Benzoyl-2- <i>C</i> - $\beta$ -glucopyranosyl-3,5-dihydroxy-6-methylphenyl- <i>O</i> - $\beta$ -D-glucopyranoside	Leaves of <i>M. quinquenervia</i>	[54]
6.2	Castalin	Leaves of <i>M. quinquenervia</i>	[52]
6.3	Castalagin	Fruits of <i>M. leucadendron</i>	[4]
6.4	Casuarinin	Fruits of <i>M. leucadendron</i>	[4] [55] [53]
6.5	1,6-Di- <i>O</i> -galloyl- $\beta$ -glucose	Leaves of <i>M. ericifolia</i>	[5]
6.6	2,6-Di- <i>O</i> -galloyl-glucose	Leaves of <i>M. ericifolia</i>	[5]
6.7	1- <i>O</i> - <i>p</i> -Hydroxybenzoyl-6- <i>O</i> -galloyl- $\beta$ -glucose	Leaves of <i>M. ericifolia</i>	[5]
6.8	2- <i>O</i> - <i>p</i> -Hydroxybenzoyl-6- <i>O</i> -galloyl-( $\alpha/\beta$ )- <sup>4</sup> C <sub>1</sub> -glucopyranose	Leaves of <i>M. ericifolia</i>	[5]
6.9	2,3-Di- <i>O</i> -galloyl glucose	Leaves of <i>M. ericifolia</i>	[5]
6.10	1,2,3-Tri- <i>O</i> -galloyl- $\beta$ -D-glucose	Fruits of <i>M. leucadendron</i>	[4]
6.11	1,2,3,6-Tetra- <i>O</i> -galloyl- $\beta$ -D-glucose	Fruits of <i>M. Leucadendron</i>	[4]
6.12	1,2,3,4,6-penta- <i>O</i> -galloyl- $\beta$ -D-glucose	Leaves of <i>M. styphelioides</i>	[45]
6.13	1,2-Di- <i>O</i> -galloyl-3- <i>O</i> -digalloyl-4,6- <i>O</i> -hexahydroxydiphenyl- $\beta$ -D-glucose	Fruits of <i>M. Leucadendron</i>	[4]
6.14	Ericifolin; eugenol-5- <i>O</i> - $\beta$ -6'- <i>O</i> -galloylglucopyranoside	Leaves of <i>M. styphelioides</i>	[45]
6.15	Grandinin	Fruits of <i>M. leucadendron</i>	[4]
6.16	2,3- <i>O</i> -Hexahydroxydiphenol-( $\alpha/\beta$ )-D- <sup>4</sup> C <sub>1</sub> -glucopyranose	Leaves of <i>M. quinquenervia</i>	[52]
6.17	3-Hydroxy-5-methoxy-4-methylphenyl $\beta$ -D-glucopyranoside	Leaves of <i>M. quinquenervia</i>	[54]
6.18	Melasquanin A	Leaves of <i>M. squarrosa</i>	[6] [55]
6.19	Melasquanin C	Leaves of <i>M. squarrosa</i>	[6]
6.20	Oenothin	Fruits of <i>M. leucadendron</i>	[4]
6.21	Pedunculagin	Leaves of <i>M. styphelioides</i>	[45]
6.22	Praecoxin A	Leaves of <i>M. ericifolia</i>	[56]
6.23	Pterocarinin A	Leaves of <i>M. styphelioides</i>	[45]
6.24	Rugosin D	Fruits of <i>M. leucadendron</i>	[4]
6.25	Stachyurin	Leaves of <i>M. styphelioides</i>	[4],[45]
6.26	Squarrosanin A	Fruits of <i>M. leucadendron</i>	[55],[53]
6.27	Squarrosanin B	Leaves of <i>M. squarrosa</i>	[6]
6.28	Squarrosanin C	Leaves of <i>M. squarrosa</i>	[6]
6.29	Tellimagrandin I	Leaves of <i>M. squarrosa</i>	[6]
6.30	Tellimagrandin II	Leaves of <i>M. styphelioides</i>	[45]
		Fruits of <i>M. Leucadendron</i>	[4]
		Leaves of <i>styphelioides</i>	[45]



**Figure 11.** Structures of hydrolysable tannins (**6.1-6.17**) from the genus *Melaleuca*



**Figure 12.** Structures of hydrolysable tannins (**6.18-6.23**) from the genus *Melaleuca*




**Figure 13.** Structures of hydrolysable tannins (6.24-6.30) from the genus *Melaleuca*

#### 4.7. Benzylic Phloroglucinol-Terpene Hybrids

As a member of the family Myrtaceae, plants of the genus *Melaleuca* are rich in phloroglucinol derivatives, including phloroglucinol-terpene adducts [57]. A unique class of benzylic phloroglucinol-terpene hybrids namely leumelaleucols A-L (7.1-7.13) and melaleucadines A and B (7.14 and 7.15), respectively, has been recently discovered [12,13], Table 7 and Figure 14. However, this class of meroterpenoids has been only identified from different parts of *M. leucadendron*, which paved the way towards more researches on other *Melaleuca* spp.

**Table 7.** Benzylic phloroglucinol-terpene hybrids from the genus *Melaleuca*


Compound	Name	Plant/ part	Ref.
7.1	(+)-Leumelaleucol A	Fruits of <i>M. leucadendron</i>	[13]
7.2	(-)-Leumelaleucol A	Fruits of <i>M. leucadendron</i>	[13]
7.3	Leumelaleucols B	Fruits of <i>M. leucadendron</i>	[13]
7.4	Leumelaleucols C	Fruits of <i>M. leucadendron</i>	[13]
7.5	Leumelaleucols D	Fruits of <i>M. leucadendron</i>	[13]
7.6	Leumelaleucols E	Fruits of <i>M. leucadendron</i>	[13]
7.7	Leumelaleucols F	Fruits of <i>M. leucadendron</i>	[13]
7.8	Leumelaleucols G	Fruits of <i>M. leucadendron</i>	[13]
7.9	Leumelaleucols H	Fruits of <i>M. leucadendron</i>	[13]
7.10	Leumelaleucols I	Fruits of <i>M. leucadendron</i>	[13]
7.11	Leumelaleucols J	Fruits of <i>M. leucadendron</i>	[13]
7.12	Leumelaleucols K	Fruits of <i>M. leucadendron</i>	[13]
7.13	Leumelaleucols L	Fruits of <i>M. leucadendron</i>	[13]
7.14	Melaleucadine A	Branches and leaves of the <i>M. leucadendron</i>	[12]
7.15	Melaleucadine B	Branches and leaves of the <i>M. leucadendron</i>	[12]



**Figure 14.** Structures of benzylic phloroglucinol-terpene hybrids (7.1-7.15)

#### 4.8. Miscellaneous Group

This class of miscellaneous compounds (Table 8 and Figure 15), comprising a group of simple terpenoidal glucosides, including citroside A (8.1), 2-endo- $\beta$ -D-glucopyranosyloxy-1,8-cineole (8.2), 2-exo- $\beta$ -D-glucopyranosyloxy-1,8-cineole (8.3), and roseoside (8.4), was identified from the leaves of *M. quinquenervia*. A minor group of alkaloids represented by proline derivatives (8.5-8.8) has been reported from several *Melaleuca* spp., including *M. alternifolia* L., *M. lanceolata* Otto., and *M. uncinata* R.Br.




**Figure 15.** Structures of miscellaneous group (8.1-8.8) from the genus *Melaleuca*

A high content of fatty acids and hydrocarbon derivatives (8.9-8.34) was detected in several parts of these plants, Table 8 and Figure 16. The leaves and stems of *M. quinquenervia* showed the presence of palmitic acid as the major component of the saturated fatty acids content [32], whereas linolenic acid was the main constituent of the unsaturated fatty acids [32]. Gas chromatographic analysis of the hexane fraction of the barks and stems of *M. alternifolia* resulted in the identification of palmitate, stearate, and linolenate, in addition to several hydrocarbon derivatives, including 4-methyl-pentan-3-one (8.25), undecan-2-one (8.33), hepta-2,4-dien-6-al (8.23), and others [28]. Other miscellaneous compounds are represented by the aromatic hydrocarbon, naphthalene (8.35) [8].

**Table 8.** Miscellaneous group of compounds from the genus *Melaleuca*

Compound	Name	Plant/ part	Ref.
<b>I- TERPENOIDAL GLUCOSIDES</b>			
8.1	Citroside A	Leaves of <i>M. quinquenervia</i>	[54]
8.2	2-Endo- $\beta$ -D-glucopyranosyloxy-1,8-cineole	Leaves of <i>M. quinquenervia</i>	[54]
8.3	2-Exo- $\beta$ -D-glucopyranosyloxy-1,8-cineole	Leaves of <i>M. quinquenervia</i>	[54]
8.4	Roseoside	Leaves of <i>M. quinquenervia</i>	[54]
<b>II- PROLINE DERVATIVES</b>			
8.5	4-Hydroxy- <i>N</i> -methyl- proline	<i>Melaleuca</i> spp.	[58]
		<i>M. lanceolata</i> and <i>M. uncinata</i>	[59]
		<i>M. lanceolata</i>	[60]
8.6	4-Hydroxy- <i>N,N</i> -dimethyl-proline (=Betonidine)	<i>Melaleuca</i> spp.	[58]
		<i>M. lanceolata</i> and <i>M. uncinata</i>	[59]
		Bark and stem of <i>M. alternifolia</i>	[28]
8.7	<i>N</i> -Methyl-L-proline	<i>Melaleuca</i> spp.	[58]
8.8	L-Proline	<i>Melaleuca</i> spp.	[58]
<b>III- FATTY ACIDS</b>			
8.9	Capric acid	Leaf and stem of <i>M. quinquenervia</i>	[32]
8.10	Lauric acid	Leaf and stem of <i>M. quinquenervia</i>	[32]
8.11	Linoleic acid	Leaf and stem of <i>M. quinquenervia</i>	[32]
8.12	Linolenic acid	Leaf and stem of <i>M. quinquenervia</i>	[32]
		Bark and stem of <i>M. alternifolia</i>	[28]
8.13	Oleic acid	Leaf and stem of <i>M. quinquenervia</i>	[32]
8.14	Pelargonic acid	Leaf and stem of <i>M. quinquenervia</i>	[32]
8.15	Palmitic acid	Leaf and stem of <i>M. quinquenervia</i>	[32]
8.16	Palmitoleic acid	Leaf and stem of <i>M. quinquenervia</i>	[32]
		Bark and stem of <i>M. alternifolia</i>	[28]
8.17	Stearic acid	Leaf and stem of <i>M. quinquenervia</i>	[32]
		Bark and stem of <i>M. alternifolia</i>	[28]
<b>IV- HYDROCARBONS AND THEIR DERIVATIVES</b>			
8.18	Decane (C10)	Bark and stem of <i>M. alternifolia</i>	[28]
8.19	Docosane (C22)	Leaf and stem of <i>M. quinquenervia</i>	[32]
8.20	Dodecane	Bark and stem of <i>M. alternifolia</i>	[28]
8.21	Eicosane (C20)	Leaf and stem of <i>M. quinquenervia</i>	[32]
8.22	Hexadecane (C16)	Leaf and stem of <i>M. quinquenervia</i>	[32]
8.23	Hepta-2,4-dien-6-inol	Bark and stem of <i>M. alternifolia</i>	[28]
8.24	6-Hydroxy-4,6-dimethyl-3-hepten-2-one	Heartwood of <i>M. leucadendron</i>	[8]
8.25	4-Methyl-pentan-3-one	Bark and stem of <i>M. alternifolia</i>	[28]
8.26	6-Methyldec-4-ene ( $^4\Delta$ C11)	Bark and stem of <i>M. alternifolia</i>	[28]
8.27	Methyl 6-hydroxyhexanoate	Bark and stem of <i>M. alternifolia</i>	[28]
8.28	Octadecane (C18)	Leaf and stem of <i>M. quinquenervia</i>	[32]
8.29	Octcosane (C28)	Leaf and stem of <i>M. quinquenervia</i>	[32]
8.30	Tridecane (C13)	Bark and stem of <i>M. alternifolia</i>	[28]
8.31	Tetradecane (C14)	Leaf and stem of <i>M. quinquenervia</i>	[32]
		Bark and stem of <i>M. alternifolia</i>	[28]
8.32	Triacontane (C30)	Leaf and stem of <i>M. quinquenervia</i>	[32]
8.33	Undecan-2-one	Bark and stem of <i>M. alternifolia</i>	[28]
8.34	Undecane (C11)	Bark and stem of <i>M. alternifolia</i>	[28]
<b>V- OTHER MISCELLANEOUS COMPOUNDS</b>			
8.35	Naphthalene	Heartwood of <i>M. leucadendron</i>	[8]





**Figure 16.** Structures of fatty acids and hydrocarbon derivatives (8.9-8.35)

It is worth mentioning that some compounds have been identified from the essential oils of *Melaleuca* spp., such as anthraquinones, coumarins, and alkaloids, which are commonly distilled with the oil. Additionally, other compounds, including diterpenes, fat, and other non-volatile compounds can be isolated from volatile oils using methods other than distillation [61].

## 5. Pharmacological Activities of the Non-volatile Components

Diverse biological activities have been reported for plants of the genus *Melaleuca*. The reported activities, including antimicrobial, antiparasitic, antioxidant, anti-inflammatory, acetylcholinesterase inhibition, neuroprotective, molluscicidal, antischistosomal, immunostimulant, anticancer, antihistaminic, cardiovascular, hepatoprotective, antisecretory, antiulcerogenic, and liver microsomal enzyme inhibition activity, are discussed herein in detail.

The reported biological assays were conducted either on the crude extracts from different plant parts, such as leaves, flowers, stems, and barks, or using pure isolated compounds of known pharmacological actions, such as betulinic acid, ursolic acid, oxyresveratrol, piceatannol, flavonoids, and ellagitannins. Betulinic acid is an attractive point for many researchers towards the derivatization and biological evaluation of several pharmacological activities, especially those related to anticancer drug discovery [7, 62].

### 5.1. Antimicrobial and Antiparasitic Activity

The methanol extracts of the flowers and the leaves of *M. cajuputi* exhibited antibacterial effects against *Staphylococcus epidermidis*, *Staphylococcus aureus*, and *Bacillus cereus*. It was suggested that the antimicrobial activities may be attributable to the high flavonoid and phenolic contents of these extracts [63]. Also, the methanol extract of *M. cajuputi* flowers exhibited inhibitory activity against the filarial worm *Brugia pahangi* through reduction in worm motility and viability and in microfilarial release. Furthermore, *M. cajuputi* flower extract inhibited the endosymbiont *Wolbachia* bacteria of *B. pahangi* through significant reduction in worm viability, enhancing the antifilarial activity [64]. The methanol extract of the fresh *M. cajuputi* leaf waste, left after distillation of the essential oil, exhibited a low to strong antibacterial activity against the tested Gram positive bacteria, including *Bacillus* sp., *Enterococcus faecalis*, *S. aureus*, and Gram negative bacteria, including *Vibrio cholerae* and *Shigella dysenteriae* [65]. In addition, three isolated neolignans from the leaves of *M. bracteata* named melaleucins A–C (**5.35–5.37**) were investigated for their antibacterial activity against *S. aureus*, *E. coli*, and methicillin-resistant *S. aureus* (MRSA). Both melaleucins A and C exhibited significant inhibitory activities against *S. aureus* and MRSA [48]. Laribi et al. (2020) compared the antifungal activities of the volatile oil and non-volatile extracts of *M. styphelioides* Sm. leaves against *Aspergillus niger*, *Rhizopus nigricans*, and *Penicillium digitatum*. The tested samples showed different selectivity for each microorganism studied, where the volatile oil was predominantly effective against *A. niger*, whereas *P. digitatum* was found to be the most susceptible species for the aqueous extract and *R. nigricans* was more susceptible to the methanol extract [66].

### 5.2. Antioxidant and Anti-inflammatory Effects

Grandinin (**6.20**), the major polyphenol in *M. quinquenervia* leaves, was found to display antioxidant activity [52]. Also, the C-glucosidic ellagitannins isolated from the leaves of *M. squarrosa*, including complex tannins and oligomers have been shown to be more effective radical scavengers in 2,2-diphenyl-1-picrylhydrazyl (DPPH) assay than its flavonoids and ordinary ellagitannins [6]. Al-Abd et al. (2015) investigated the antioxidant and radical scavenging activity of the methanol extracts of *M. cajuputi* using DPPH radical scavenging,  $\beta$ -carotene bleaching, and ferric reducing antioxidant power (FRAP) assays. The flower extract showed a better activity than that of the leaf extract explained by the higher total phenolic contents in the flower extract compared to that of the leaf extract [63]. The nephroprotective effect of phenolic compounds isolated from the leaf extract of *M. styphelioides*, including pedunculagin (**6.21**), pterocarinin A (**6.23**), tellimagrandin I (**6.29**), and casuarinin (**6.4**), towards diabetes mellitus and kidney stone oxidative stress-mediated kidney damage has been studied on a glucose- and oxalate-challenged NRK-49F cell model [67]. The results showed that the tested compounds restored catalase and superoxide dismutase antioxidant activity, reversed lipid peroxidation, and protected the kidney fibroblast cells from glucose- and oxalate-mediated stress [67]. Also, the methanolic extract of the leaves of *M. styphelioides* exhibited antioxidant activity using DPPH, ABTS<sup>•+</sup>, and FRAP assays. It inhibited pro-inflammatory mediators, including iNOS, NF-kB,

COX-2, and inter-cellular adhesion molecule-1 (ICAM-1) in interferon gamma/histamine (IFN-g/H)-stimulated inflammation in NCTC 2544 human keratinocytes. LC/MS-MS analysis of the tested extract suggested that the phenolic compounds, mainly as quercetin, gallic acid, and ellagic acid, are responsible of these activities [68]. The ethanolic extract of *M. leucadendron* flowers exhibited an *in vitro* photoprotective effect towards UVB-induced oxidative stress in human skin keratinocytes (HaCaT). This has been concluded from the obtained DPPH antioxidant capacity, the maintained levels of ROS, GSH, glutathione peroxidase (GPx), SOD, catalase, and COX-2 expressions and reduced impairment of transmembrane transport and pro-apoptotic protein expression of the tested cells [69]. Casuarinin (**6.4**), an ellagitannin isolated from *M. leucadendra* L. leaves, suppressed the expression of NF- $\kappa$ B, COX-2, and iNOS to their normal levels. It also increased the level of reduced glutathione, catalase, and prostaglandin E<sub>2</sub> in ethanol-induced gastric ulcer in rats [70]. Herbal infusion of *M. citrolens* Barlow. leaves (gulbarn) infusions showed similar antioxidant activity as commercial green tea [71]. Similarly, the leaf extract of *M. leucadendra* and its isolate myricetin-3-*O*- $\beta$ -<sup>4</sup>C<sub>1</sub>-galactopyranuronoid (**4.24**) exhibited strong antioxidant activity using DPPH and oxygen radical absorption capacity (ORAC) assay [44]. Regarding the non-phenolic components, betulinic acid (**1.26**) isolated from *M. bracteata* displayed antioxidant potential through preventing catalase inhibition as catalase enzyme catalyzes the decomposition of harmful H<sub>2</sub>O<sub>2</sub> into H<sub>2</sub>O, and O<sub>2</sub> [72]. Also, evaluation of the anti-inflammatory activity of betulinic acid isolated from *M. bracteata* using cotton pellet-induced granuloma model showed significant dose-dependent inhibition. Furthermore, betulinic acid significantly inhibited COX-1 and COX-2 activity in concentration-dependent manner [72]. Additionally, betulinic acid exhibited antioxidant activity through preventing SOD inhibition [72 73]. Superoxide dismutase (SOD) acts as an inhibitory agent of neutrophil-mediated inflammation and serves as an antioxidant through splitting superoxide radicals into the less harmful O<sub>2</sub> and H<sub>2</sub>O [72, 73].

### 5.3. Acetylcholinesterase and Neuroprotective Activity

The benzylic phloroglucinol-terpene hybrids, melaleucadines A (**7.14**) and B (**7.15**), Table 7 and Figure 14, isolated from the leaves and branches of *M. leucadendron*, exhibited potential neuroprotective effect on the dopaminergic neuron cells (PC12) from cortisone-induced neural damage [12]. Recently, a larger group of benzylic phloroglucinol-terpene hybrids (*Syn.* benzyl phloroglucinol-based meroterpenoids), isolated from the fruits of *M. leucadendron*, Table 7 and Figure 14, has been *in vitro* evaluated for its neuroprotective effect of cortisone-induced neurotoxicity on PC12 cells. The reported results showed that (+)-leumelaleucol A (**7.1**), leumelaleucol B (**7.3**), leumelaleucol D (**7.5**), leumelaleucol E (**7.6**), leumelaleucol F (**7.7**), and leumelaleucol G (**7.8**) increased the survival of treated PC-12 cells with the highest activity attributed to leumelaleucol G and E, respectively [13]. Betulinic acid (**2.26**) isolated from *M. bracteata* showed significant inhibition of acetylcholinesterase activity [72].

### 5.4. Molluscicidal and Antischistosomal Activity

A moderate biocidal effect of the methanol extract of *M. styphelioides* leaves and the isolated compounds, kaempferol-3-*O*- $\alpha$ -rhamnoside **1.12**, pedunculagin **6.21**, and casuarinin **6.4**, has been revealed against *Biomphalaria alexandrina* snails [74].

### 5.5. Immunostimulant Activity

Hashim et al. (2018) reported that *M. leucadendra* fresh leaf extract significantly increased the phagocytic activity of neutrophils, suggesting that it has an immunostimulant effect [44].

### 5.6. Anticancer Activity

Antiproliferative activity to cultured mouse T-lymphoma cells by hexane and dichloromethane extracts, as well as eugenitin (**5.14**) from *M. leucadendron* leaves, has been reported [50]. Also, the

antiproliferative effect of the triterpenes of *M. ericifolia* leaf extract on malignant mouse +SA mammary epithelial cells has been investigated. A group of isolated 28-norlupanes (Table 3 and Figure 3), such as 28-norlup-20(29)-en-3 $\beta$ ,17 $\beta$ -diol (**1.30**), 28-norlup-20(29)-en-3 $\beta$ -hydroxy-17 $\alpha$ -hydroperoxide (**1.31**), 28-norlup-20(29)-en-3 $\beta$ -hydroxy-17 $\beta$ -hydroperoxide (**1.32**), and 20S-17 $\beta$ ,29-epoxy-28-norlupan-3 $\beta$ -ol (**1.28**), showed higher antiproliferative effect compared to other triterpene derivatives, including ursolaldehyde (**1.10**), ursolic acid (**1.11**), asiatic acid (**1.1**), betulin (**1.24**), betulinaldehyde (**1.25**), betulinic acid (**1.26**), and platanic acid (**1.33**) [7]. The cytotoxicity of *M. leucadendra* leaf extract alongside its isolate, myricetin-3-*O*- $\beta$ -<sup>4</sup>C<sub>1</sub>-galactopyranuronoid (**4.24**), has been evaluated against three different tumor cell lines: liver carcinoma cell line (Huh-7), breast cancer (MCF-7), and colon cancer (HCT 116). The results showed that the tested natural products exhibited higher cytotoxic activity towards HCT 116 [44].

### 5.7. Antihistaminic Activity

The chloroform and methanol extracts of the fruits of *M. leucadendron* strongly inhibited histamine release from rat mast cells. Ursolic acid (**1.11**) from the chloroform extract and two stilbenes, oxyresveratrol (**5.32**) and piceatannol (**5.34**) from the methanol extract, have been isolated as the bioactive compounds [9].

### 5.8. Cardiovascular Activity

Glycosides from the leaves of *M. quinquenervia* inhibit the vascular contraction in aortic rings of rats [54]. Furthermore, betulinic acid (**1.26**) isolated from *M. bracteata* leaves possesses antithrombotic effect, antiplatelet aggregations, and anticoagulants potential [75].

### 5.9. Hepatoprotective Activity

The total leaf extract of *M. styphelioides* showed a significant hepatoprotective effect in a CCl<sub>4</sub>-induced toxicity model in mice. It decreased the levels of AST, ALT, ALP, and MDA and increased the level of GSH and antioxidant enzymes compared to those of the CCl<sub>4</sub>-treated group. It has restored the liver architecture and reduced the fatty changes, central vein congestion, Kupffer cell hyperplasia, inflammatory infiltration, and necrosis induced by CCl<sub>4</sub> [45]. Al-Sayed and Esmat (2016) investigated the *in vitro* hepatoprotective compounds in *M. styphelioides* leaves using the CCl<sub>4</sub>-challenged HepG2 cell model. Marked hepatoprotections by 1,2,3,4,6-penta-*O*-galloyl- $\beta$ -D-glucose (**6.12**), pedunculagin (**6.21**), tellimagrandin I (**6.29**), and tellimagrandin II (**6.30**), Table 6 and Figure 13, were observed as revealed from decreased AST and ALT levels and increased GSH levels [76]. Praecoxin A (**6.22**), an ellagitannin from the leaves *M. ericifolia*, showed promising hepatoprotective activity in CCl<sub>4</sub>-induced toxicity in mice, where it decreased the CCl<sub>4</sub>-induced increase in AST, ALP, ALT, total bilirubin, and malondialdehyde (MDA). Histological results indicated a marked improvement in hepatocyte inflammation, degeneration, cell infiltration, necrosis, and hemorrhage [56].

### 5.10. Antisecretory and Antiulcerogenic Activities

Adesanwo et al. (2009) investigated the antiulcer and antisecretory effect of the ethyl acetate extract of the stem bark of *M. bracteata* on gastric acid secretion and indomethacin-induced ulceration in albino rats. The results indicated that the tested extract significantly reduced gastric secretion and ulceration. Moreover, the researchers suggested that the presence of betulinic acid (**1.26**) and oleanolic acid (**1.20**) in the extract may account for this antiulcer action [77]. Casuarinin (**6.4**), isolated from *M. leucadendra* leaves, exhibited a gastroprotective effect on ethanol-induced ulceration in rats. The potent gastroprotective effect of casuarinin was shown to be attributed to its anti-inflammatory, antioxidant, and antiapoptotic effects [70].

### 5.11. Inhibition of liver microsomal enzyme

The methanol extract of *M. leucadendron* leaves exhibited strong inhibitory activity on cytochrome P450 2D6 (CYP2D6) [78].

## 6. Conclusions

Natural products have a vital role in pharmaceutical industries. *Melaleuca* species have a diverse metabolic pool of phytochemicals, and their essential oils are commonly used in ayurvedic medicines. Huge amounts of these plants' waste are left behind after the distillation of the essential oils from various species. This review unveiled that melaleuca trees are rich sources of bioactive compounds, including triterpenoids (e.g. lupane, nolupane, ursane, and oleanane), flavonoids, and phenolics. Also, a large group of hydrolysable tannins, reported from different *Melaleuca* spp., has been shown to be biologically and pharmacologically active, such as grandinin, pterocarinin, casuarinin, etc. Being a member of the Myrtaceae family, *Melaleuca* spp. are also characterized by the presence of benzylic phloroglucinol-terpene hybrids, which have been reported as neuroprotective compounds. Consequently, with recent technologies in extraction, isolation, and purification and with advancement in biological and pharmacological assay models, this plant waste could participate in new drug discoveries and development.

## Competing Interests

The author has declared that no competing interests exist.

## Acknowledgments

This publication was supported by the Deanship of Scientific Research at Prince Sattam bin Abdulaziz University, Alkharij, Saudi Arabia. Sincere thanks to Miss Rokaia Badreldin for English language editing.

ORCID 

Fatma M. Abdel Bar: [0000-0002-4690-4975](https://orcid.org/0000-0002-4690-4975)

## References

- [1] I. Southwell and R. Lowe (1999). Tea tree: the genus *Melaleuca*, Harwood Academic Publisher, Amsterdam, Netherlands.
- [2] P. Shukla and S. P. Misra (1997). An introduction to taxonomy of Angiosperms, Vikas Publishing House Pvt. Ltd., Delhi.
- [3] E. Wollenweber, R. Wehde, M. Dorr, G. Lang and J. F. Stevens (2000). C-methyl-flavonoids from the leaf waxes of some Myrtaceae, *Phytochemistry* **55**, 965-970.
- [4] T. Yoshida, T. Maruyama, A. Nitta and T. Okuda (1996). An hydrolysable tannin and accompanying polyphenols from *Melaleuca leucadendron*, *Phytochemistry* **42**, 1171-1173.
- [5] S. A. Hussein, A. N. Hashim, R. T. El-Sharawy, M. A. Seliem, M. Linscheid, U. Lindequist and M. A. Nawwar (2007). Ericifolin: an eugenol 5-*O*-galloylglucoside and other phenolics from *Melaleuca ericifolia*, *Phytochemistry* **68**, 1464-1470.
- [6] M. Yoshimura, H. Ito, K. Miyashita, T. Hatano, S. Taniguchi, Y. Amakura and T. Yoshida (2008). Flavonol glucuronides and C-glucosidic ellagitannins from *Melaleuca squarrosa*, *Phytochemistry* **69**, 3062-3069.
- [7] F. M. Abdel Bar, A. M. Zaghoul, S. V. Bachawal, P. W. Sylvester, K. F. Ahmad and K. A. El Sayed (2008). Antiproliferative triterpenes from *Melaleuca ericifolia*, *J. Nat. Prod.* **71**, 1787-1790.
- [8] C. K. Lee (1998). New triterpenes from the heartwood of *Melaleuca leucadendron* L., *J. Chin. Chem. Soc.* **45**, 303-306.

- [9] T. Tsuruga, Y. T. Chun, Y. Ebizuka and U. Sankawa (1991). Biologically active constituents of *Melaleuca leucadendron*: inhibitors of induced histamine release from rat mast cells, *Chem. Pharm. Bull. (Tokyo)* **39**, 3276-3278.
- [10] C. K. Lee (1998). Ursane triterpenoids from leaves of *Melaleuca leucadendron*, *Phytochemistry* **49**, 1119-1122.
- [11] C. K. Lee (1998). A new norlupene from the leaves of *Melaleuca leucadendron*, *J. Nat. Prod.* **61**, 375-376.
- [12] X. Xie, L. Wu, Z. Cui, M. Yang, Y. Yin, J. Luo and L. Kong (2019). Melaleucadines A and B: Two rare benzylic phloroglucinol-terpene hybrids from *Melaleuca leucadendron*, *Tetrahedron Lett.* **60**, 1011-1013.
- [13] L. Wu, X. Xie, X.-B. Wang, M.-H. Yang, J. Luo and L.-Y. Kong (2020). Diverse benzyl phloroglucinol-based meroterpenoids from the fruits of *Melaleuca leucadendron*, *Tetrahedron* **76**, 131326.
- [14] S. B. Jones and A. E. Luchsinger (1986). Plant systematics, McGraw-Hill Book Co., New York.
- [15] G. Singh (2019). Plant systematics: An integrated approach, CRC Press - Taylor & Francis Group, New York.
- [16] A. Padovan, A. Keszei, C. Külheim and W. J. Foley (2014). The evolution of foliar terpene diversity in Myrtaceae, *Phytochem. Rev.* **13**, 695-716.
- [17] S. K. T. Seraglio, M. Schulz, P. Nehring, F. Della Betta, A. C. Valse, H. Daguer, L. V. Gonzaga, R. Fett and A. C. O. Costa (2018). Nutritional and bioactive potential of Myrtaceae fruits during ripening, *Food Chem.* **239**, 649-656.
- [18] E. Wollenweber, R. Wehde, M. Dörr, G. Lang and J. F. Stevens (2000). C-methyl-flavonoids from the leaf waxes of some Myrtaceae, *Phytochemistry* **55**, 965-970.
- [19] E. A. Porter, E. N. Lughadha and M. S. J. Simmonds (2000). Taxonomic significance of polyhydroxyalkaloids in the Myrtaceae, *Kew Bull.* **55**, 615-632.
- [20] C. F. Carson, K. A. Hammer and T. V. Riley (2006). *Melaleuca alternifolia* (Tea Tree) oil: a review of antimicrobial and other medicinal properties, *Clin. Microbiol. Rev.* **19**, 50-62.
- [21] R. A. Hill and J. D. Connolly (2018). Triterpenoids, *Nat. Prod. Rep.* **35**, 1294-1329.
- [22] P. Dzubak, M. Hajdich, D. Vydra, A. Hustova, M. Kvasnica, D. Biedermann, L. Markova, M. Urban and J. Sarek (2006). Pharmacological activities of natural triterpenoids and their therapeutic implications, *Nat. Prod. Rep.* **23**, 394-411.
- [23] H. Senol, K. Cokuludag, A.S. Aktas, S. Atasoy, A. Dag and G. Topcu (2020). Synthesis of new fatty acid derivatives of oleanane and ursane triterpenoids and investigation of their in vitro cytotoxic effects on 3T3 fibroblast and PC3 prostate cancer cell lines, *Org. Commun.* **13**, 114-126.
- [24] A. Kapil and S. Sharma (1995). Effect of oleanolic acid on complement in adjuvant- and carrageenan-induced inflammation in rats, *J. Pharm. Pharmacol.* **47**, 585-587.
- [25] T. Ringbom, L. Segura, Y. Noreen, P. Perera and L. Bohlin (1998). Ursolic acid from *Plantago major*, a selective inhibitor of cyclooxygenase-2 catalyzed prostaglandin biosynthesis, *J. Nat. Prod.* **61**, 1212-1215.
- [26] J. Zhou, L. Huang, D. L. Hachey, C. H. Chen and C. Aiken (2005). Inhibition of HIV-1 maturation via drug association with the viral Gag protein in immature HIV-1 particles, *J. Biol. Chem.* **280**, 42149-42155.
- [27] O. B. Kazakova, L.V. Rubanik, I. E. Smirnova, O. V. Savinova, A. V. Petrova, N. N. Poleschuk, E. F.Khusnutdinova, E. I. Boreko and Y. M. Kapustina (2019). Synthesis and in vitro activity of oleanane type derivatives against *Chlamydia trachomatis*, *Org. Commun.* **12**, 1169-175.
- [28] T. R. Vieira, L. C. A. Barbosa, C. R. A. Maltha, V. F. Paula and E. A. Nascimento (2004). Chemical constituents from *Melaleuca alternifolia* (Myrtaceae), *Quim. Nova* **27**, 536-539.
- [29] F. B. H. Ahmad, V. U. Hassan, Z. R. B and A. J. H. (1997). Betulinic and ursolic acid from seed of *Melaleuca cajuputi* (Myrtaceae). *Orient. J. Chem.* **13**, 231-232.
- [30] W.-H. Hui and M.-M. Li (1976). Neutral triterpenoids from *Melaleuca leucadendron*, *Phytochemistry* **15**, 563.
- [31] C. K. Lee and M. H. Chang (1999). Four new triterpenes from the heartwood of *Melaleuca leucadendron*, *J. Nat. Prod.* **62**, 1003-1005.
- [32] M. A. Abdel-Alim, F. A. Moharram, E. A. Aboutabl and S. F. El-Tohamy (1998). Constituents of *Melaleuca quinquenervia* Cav. grown in Egypt, *J. Herbs Spices Med. Plants* **5**, 29-37.
- [33] J. L. Courtney, E. V. Lassak and G. B. Speirs (1983). Leaf wax constituents of some Myrtaceous species, *Phytochemistry* **22**, 947-949.
- [34] C. S. Chopra, A. R. H. Cole, K. J. L. Theiberg, D. E. White and H. R. Arthur (1965). Triterpene compounds—VII: The constitution of melaleucic acid, *Tetrahedron* **21**, 1529-1536.

- [35] F. M. Abdel Bar (2019). Radical scavenging potential of compounds from *Melaleuca ericifolia* SM., *World J. Pharm. Res.* **8**, 110-123.
- [36] R. A. Moreau, L. Nyström, B. D. Whitaker, J. K. Winkler-Moser, D. J. Baer, S. K. Gebauer and K. B. Hicks (2018). Phytosterols and their derivatives: Structural diversity, distribution, metabolism, analysis, and health-promoting uses, *Prog. Lipid Res.* **70**, 35-61.
- [37] S. Rattanaburi, W. Mahabusarakam, S. Phongpaichit and A. R. Carroll (2013). A new chromone from the leaves of *Melaleuca cajuputi* Powell, *Nat. Prod. Res.* **27**, 221-225.
- [38] A. N. Panche, A. D. Diwan and S. R. Chandra (2016). Flavonoids: An overview, *J. Nutr. Sci.* **5**, e47.
- [39] D. Tungmunthum, A. Thongboonyou, A. Pholboon and A. Yangsabai (2018). Flavonoids and other phenolic compounds from medicinal plants for pharmaceutical and medical aspects: an overview, *Medicines (Basel)* **5**, e93.
- [40] L. Li, W. Du and W. Wang (2019). Evaluation of the antioxidant and anti-inflammatory effects of the extract of *Ribes mandshuricum* (Maxim.) Kom. leaves, *Rec. Nat. Prod.* **13**, 141-155.
- [41] O. Seligmann and H. Wagner (1981). Structure determination of melanerin, the first naturally occurring flavonoid of the triphenylmethane family, *Tetrahedron* **37**, 2601-2606.
- [42] C.-K. Lee (1999). Leucadenone A-D, the novel class flavanone from the leaves of *Melaleuca leucadendron* L., *Tetrahedron Lett.* **40**, 7255-7259.
- [43] G. M. Kitanov and I. Assenov (1992). Flavonols from *Melaleuca leucadendron* leaves, *Fitoterapia* **63** 379-380.
- [44] A. N. Hashim, N. F. Swilam, E. S. Moustafa, S. M. Bakry, R. M. Labib, H. H. Barakat, A. B. Singab, M. W. Linscheid and M. A. Nawwar (2018). A cytotoxic flavonol glycoside from *Melaleuca leucadendra* leaves extract with immunostimulant activity, *Pharmazie* **73**, 61-64.
- [45] E. Al-Sayed, N. M. El-Lakkany, S. H. Seif El-Din, A. N. Sabra and O. A. Hammam (2014). Hepatoprotective and antioxidant activity of *Melaleuca styphelioides* on carbon tetrachloride-induced hepatotoxicity in mice, *Pharm. Biol.* **52**, 1581-1590.
- [46] L. Yao, Y. Jiang, R. Singanusong, B. D'arcy, N. Datta, N. Caffin and K. Raymont (2004). Flavonoids in Australian *Melaleuca*, *Guioa*, *Lophostemon*, *Banksia* and *Helianthus* honeys and their potential for floral authentication, *Food Res. Int.* **37**, 166-174.
- [47] S. A. El-Toumy, M. S. Marzouk, F. A. Moharram and E. A. Aboutabl (2001). Flavonoids of *Melaleuca quinquenervia*, *Pharmazie* **56**, 94-95.
- [48] C. Li, H. Liu, L. Zhao, W. Zhang, S. Qiu, X. Yang and H. Tan (2017). Antibacterial neolignans from the leaves of *Melaleuca bracteata*, *Fitoterapia* **120**, 171-176.
- [49] E. Durães, C. Clementino, L. Fari, L. Ramos, M. Oliveira, J. Paula and P. L. Naves (2017). Phytochemical study, toxicity and antimicrobial activity of *Psidium myrsinites* DC. (Myrtaceae) leaves, *Biosci. J.* **33**, 1305-1313.
- [50] S. Sawada, H. Mori, Y. Sawanakunant, R. Nishida, Y. Yamamoto and T. Hosokawa (2000). Polysubstituted polyphenols of cajuput tree leaf and their anti-proliferative activity to cultured mouse T-lymphoma cells, EL4, *Bull. Kyoto Univ. Ed. Series B, Math. Nat. Sci.* **95/96**, 1-9.
- [51] L. Yao, Y. Jiang, R. Singanusong, N. Datta and K. Raymont (2005). Phenolic acids in Australian *Melaleuca*, *Guioa*, *Lophostemon*, *Banksia* and *Helianthus* honeys and their potential for floral authentication, *Food Res. Int.* **38**, 651-658.
- [52] F. A. Moharram, M. S. Marzouk, S. A. El-Toumy, A. A. Ahmed and E. A. Aboutabl (2003). Polyphenols of *Melaleuca quinquenervia* leaves--pharmacological studies of grandinin, *Phytother. Res.* **17**, 767-773.
- [53] T. Yoshida, Y. Amakura and M. Yoshimura (2010). Structural features and biological properties of ellagitannins in some plant families of the order Myrtales, *Int. J. Mol. Sci.* **11**, 79-106.
- [54] T. H. Lee, G. J. Wang, C. K. Lee, Y. H. Kuo and C. H. Chou (2002). Inhibitory effects of glycosides from the leaves of *Melaleuca quinquenervia* on vascular contraction of rats, *Planta Med.* **68**, 492-496.
- [55] T. Yoshida, H. Ito, M. Yoshimura, K. Miyashita and T. Hatano (2008). C-Glucosidic ellagitannin oligomers from *Melaleuca squarrosa* Donn ex Sm., Myrtaceae, *Phytochemistry* **69**, 3070-3079.
- [56] E. Al-Sayed, M. M. Abdel-Daim and M. A. Khatib (2019). Hepatoprotective activity of praecoxin A isolated from *Melaleuca ericifolia* against carbon tetrachloride-induced hepatotoxicity in mice. Impact on oxidative stress, inflammation, and apoptosis, *Phytother. Res.* **33**, 461-470.
- [57] O. Celaj, A. G. Durán, P. Cennamo, M. Scognamiglio, A. Fiorentino, A. Esposito and B. D'abrosca (2020). Phloroglucinols from Myrtaceae: attractive targets for structural characterization, biological properties and synthetic procedures, *Phytochem. Rev.* 1-41. doi:10.1007/s11101-020-09697-2
- [58] B. P. Naidu (2000). Accumulation of proline analogues and adaptation of *Melaleuca* species to diverse environments in Australia, *Aust. J. Bot.* **48**, 611-620.
- [59] G. P. Jones, B. P. Naidu, L. G. Paleg, E. R. T. Tiekink and M. R. Snow (1987). 4-Hydroxy-N-methylproline analogues in *Melaleuca* spp., *Phytochemistry* **26**, 3343-3344.

- [60] A. Poljakoff-Mayber, D. E. Symon, G. P. Jones, B. P. Naidu and L. G. Paleg (1987). Nitrogenous compatible solutes in native South Australian plants, *Funct. Plant Biol.* **14**, 341-350.
- [61] J.-L. Ríos (2016). Chapter 1 - Essential Oils: What They Are and How the Terms Are Used and Defined, In: Book, Academic Press, San Diego. 3-10.
- [62] Y. Bi, J. Xu, F. Sun, X. Wu, W. C. Ye, Y. Sun, Q. Meng and W. Huang (2013). Synthesis and antitumor activity of 17-carboxylic acid modified amide derivatives of 23-hydroxy betulinic acid, *Rec. Nat. Prod.* **7**, 254-265.
- [63] N. M. Al-Abd, Z. Mohamed Nor, M. Mansor, F. Azhar, M. S. Hasan and M. Kassim (2015). Antioxidant, antibacterial activity, and phytochemical characterization of *Melaleuca cajuputi* extract, *BMC Complement. Altern. Med.* **15**, e385.
- [64] N. M. Al-Abd, Z. M. Nor, M. Mansor, M. S. Hasan and M. Kassim (2016). Antifilarial and antibiotic activities of methanolic extracts of *Melaleuca cajuputi* flowers, *Korean J. Parasitol.* **54**, 273-280.
- [65] U. Ukit, A. Widiana, E. Rahmawati and R. M. Hasby (2019). Antibacterial activities test of cajuput leaf waste extract (*Melaleuca cajuputi* Powell) on pathogenic bacteria, *J. Phys. Conf. Ser.* **1402**, 033030.
- [66] B. Laribi, A. Ismail, T. Bettaeib and L. Hamrouni (2020). Phytochemical evaluation and *in vitro* antifungal activities of *Melaleuca styphelioides* leaves: comparison between volatile and non-volatile extracts, *Plant Biosyst.* 1-21. doi: 10.1080/11263504.2020.1727986
- [67] D. Ganesan, E. Al-Sayed, A. Albert, E. Paul, A. N. B. Singab, S. Govindan Sadasivam and L. Saso (2018). Antioxidant activity of phenolic compounds from extracts of *Eucalyptus globulus* and *Melaleuca styphelioides* and their protective role on D-glucose-induced hyperglycemic stress and oxalate stress in NRK-49Fcells, *Nat. Prod. Res.* **32**, 1274-1280.
- [68] F. Albouchi, R. Avola, G. M. L. Dico, V. Calabrese, A. C. E. Graziano, M. Abderrabba and V. Cardile (2018). *Melaleuca styphelioides* Sm. polyphenols modulate interferon gamma/histamine-induced inflammation in human NCTC 2544 keratinocytes, *Molecules* **23**, e2526.
- [69] L. S. Bianchini Silva, F. B. Perasoli, K. V. Carvalho, K. M. Vieira, M. T. Paz Lopes, G. H. Bianco De Souza, O. D. Henrique Dos Santos and K. M. Freitas (2020). *Melaleuca leucadendron* (L.) L. flower extract exhibits antioxidant and photoprotective activities in human keratinocytes exposed to ultraviolet B radiation, *Free Radic. Biol. Med.* **159**, 54-65.
- [70] E. Al-Sayed, H. E. Michel, M. A. Khatlab, M. El-Shazly and A. N. Singab (2020). Protective role of casuarinin from *Melaleuca leucadendra* against ethanol-induced gastric ulcer in rats, *Planta Med.* **86**, 32-44.
- [71] N. P. Nirmal, D. Webber, R. Mereddy and Y. Sultanbawa (2018). Biochemical and functional properties of indigenous Australian herbal infusions, *Food Biosci.* **26**, 133-138.
- [72] F. O. Osunsanmi, G. E. Zharare, R. Mosa, M. I. Ikhile, F. O. Shode and A. J. T. J. O. P. R. Opoku (2019). Anti-oxidant, anti-inflammatory and antiacetylcholinesterase activity of betulinic acid and 3 $\beta$ -acetoxybetulinic acid from *Melaleuca bracteata* 'Revolution Gold', *Trop. J. Pharm. Res.* **18**, 303-309.
- [73] H. Younus (2018). Therapeutic potentials of superoxide dismutase, *Int. J. Health. Sci.* **12**, 88-93.
- [74] E. Al-Sayed, H. A. Hamid and H. M. Abu El Einin (2014). Molluscicidal and antischistosomal activities of methanol extracts and isolated compounds from *Eucalyptus globulus* and *Melaleuca styphelioides*, *Pharm. Biol.* **52**, 698-705.
- [75] F. Osunsanmi, G. Zaharare, B. Oyinloye, R. Mosa, M. Ikhile, F. Shode, I. Ogunyinka and A. Opoku (2018). Antithrombotic, anticoagulant and antiplatelet activity of betulinic acid and 3 $\beta$ -acetoxybetulinic acid from *Melaleuca bracteata* 'Revolution Gold' (Myrtaceae) Muell leaf, *Trop. J. Pharm. Res.* **17**, 1983-1989.
- [76] E. Al-Sayed and A. Esmat (2016). Hepatoprotective and antioxidant effect of ellagitannins and galloyl esters isolated from *Melaleuca styphelioides* on carbon tetrachloride-induced hepatotoxicity in HepG2 cells, *Pharm. Biol.* **54**, 1727-1735.
- [77] J. Adesanwo, F. Shode, O. Aiyelaagbe, O. Rabiun and R. Oyede (2009). Antisecretory and antiulcerogenic activities of the stem bark extract of *Melaleuca bracteata* and isolation of principles, *J. Med. Plant Res.* **3**, 822-824.
- [78] S. Subehan, T. Usia, H. Iwata, S. Kadota and Y. Tezuka (2006). Mechanism- based inhibition of CYP3A4 and CYP2D6 by Indonesian medicinal plants, *J. Ethnopharmacol.* **105**, 449-455.