

Potassium ferrocyanide promoted an efficient synthesis of benzoxazoles and benzothiazoles under solvent free condition

Vishal S. Patil^{1*}, Dhanshri V. Patil² and Sachin S. Potdar³

¹Department of Chemistry, Sanjeevan Engineering and Technology Institute, Panhala, Kolhapur, Maharashtra, India, 416201

²Department of Chemistry, Krishna Mahavidyalaya, Rethare Bk. Karad, Maharashtra, India, 415108

³Department of Physics, Sanjeevan Engineering and Technology Institute, Panhala, Kolhapur, Maharashtra, India, 416201

(Received December 03, 2021; Revised March 20, 2022; Accepted March 21, 2022)

Abstract: In the family of heterocycles that includes benzoxazoles and benzothiazoles, there exist compounds with a wide range of biological activity. Because of this characteristic, we designed a moderate and effective technique for the synthesis of 2-substituted benzoxazole and benzothiazole using condensation of aldehyde and 2-aminophenol or 2-aminothiophenol via oxidation of carbon-nitrogen bond. Potassium ferrocyanide catalyzed one-pot synthesis is efficient and provides for quick reaction times, simple set-up and high yields. As a result, we provide here a technique for the rapid solvent free synthesis of benzoxazoles and benzothiazoles. Some synthesized products were identified by ¹H-NMR, ¹³C-NMR and MASS. The role of potassium ferrocyanide as a catalyst is represented by plausible reaction mechanism.

Keywords: Aldehyde; potassium ferrocyanide; benzoxazoles; benzothiazoles; solvent free.

©2022 ACG Publication. All right reserved.

1. Introduction

Benzoxazoles and benzothiazoles are frequent heterocyclic scaffolds in physiologically active and pharmaceutically relevant chemicals and they belong to a large family of molecules. Benzoxazoles are essential scaffolds in natural compounds¹⁻² and drug development³⁻⁵. Benzoxazole compounds with appropriate substitutions have been shown to exhibit a variety of medicinal properties including antibacterial activity⁶, antimicrobial⁷⁻¹⁰, antiviral¹¹, topoisomerase I, II inhibitory¹², antitumor activities¹³, anticancer agent¹⁴⁻¹⁵ NSC-693638, L-697,661, antiviral¹⁶ and antibacterial¹⁷ UK-1, AJ19561. According to recent research, substituted 2-benzylbenzoxazoles exhibit antibacterial, antifungal¹⁸, antimicrobial¹⁹⁻²¹ and anti-measles virus²² properties (Figure 1).

The tiny and simple benzothiazole nucleus is found in compounds with intriguing biological properties such as anticonvulsant²³⁻²⁴, antimalarial²⁵, antitubercular²⁶, antimicrobial²⁷⁻²⁸, antitumour²⁹⁻³², anthelmintic³³, anti-inflammatory, analgesic properties³⁴. The benzothiazole ring may be found in a variety of natural substances, both marine and terrestrial, that have significant biological activity. Many natural products, such as epothilone-A, lyngbyabellin A, dolastatin 10 & bleomycin, include thiazole nucleus molecules³⁵. The synthesis of these molecules is of significant interest due to their substantial medicinal value. Riluzole is a benzothiazole derivative-containing medication used to treat amyotrophic lateral sclerosis. In certain patients, it may postpone the need for a tracheostomy or a ventilator and it

*Corresponding author: E-Mail: vishalpatil.chem@gmail.com

Synthesis of benzoxazoles and benzothiazoles

can extend life by 3–5 months. Benzothiazoles and benzoxazoles are heterocyclic compounds with a wide range of biological characteristics. (Figure 1)³⁶⁻⁴¹.




Figure 1. Some biologically active benzoxazole and benzothiazoles

However, a variety of techniques for synthesizing benzoxazoles⁴²⁻⁴⁶ and benzothiazoles⁴⁷⁻⁵² are now available, there is still a need for novel and easy processes that enable a broad range of structural diversity and substitution patterns in the target library.

The condensation of o-aminobenzenethiol or o-aminophenol with substituted aldehydes⁵³, nitriles⁵⁴, acyl chlorides⁵⁵ or carboxylic acids⁵⁶⁻⁵⁸ is the most frequently used synthesis technique for obtaining benzoxazole and benzothiazole. These techniques often require lengthy response times and harsh circumstances. An example from recent times is the synthesis of benzoxazoles and benzothiazoles from alcohol⁵⁹. However, several of these techniques have one or more disadvantages, such as high acidic conditions, lengthy reaction durations, poor product yields, laborious workup, the requirement for excessive quantities of reagent and the use of hazardous reagents, catalysts or solvents. As a result, there is a significant need for a highly efficient and ecologically friendly technique of synthesizing these heterocycles. Thus, developing an effective and easy chemical process or technique for the synthesis of physiologically active molecules from a simple reagent is always a difficult challenge for chemists working in the area of organic synthesis. Thus, the superiority of this research article is we overcome all the above problems to synthesize benzoxazoles and benzothiazoles by using potassium ferrocyanide as an environment friendly catalyst.

$\text{K}_4[\text{Fe}(\text{CN})_6]$ has recently played an important role as a catalyst in the production of the anti-Alzheimer medication (-) Galanthamine⁶⁰. Because of its high stability, oxidising power selectivity and harmless byproduct $\text{Fe}(\text{III})$, it also facilitated oxidative cyclization of 5-S Cysteinyl-dopa⁶¹. Yu and Gu investigated the release of cyanide into the environment, which has terrestrial implications for ecosystems⁶². Other researchers are interested in $\text{K}_4[\text{Fe}(\text{CN})_6]$ because Gaffar and Abu-El Fadl investigated the kinetics of potassium ferro cyanide⁶³ since it has several benefits such as good solubility in water, simple handling, low cost, eco-friendliness, availability and strong reactivity.

2. Experimental

2.1 General Procedure for the Synthesis of Benzoxazoles and Benzothiazoles on Grinding

At room temperature, a mortar with a pestle was used to grind a combination of 2-aminophenol or 2-aminothiophenol (1 mmol), aldehyde (1 mmol) and potassium ferro-cyanide (10 mol%). TLC was used to track the progress of the reaction. The crude product was washed with water, dried and recrystallized with ethanol once the reaction was completed (< 2 min).

2.2 General Experimental Methods

On a Stuart-SMP10 melting point apparatus, the melting points of the prepared compounds were measured in open-glass capillaries. IR absorption spectra were recorded on a Perkin Elmer 1650 FTIR using KBr pellets in the range of 4000-450 cm^{-1} . A Bruker spectrometer running at 300 MHz was used to record ^1H -NMRs. The chemical shifts in ^1H -NMR are given as parts per million (ppm) downfield from the internal standard TMS (Me_4Si). The chemical shifts (scale) are given in parts per million in the ^{13}C NMR spectra obtained at 75 MHz (ppm). On an LCQ ion trap mass spectrometer, mass spectra were captured. Purity of the compounds were checked by thin layer chromatography (TLC) on Merck silica gel 60 F254 pre-coated sheets in benzene/methanol mixture and spots were developed using iodine vapor as visualizing agents.

2-phenyl-1,3-benzoxazole⁶⁵(a): White solid, mp: 102 °C. IR: (KBr, cm^{-1}) ν 3019.50, 1634.07, 1526.79, 1453.36, 1225.91, 1054.34; ^1H NMR: (CDCl_3 , 300 MHz), δ : 8.18 (d, 2H), 7.76-7.72 (m, 2H), 7.67-7.60 (m, 3H), 7.40-7.36 (d, 2H) ppm; ^{13}C NMR: (CDCl_3 , 75 MHz), δ : 161.4, 151.4, 138.5, 132.7, 128.6, 128, 127.8, 127.6, 124.3, 119.9, 110.6; MS (ES): Calculated for $\text{C}_{13}\text{H}_9\text{NO}$ m/z : 195.21, found 195.10.

2-(4-bromophenyl)-1,3-benzoxazole⁶⁵(b): White solid, mp: 158 °C.

2-(4-methylphenyl)-1,3-benzoxazole⁶⁶(c): Light brown solid, mp: 113 °C. ^1H NMR: (CDCl_3 , 300 MHz), δ : 8.10-8.08 (d, 2H), 7.78 (t, 2H), 7.41-7.39 (m, 4H), 2.41 (s, 3H) ppm; ^{13}C NMR: (CDCl_3 , 75 MHz) δ : 161.4, 151.4, 141.6, 138.5, 129, 127.6, 126.5, 124.3, 119.4, 110.6, 21.5; MS (ES): Calculated for $\text{C}_{14}\text{H}_{11}\text{NO}$ m/z : 209.24, found 209.16.

2-(4-chlorophenyl)-1,3-benzoxazole⁶⁵(d): White solid, mp: 148 °C. ^1H NMR: (CDCl_3 , 300 MHz), δ : 8.21-8.26 (d, 2H), 7.53-7.57 (d, 2H), 7.80-7.81 (m, 1H), 7.61, 7.63 (m, 1H), 7.40-7.43 (m, 2H) ppm; ^{13}C NMR: (CDCl_3 , 75 MHz) δ : 161.4, 151.4, 138.5, 137.3, 129.3, 128.3, 127.6, 124.3, 119.9, 110.6; MS (ES): Calculated for $\text{C}_{13}\text{H}_8\text{ClNO}$ m/z : 229.66, found 229.46.

2-[3-fluoro-4-(trifluoromethyl)phenyl]-1,3-benzoxazole(e): White Sticky Semi-solid. IR: (KBr, cm^{-1}) ν 3253.19, 2923.60, 1641.42, 1501.51, 1258.21, 1134.63, 1019.10; ^1H NMR: (CDCl_3 , 300 MHz), δ : 7.76-7.69 (m, 3H), 7.61 (d, 1H), 7.40-7.36 (m, 2H) ppm; ^{13}C NMR: (75 MHz, CDCl_3) δ : 162.7, 162.7, 162.7, 161.5, 160.7, 160.6, 151.3, 138.5, 132, 132.9, 128.1, 127.6, 125.1, 124.3, 121.7, 119.6; MS (ES): Calculated for $\text{C}_{14}\text{H}_7\text{F}_4\text{NO}$ m/z : 281.20, found 281.10.

2-(4-methoxyphenyl)-1,3-benzoxazole⁶⁵(f): White solid, mp: 99 °C.

5-chloro-2-(4-nitrophenyl)-1,3-benzoxazole⁶⁹(g): White Sticky Semi-solid. IR: (KBr, cm^{-1}) ν 3025.54, 2871.9, 1641.32, 1503.51, 1134.63, 1020.10, 748.10; ^1H NMR: (CDCl_3 , 300 MHz), δ : 8.35 (d, 2H), 8.22 (d, 2H), 7.81 (s, 1H), 7.75 (d, 1H), 7.40 (d, 1H) ppm; ^{13}C NMR: (75 MHz, CDCl_3) δ : 161.5, 150, 148.9, 137.8, 134.6, 129.1, 128.9, 126.3, 124.2, 117.4, 111.7; MS (ES): Calculated for $\text{C}_{13}\text{H}_7\text{ClN}_2\text{O}_3$ m/z : 274.65, found 274.40.

Synthesis of benzoxazoles and benzothiazoles

5-chloro-2-[3-fluoro-4-(trifluoromethyl)phenyl]-1,3-benzoxazole(h): White Sticky Semi-solid. ^1H NMR: (CDCl_3 , 300 MHz) δ : 7.15-7.49 (m, 6H) ppm; ^{13}C NMR: (75 MHz, CDCl_3), δ : 160.4, 151.6, 148.1, 141.5, 130.7, 127.4, 122.9, 120.9, 118, 114.3, 112.2, 108.5 MS (ES): Calculated for $\text{C}_{14}\text{H}_6\text{ClF}_4\text{NO}$ m/z : 315.65 found 315.38.

5-chloro-2-phenyl-1,3-benzoxazole⁶⁷(i): Yellow solid, mp: 111 °C. ^1H NMR: (CDCl_3 , 300 MHz), δ : 8.31-8.33 (d, 2H), 7.63 (s, 1H), 7.56-7.58 (m, 3H), 7.50-7.51 (d, 1H), 7.21-7.22 (d, 1H) ppm; ^{13}C NMR: (CDCl_3 , 75 MHz) δ : 161.5, 150, 137.8, 132.7, 129.1, 128, 127.8, 126.3, 117.4, 111.7; MS (ES): Calculated for $\text{C}_{13}\text{H}_8\text{ClNO}$ m/z : 229.66, found 229.40.

2-(4-methylphenyl)-5-nitro-1,3-benzoxazole(j): Yellow solid, mp: 192 °C. ^1H NMR: ($\text{DMSO}-d_6$, 300 MHz), δ : 8.58 (d, 1H), 7.85 (m, 1H), 7.78 (d, 2H), 7.32 (m, 1H), 7.23 (m, 2H), 2.40 (d, 3H) ppm; ^{13}C NMR: (CDCl_3 , 75 MHz) δ : 161.5, 151.3, 141.6, 141.2, 139.4, 129, 126.3, 120.8, 119.2, 110.9, 21.1; MS (ES): Calculated for $\text{C}_{14}\text{H}_{10}\text{N}_2\text{O}_3$ m/z : 254.24, found 254.18.

2-(4-nitrophenyl)-5-nitro-1,3-benzoxazole⁷³(k): White solid, mp: 262 °C.

2-phenyl-1,3-benzothiazole⁶⁵(l): Yellow solid, mp: 112 °C. IR: (KBr, cm^{-1}) ν : 3246.73, 2923.64, 1507.77; ^1H NMR: (CDCl_3 , 300 MHz), δ : 8.19 (d, 1H), 8.04-8.00 (m, 3H), 7.54-7.48 (m, 5H) ppm; ^{13}C NMR: (CDCl_3 , 75 MHz) δ : 166.6, 151.8, 136, 134.9, 132.1, 129.3, 127.3, 126.3, 126, 123.5, 123.2; MS (ES): Calculated for $\text{C}_{13}\text{H}_9\text{NS}$ m/z : 211.28, found 212.20.

2-(4-methylphenyl)-1,3-benzothiazole⁶⁷(m): Yellow solid, mp: 88 °C. ^1H NMR: (CDCl_3 , 300 MHz) δ : 8.29–7.18 (m, 8H), 2.64 (s, 3H) ppm; ^{13}C NMR: (CDCl_3 , 75 MHz) δ : 166.6, 151.8, 139, 136, 132.7, 129.3, 127.9, 126, 123.5, 123.2, 21.2; MS (ES): Calculated for $\text{C}_{14}\text{H}_{11}\text{NS}$ m/z : 225.30, found 225.20.

2-(4-methoxyphenyl)-1,3-benzothiazole⁶⁸(n): Yellow solid, mp: 123 °C.

4-(1,3-benzothiazol-2-yl)phenol⁷⁰(o): Yellow solid, mp: 225 °C. IR: (KBr, cm^{-1}) ν : 3404.14, 2928.27, 2845.32, 1626.86, 1491.60, 1021.10; ^1H NMR: (CDCl_3 , 300 MHz) δ : 8.17 (d, 1H), 8.02 (d, 1H), 7.73 (d, 2H), 7.54-7.48 (m, 2H), 6.86 (d, 2H), 5.13 (s, 1H) ppm; ^{13}C NMR: (CDCl_3 , 75 MHz) δ : 166.6, 161.4, 151.8, 136, 130, 126.3, 126, 123.6, 123.5, 123.2, 116.7; MS (ES): Calculated for $\text{C}_{13}\text{H}_9\text{NOS}$ m/z : 227.28, found 227.17

2-(4-bromophenyl)-1,3-benzothiazole⁷²(p): Yellow solid, mp: 133 °C.

2-(4-chlorophenyl)-1,3-benzothiazole⁷¹(q): White solid, mp: 117 °C.

2-(2-chlorophenyl)-1,3-benzothiazole⁶⁸(r): White solid, mp: 70 °C. ^1H NMR: (CDCl_3 , 300 MHz) δ : 8.17 (d, 1H), 8.03 (d, 1H), 7.73 (d, 1H), 7.54-7.48 (m, 3H), 7.39-7.33 (m, 2H) ppm; ^{13}C NMR: (75 MHz, CDCl_3) δ : 166.1, 151.8, 136, 133.9, 132.3, 131.3, 130, 128.2, 128.1, 126.3, 126, 123.5, 123.2; MS (ES): Calculated for $\text{C}_{13}\text{H}_8\text{ClNS}$ m/z : 245.72, found 247.2.


2-[3-fluoro-4-(trifluoromethyl)phenyl]-1,3-benzothiazole(s): White solid, mp: 185 °C. ^1H NMR: (CDCl_3 , 300 MHz), δ : 8.13 (s 1H, Ar-H), 8.11 (d, 1H, $J = 8.5$ Hz, Ar-H), 7.95 (d, 1H, $J = 8$ Hz, Ar-H), 7.27–7.53 (m, 4H, Ar-H) ppm; ^{13}C NMR: (75 MHz, CDCl_3), δ : 165.7, 163.4, 161.4, 161.3, 151.8, 141.5, 141.4, 136, 128.5, 126, 125, 124.7, 123.9, 121.8, 121.2, 121.1, 119.6, 119.6; MS (ES): Calculated for $\text{C}_{13}\text{H}_8\text{ClNS}$ m/z : 297.27, found 297.20.

2-(4-nitrophenyl)-1,3-benzothiazole⁶⁸(**t**): Yellow solid, mp: 229 °C. IR: (KBr, cm⁻¹) ν 3120.46, 1622.97, 1595.78, 1470.85, 1188.49; ¹H NMR: (CDCl₃, 300 MHz), δ : 8.27 (d, 2H), 8.18 (s, 1H), 8.12 (d, 2H), 8.02 (d, 1H), 7.55-7.49 (m, 2H) ppm; ¹³C NMR: (75 MHz, CDCl₃) δ : 166.6, 151.8, 148.9, 142.3, 136, 127.8, 126.3, 126, 124.9, 123.5, 123.2; MS (ES): Calculated for C₁₃H₈ClNS m/z : 256.28, found 256.16.

2-(4-florophenyl)-1,3-benzothiazole⁷²(**u**): White solid, mp: 99 °C.

3. Results and Discussion

We present an effective and ecologically friendly approach for synthesizing benzoxazole and benzothiazole under solvent-free conditions using catalytic quantities of K₄[Fe(CN)₆].









Scheme 1. K₄[Fe(CN)₆] catalyzed synthesis of benzoxazoles on grinding

To determine the best reaction conditions, o-aminophenol was treated with benzaldehyde in the presence of K₄[Fe(CN)₆] (Scheme 1, Table 1). The best molar ratio of o-aminophenol:aldehyde at room temperature under solvent-free conditions is 1:1 and 2-phenylbenzoxazole was produced in 94 percent yield after 2 minutes under these circumstances (Entry 1, a, Table 1). To establish the involvement of K₄[Fe(CN)₆], the same reaction was carried out at ambient temperature in the absence of a catalyst and no product production was seen after 20 minutes. These findings imply that K₄[Fe(CN)₆] has a high catalytic activity in this reaction. The procedure's generality was next tested using reactions of different aldehydes with o-aminophenols. The result demonstrated that the reaction completed within 2-3 minutes with excellent yield of the products (Entry 1-3, Table 1).

Table 1. Synthesis of benzoxazoles and benzothiazoles on grinding


Entry	Substituted 2-aminophenol/ aminothiophenol	Aldehydes	Product	Yield ^a %
1			 R = a. H; b. 4-Br; c. 4-CH ₃ ; d. 4-Cl; e. 3-F, 4-CF ₃ ; f. 4-OCH ₃	a. 94 b. 92 c. 90 d. 92 e. 93 f. 87
2			 R = g. 4-NO ₂ ; h. 3-F, 4-CF ₃ ; i. H	g. 96 h. 88 i. 90

Synthesis of benzoxazoles and benzothiazoles


3				j. 94
				k. 92
4				l. 92
				m. 90
				n. 89
				o. 90
				p. 92
				q. 94
				r. 91
				s. 94
				t. 95
				u. 90

R = j. 4-CH₃; k. 4-NO₂R = l. H; m. 4-CH₃; n. 4-OCH₃; o. 4-OH; p. 4-Br; q. 4-Cl; r. 2-Cl; s. 3-F, 4-CF₃; t. 4-NO₂; u. 4-F^a Isolated yield of the products

This protocol also extended by the reaction of various aldehydes with o-aminothiophenol (Scheme 2). 2-aminothiophenol was grind with a variety of aldehydes at room temperature in K₄[Fe(CN)₆] catalysis under solvent free condition (Entry 4, Table 1). All the reactions proceed to completion in just 2-3 min. at room temperature without any organic solvent. In all instances, the corresponding benzothiazoles could be recovered with high yields (Entry 4, Table 1).

**Scheme 2** K₄[Fe(CN)₆] catalyzed synthesis of benzothiazoles on grinding

Scheme 3 depicts one proposed mechanism for this process. The K₄[Fe(CN)₆] increases the electrophilic character of the carbonyl carbon, allowing for the nucleophilic addition of 2-aminothiophenol or 2-aminophenol to provide an intermediate I, which upon cyclisation followed by oxidation produces the desired product.

**Scheme 3.** Proposed mechanism for the synthesis of benzoxazoles and benzothiazoles

4. Conclusion

Potassium ferro-cyanide was discovered to be a gentle and effective catalyst for the production of benzoxazoles and benzothiazoles. This technique is feasible, environmentally benign and economically appealing due to the utilization of this affordable, readily accessible and reusable catalyst under solvent-free conditions. The proposed approach also has the benefits of a simple work-up process, moderate reaction conditions, extremely short reaction periods, excellent product yields and non-toxicity of the catalyst.

Supporting Information

Supporting information accompanies this paper on <http://www.acgpubs.org/journal/organic-communications>

ORCID ID

Vishal S. Patil: [0000-0003-3911-5107](https://orcid.org/0000-0003-3911-5107)

Dhanshri V. Patil: [0000-0002-5305-7719](https://orcid.org/0000-0002-5305-7719)

Sachin S. Potdar: [0000-0003-4409-8364](https://orcid.org/0000-0003-4409-8364)

References

- [1] Chaney, M. O.; Demarco, P. V.; Jones, N. D.; Occolowitz, J. L. Structure of A23187, a divalent cation ionophore. *J. Am. Chem. Soc.* **1974**, 96, 1932-1933.
- [2] Sato, S.; Kajiura, T.; Noguchi, M.; Takehana, K.; Kobayasho, T.; Tsuji, T. AJI9561, a new cytotoxic benzoxazole derivative produced by streptomyces sp. *J. Antibiot.* **2001**, 54, 102-104.
- [3] Brown, R. N.; Cameron, R.; Chalmers, D. K.; Hamilton, S.; Luttick, A.; Krippner, G. Y.; McConnell, D. B.; Nearn, R.; Stanislawski, P. C.; Tucker, S. P.; Watson, K. G. 2-Ethoxybenzoxazole as a bioisosteric replacement of an ethyl benzoate group in a human rhinovirus (HRV) capsid binder. *Bioorg. Med. Chem. Lett.* **2005**, 15, 2051-2055
- [4] Manas, E. S.; Unwalla, R. J.; Xu, Z. B.; Malamas, M. S.; Miller, C. P.; Harris, H. A.; Hsiao, C.; Akopian, T.; Hum, W. T.; Malakian, K.; Wolfrom, S.; Bapat, A.; Bhat, R. A.; Stahl, M. L.; Somers, W. S.; Alvarez, J. C. Structure-based design of estrogen receptor- β selective ligands. *J. Am. Chem. Soc.* **2004**, 126, 15106-15119.
- [5] Malamas, M. S.; Manas, E. S.; McDevitt, R. E.; Gunawan, I.; Xu, Z. B.; Collini, M. D.; Miller, C. P.; Dinh, T.; Henderson, R. A.; Keith, J. C.; Harris, H. A. Design and synthesis of aryl diphenolic azoles as potent and selective estrogen receptor- β ligands. *J. Med. Chem.* **2004**, 47, 5021.
- [6] Prudhomme, M.; Guyot, J.; Jeminet, G. Semi-synthesis of a23187 (calcimycin) analogs. *J. Antibiotics* **1986**, 39, 934-937.
- [7] Ren, I. O.; Temiz, O.; Yalcin, I.; Sener, E. Akin, A.; Ucarturk N. Synthesis and microbiological activity of 5(or 6)-methyl-2-substituted benzoxazole and benzimidazole derivatives. *Arzneim Forsch. Drug Res.* **1997**, 47, 1393-1397.
- [8] Ren, I. O.; Temiz, O.; Yalcin I.; Aki-Sener, E.; Altanlar, N. Synthesis and antimicrobial activity of some novel 2,5- and/or 6-substituted benzoxazole and benzimidazole derivatives. *Eur. J. Pharm. Sci.* **1999**, 7, 153-160.
- [9] Temiz-Arpaci, O.; Ozdemir, A.; Yalcin, I.; Yildiz, I.; Aki-Sener, E.; Altanlar, N. Synthesis and antimicrobial activity of some 5-[2-(Morpholin-4-yl)acetamido] and/or 5-[2-(4-Substituted piperazin-1-yl)acetamido]-2-(p-substituted phenyl)benzoxazoles. *Arch. Pharm.* **2005**, 338, 105-111.
- [10] Vinsova, J.; Horak, V.; Buchta, V.; Kaustova, J. Highly lipophilic benzoxazoles with potential antibacterial activity. *Molecules* **2005**, 10, 783-793.
- [11] Song, X.; Vig, B. S.; Lorenzi, P. L.; Drach, J. C.; Townsend, L. B.; Amidon, G. L. Amino acid ester prodrugs of the antiviral agent 2-bromo-5,6-dichloro-1-(β -D-ribofuranosyl)benzimidazole as potential substrates of hPEPT1 transporter. *J. Med. Chem.* **2005**, 48, 1274-1277.
- [12] Pinar, A.; Yurdakul, P.; Yildiz, I.; Temiz-Arpaci, O.; Acan, N. L.; Aki-Sener, E.; Yalcin, I. Some fused heterocyclic compounds as eukaryotic topoisomerase II inhibitors. *Biochem. Biophys. Res. Commun.* **2004**, 317, 670-674.

Synthesis of benzoxazoles and benzothiazoles

- [13] Ueki, M.; Taniguchi, M. J. UK-1, a novel cytotoxic metabolite from *Streptomyces* sp. 517-02. III. antibacterial action of demethyl UK-1. *Antibiotics* **1997**, *50*, 788–790.
- [14] McKee, M. L.; Kerwin, S. M. Synthesis, metal ion binding and biological evaluation of new anticancer 2-(2'-hydroxyphenyl)benzoxazole analogs of UK-1. *Bioorg. Med. Chem.* **2008**, *16*, 1775.
- [15] Oksuzoglu, E.; Temiz-Arpaci, O.; Tekiner-Gulbas, B.; Eroglu, H.; Sen, G.; Alper, S.; Yildiz, I.; Diril, N.; Aki-Sener, E.; Yalcin, I. A study on the genotoxic activities of some new benzoxazoles. *Med. Chem. Res.* **2008**, *16*, 1.
- [16] Gong, B.; Hong, F.; Kohm, C.; Bonham, L.; Klein, P. Synthesis and SAR of 2-arylbenzoxazoles, benzothiazoles and benzimidazoles as inhibitors of lysophosphatidic acid acyltransferase- β . *Bioorg. Med. Chem. Lett.* **2004**, *14*, 1455.
- [17] Sun, L.-Q.; Chen, J.; Bruce, M.; Deskus, J. A.; Epperson, J. R.; Takaki, K.; Johnson, G.; Iben, L.; Mahle, C. D.; Ryan, E.; Xu, C. Synthesis and structure–activity relationship of novel benzoxazole derivatives as melatonin receptor agonists. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 3799.
- [18] Ertan, T.; Yildiz, I.; Tekiner-Gulbas, B.; Bolelli, K.; Temiz-Arpaci, O.; Yalcin, I.; Aki, E.; Ozkan, S.; Kaynak, F. Synthesis, biological evaluation and 2D-QSAR analysis of benzoxazoles as antimicrobial agents. *Eur. J. Med. Chem.* **2009**, *44*, 501.
- [19] Alper-Hayta, S.; Arisoy, M.; Temiz-Arpaci, O.; Yildiz, I.; Aki, E.; Özkan, S.; Kaynak, F. Synthesis, antimicrobial activity, pharmacophore analysis of some new 2-(substitutedphenyl/benzyl)-5-[(2-benzofuryl)carboxamido]benzoxazoles. *Eur. J. Med. Chem.* **2008**, *43*, 2568.
- [20] Tekiner-Gulbas, B.; Temiz-Arpaci, O.; Yildiz, I.; Altanlar, N. Synthesis and *in vitro* antimicrobial activity of new 2-[p-substituted-benzyl]-5-[substituted-carbonylamino] benzoxazoles. *Eur. J. Med. Chem.* **2007**, *42*, 1293.
- [21] Yildiz-Oren, I.; Tekiner-Gulbas, B.; Yalcin, I.; Temiz-Arpaci, O.; Aki-Sener, E.; Altanlar, N. Synthesis and antimicrobial activity of new 2-[p-Substituted-benzyl]-5-[substituted-carbonylamino]benzoxazoles. *Arch. Pharm.* **2004**, *337*, 402.
- [22] Sun, A.; Prussia, A.; Zhan, W.; Murray, E. E.; Doyle, J.; Cheng, L.-T.; Yoon, J. J.; Radchenko, E. V.; Palyulin, V. A.; Compans, R. W.; Liotta, D. C.; Plemper, R. K.; Snyder, J. A. Nonpeptide inhibitors of measles virus entry. *J. Med. Chem.* **2006**, *49*, 5080.
- [23] Chakole, R. D.; Amnerkar, N. D.; Khedekar, P. B.; Bhusari, K. P. Synthesis of substituted benzothiazole derivatives of thioquinazoline as anticonvulsant agents. *Ind. J. Heterocyc. Chem.* **2005**, *15*, 27-30.
- [24] Alam, M.; Siddiqui, N. Synthesis of new benzothiazole incorporated sulphonamides as potential anticonvulsants. *Ind. J. Heterocyc. Chem.* **2004**, *13*, 361-364.
- [25] Burger, A.; Sawhey, S. N. Antimalarials. III. benzothiazole amino alcohols. *J. Med. Chem.* **1968**, *11*, 270-273.
- [26] Yalcin, I.; Oren, I.; Sener, E.; Akin, A.; Ucarturk, N. The synthesis and the structure-activity relationships of some substituted benzoxazoles, oxazolo(4,5-b)pyridines, benzothiazoles and benzimidazoles as antimicrobial agents. *Eur. J. Med. Chem.* **1992**, *27*, 401-406.
- [27] Gurupadayya, B. M.; Gopal, M.; Padmashali, B.; Vaidya, V. P. Synthesis and biological activity of novel pyrimidino imidazolines. *Int. J. Heterocyc. Chem.* **2005**, *15*, 169-172.
- [28] Latrofa, A.; Franco, M.; Lopodota, A.; Rosato, A.; Carone, D.; Vitali, C. Structural modifications and antimicrobial activity of *N*-cycloalkenyl-2-acylalkylidene-2,3-dihydro-1,3-benzothiazoles. *I L Farmaco* **2005**, *60*, 291-297.
- [29] Beneteau, V.; Besson, T.; Guillard, J.; Leonce, S.; Pfeiffer, B. Synthesis and *in vitro* antitumour evaluation of benzothiazole-2-carbonitrile derivatives. *Eur. J. Med. Chem.* **1999**, *34*, 1053-1060.
- [30] Wells, G.; Bradshaw, T. D.; Diana, P. Antitumour benzothiazoles. Part 10: The synthesis and antitumour activity of benzothiazole substituted quinol derivatives. *Bioorg. Med. Chem. Lett.* **2000**, *10*, 513-515.
- [31] Shi, D.; Bradshaw, T. D.; Chua, M.; Westwell, A. D. Stevens, M. F. G. Antitumour benzothiazoles. part 15: the synthesis and physico-chemical properties of 2-(4-aminophenyl)benzothiazole sulfamate salt derivatives. *Bioorg. Med. Chem.* **2001**, *11*, 1093-1095.
- [32] Hutchinson, I.; Bradshaw, T. D.; Matthews, C. S.; Stevens, M. F. G.; Westwell, A. D. Antitumour benzothiazoles. Part 20: 3'-cyano and 3'-alkynyl-Substituted 2-(4'-aminophenyl)benzothiazoles as new potent and selective analogues. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 471-474.
- [33] Jayachandran, E.; Bhatia, K.; Naragud, L. V. G.; Roy, A. Anthelmintic activity of 2-[3-amino, 5-S-methyl 4-carboxamido pyrazol-1-YL] 6-fluoro, 7-substituted (1,3) benzothiazoles on *Perituma* posthuman. *Ind. Drugs.* **2003**, *40*, 408-411.
- [34] Siddiqui, N.; Alam, M.; Siddiqui, A. A. Synthesis and analgesic activity of some 2-[(4-(alkyl thioureido)phenyl)sulphonamido]-6-substituted benzothiazoles. *Asian J. Chem.* **2004**, *16*, 1005-1008.
- [35] Lee, C. L.; Lam, Y.; Lee, S. Solid-phase combinatorial synthesis of benzothiazole and 2,3-dihydro-[1,5]-benzothiazepine derivatives. *Tetrahedron Lett.* **2001**, *42*, 109-11.

- [36] Ruiz, F.; Hazemann, I.; Mitschler, A.; Joachimiak, A.; Schneider, T.; Karplud, M.; Podjarnya A. The crystallographic structure of the aldose reductase±IDD552 complex shows direct proton donation from tyrosine 48. *Acta Crystallogr. Sect. D.* **2004**, *D60*, 1347-1354.
- [37] Nagoshi, N.; Nakashima, H.; Fehlings, M. G. Riluzole as a neuroprotective drug for spinal cord injury: from bench to bedside. *Molecules* **2015**, *20*, 7775-7789.
- [38] Rahman, M. M.; Tikhomirova, A.; Modak, J. K.; Hutton, M. L.; Supuran, C. T.; Roujeinikova, A. Antibacterial activity of ethoxzalamide against helicobacter pylori strains SS1 and 26695. *Gut Pathogen.* **2020**, *12*:20.
- [39] Sato, S.; Kajiura, T.; Noguchi, M.; Takehana, K.; Kobayashi, T.; Tsuji, T. J. AJI9561, a new cytotoxic benzoxazole derivative produced by streptomyces sp. *Antibiot.* **2001**, *54*, 102.
- [40] Kosmowska, B.; Wardas, J.; Głowacka, U.; Ananthan, S.; Ossowska, K. Pramipexole at a low dose induces beneficial effect in the harmaline-induced model of essential tremor in rats. *CNS Neurosci.Therapeut.* **2016**, *22*, 53-62.
- [41] Costanzo, M. J.; Yabut, S. C.; Almond, H. R. Jr.; Andrade-Gordon, P.; Corcoran, T. W.; Garavilla, L. de.; Kauffman, J. A.; Abraham, W. M.; Recacha, R.; Chattopadhyay, D.; Maryanoff, B. E. Potent, small-molecule inhibitors of human mast cell tryptase. antiasthmatic action of a dipeptide-based transition-state analogue containing a benzothiazole ketone. *J. Med. Chem.* **2003**, *46*, 3865-3876.
- [42] Shi, D.-F.; Bradshaw, T. D.; Wrigley, S.; McCall, C. J.; Lelieveld, P.; Fichtner, I.; Stevens, M. F. G. Antitumor benzothiazoles, synthesis of 2-(4-Aminophenyl)benzothiazoles and evaluation of their activities against breast cancer cell lines *in vitro* and *in vivo*. *J. Med. Chem.* **1996**, *39*, 3375–3384.
- [43] Beebe, X.; Wodka, D.; Sowin, T. J. Solid-phase synthesis of benzoxazoles from 3-nitrotyrosine. *J. Comb. Chem.* **2001**, *3*, 360–366.
- [44] Hari, A.; Karan, C.; Rodrigues, W. C.; Miller, B. L. Extending the scope of chromium–manganese redox-coupled reactions: a one-pot synthesis of benzoxazoles. *J. Org. Chem.* **2001**, *66*, 991–996.
- [45] Pottorf, R. S.; Chadha, N. K.; Katkevics, M.; Ozola, V.; Suna, E.; Ghane, H.; Regberg, T.; Player, M. R. Parallel synthesis of benzoxazoles via microwave-assisted dielectric heating. *Tetrahedron Lett.* **2003**, *44*, 175–178.
- [46] Chen, F.; Shen, C.; Yang, D. A simple protocol for the synthesis of 2-arylbenzoxazoles by oxidation with o-iodoxybenzoic acid (IBX) and its application in the synthesis of arylbenzoxazole-containing amino acids. *Tetrahedron Lett.* **2011**, *52*, 2128–2131.
- [47] Xu, W.; Zeng, M.-T.; Liu, S. S.; Li, Y. S.; Dong Z. B. Copper catalyzed synthesis of benzoxazoles and bezothiazoles via via tandem manner. *Tetrahedron Lett.* **2017**, *58*, 4289–4292.
- [48] Chua, M.-S.; Shi, D.-F.; Wrigley, S.; Bradshaw, T. D.; Hutchinson, I.; Shaw, P. N.; Barrett, D. A.; Stanley, L. A.; Stevens, M. F. G. Antitumor benzothiazoles. 7. synthesis of 2-(4-acylaminophenyl)benzothiazoles and investigations into the role of acetylation in the antitumor activities of the parent amines. *J. Med. Chem.* **1999**, *42*, 381–392.
- [49] Kashiya E.; Hutchinson I.; Chua M.S.; Stinson S. F.; Phillips L. R.; Kaur G.; Sausville E. A.; Bradshaw T. D.; Westwell A. D.; Stevens M. F. G. Antitumor benzothiazoles. 8. synthesis, metabolic formation and biological properties of the C- and N-oxidation products of antitumor 2-(4-aminophenyl)benzothiazoles. *J. Med. Chem.* **1999**, *42*, 4172–4184.
- [50] Hutchinson I.; Chua M.-S.; Browne H. L.; Trapani V.; Bradshaw T. D.; Westwell A. D.; Stevens M. F. G. Antitumor benzothiazoles. 14.¹ synthesis and in vitro biological properties of fluorinated 2-(4-aminophenyl)benzothiazoles. *J. Med. Chem.* **2001**, *44*, 1446–1455.
- [51] Leng W.; Zhou Y.; Xu Q.; Liu J. Synthesis and characterization of nonlinear optical side-chain polyimides containing the benzothiazole chromophores. *Macromolecules* **2001**, *34*, 4774–4779.
- [52] Hao A. N.; Long H. N.; Quynh N. B.; Hai T. N.; Khang Q. N.; Phuong H. T. Straightforward synthesis of benzoxazoles and benzothiazoles via photocatalytic radical cyclization of 2-substituted anilines with aldehydes. *Catalysis Comm.* **2020**, *145*, 106120.
- [53] Hutchinson I.; Jennings S. A.; Vishnuvajjala B. R.; Westwell A. D.; Stevens M. F. G. Antitumor benzothiazoles. 16. synthesis and pharmaceutical properties of antitumor 2-(4-aminophenyl)benzothiazole amino acid prodrugs. *J. Med. Chem.* **2002**, *45*, 744-747.
- [54] Blacker A. J.; Farah M. M.; Hall M. I.; Marsden S. P.; Saidi O.; Williams J. M. Synthesis of benzazoles by hydrogen-transfer catalysis. *Org. Lett.* **2009**, *11*, 2039-2042.
- [55] Shi D. F.; Bradshaw T. D.; Wrigley S.; McCall C. J.; Lelieveld P.; Fichtner I.; Stevens. M. F. Synthesis of 2-(4-aminophenyl)benzothiazoles and evaluation of their activities against breast cancer cell lines *in Vitro* and *in Vivo*. *J. Med. Chem.* **1996**, *39*, 3375–3384.
- [56] Tandon V. K.; Kumar M. BF₃·Et₂O promoted one-pot expeditious and convenient synthesis of 2-substituted benzimidazoles and 3,1,5-benzoxadiazepines. *Tetrahedron Lett.* **2004**, *45*, 4185–4187.
- [57] Charton J.; Girault-Mizzi S.; Sergheraert C. Conversion of sterically hindered diacylated 1,2-phenylenediamines into 2-substituted benzimidazoles. *Chem. Pharm. Bull.* **2005**, *53*, 492–497.

Synthesis of benzoxazoles and benzothiazoles

- [58] Seijas J. A.; Vazquez T.; Pilar M.; Carballido R.; Raquel M.; Crecente C. J.; Romar-Lopez L. Lawesson's Reagent and microwaves: a new efficient access to benzoxazoles and benzothiazoles from carboxylic acids under solvent-free conditions. *Syn. Lett.* **2007**, 313–317.
- [59] Maradolla M. B.; Allam S. K.; Mandha A.; Chandramouli G. V. P. One pot synthesis of benzoxazoles, benzthiazoles and benzimidazoles from carboxylic acids using ionic liquid. *Arkivoc* **2008**, XV, 42–46.
- [60] Raghavendra G. M.; Ramesha A. B.; Revanna C. N.; Nandeesh K. N.; Mantelingu K.; Rangappa K. S. One-pot tandem approach for the synthesis of benzimidazoles and benzothiazoles from alcohols. *Tetrahedron Lett.* **2011**, 52, 5571–5574.
- [61] Laszlo C.; Werner F.; Bernhard K.; Ursula H.; Johannes F.; Ulrich J. New kilogram-synthesis of the anti-alzheimer drug (–)-galanthamine. *Tetrahedron Lett.* **1998**, 39, 2087–2088.
- [62] Grego G.; Luucia P.; Alessandra N.; Marco D. Biologically inspired one-pot access routes to 4-hydroxybenzothiazole amino acids, red hair-specific markers of UV susceptibility and skin cancer risk. *Tetrahedron Lett.* **2009**, 50, 3095–3097.
- [63] Xiao Z. Y.; Ji D. G. Effects of available nitrogen on the uptake and assimilation of ferrocyanide and ferricyanide complexes in weeping willows. *J. Hazard. Mater.* **2008**, 156, 300–307.
- [64] Gaffar M. A.; A-Abu-El dl. Indirect band gap and optical parameters of pure and doped potassium ferrocyanide single crystals. *Physica B.* **2000**, 292, 221–232.
- [65] Ghotas E.; Batey R. A. Parallel synthesis of a library of benzoxazoles and benzothiazoles using ligand-accelerated copper-catalyzed cyclizations of ortho-halobenzanilides. *J. Org. Chem.* **2006**, 71, 1802–1808.
- [66] Naeimi H.; Rahmatinejad S. Microwave-assisted synthesis of 2-substituted benzoxazoles in the presence of potassium cyanide as catalyst. *Synth. React. Inorg. Met. Org. Nano Metal Chem.* **2016**, 46, 471–476.
- [67] Hu Wei-Ye.; Wang Pei-P.; Zhang Song-Lin An iron and copper system catalyzed C–H arylation of azoles with arylboronic acids. *Synthesis* **2015**, 47, 42–48.
- [68] Karimian A.; Kakhki R. M.; Beidokhti H. K. Magnetic Co-doped NiFe₂O₄ Nanocomposite: A heterogeneous and recyclable catalyst for the one-pot synthesis of benzimidazoles, benzoxazoles and benzothiazoles under solvent-free conditions. *J. Chin. Chem. Soc.* **2017**, 64, 11.
- [69] Praveen C.; Hemanth Kumar K.; Muralidharan D.; Perumal P.T. Oxidative cyclization of thiophenolic and phenolic Schiff's bases promoted by PCC: a new oxidant for 2-substituted benzothiazoles and benzoxazoles, *Tetrahedron* **2008**, 64, 2369–2374.
- [70] Merroun Y.; Chehab S.; Ghailane T.; Akhazzane M.; Souizi A.; Ghailane R. Preparation of tin-modified mono-ammonium phosphate fertilizer and its application as heterogeneous catalyst in the benzimidazoles and benzothiazoles synthesis. *React. Kinet. Mechanism. Catal.* **2019**, 126, 249–264.
- [71] Seth, K.; Nautiyal, M.; Purohit, P.; Parikh, N.; Chakraborti, A. K. Palladium Catalyzed Csp²-H Activation for Direct Aryl Hydroxylation: Unprecedented Role of 1,4-Dioxane as Source of Hydroxyl Radical. *Chemical Communication* **2014**, 1, 51.
- [72] Yang, X.-L.; Xu, C.-M.; Lin, S.-M.; Chen, J.-X.; Ding, J.-C.; Wu, H.-Y.; Su, W.-K. Eco-friendly synthesis of 2-substituted benzothiazoles catalyzed by cetyltrimethyl ammonium bromide (CTAB) in water. *J. Braz. Chem. Soc.* **2010**, 21, 37–42.
- [73] Li, Z.; Zhan, P.; Naesens, L.; Vanderlinden, E.; Liu, A.; Du, G.; Clercq, E. D.; Liu, X. Synthesis and preliminary biological evaluation of 5-substituted-2-(4-substituted phenyl)-1,3-benzoxazoles as a novel class of influenza virus a inhibitors. *Chem. Biol. Drug. Des.* **2012**, 79, 1018–1024.