SHORT REPORT



Rec. Nat. Prod. 17:1 (2023) 174-178

records of natural products

Three New Chromone Derivatives from the Deep-Sea-Derived

Fungus Penicillium thomii

Fan Yang ⁽ⁱ⁾, Yu Liu ⁽ⁱ⁾, Xiaoqian Zhang ⁽ⁱ⁾, Wan Liu ⁽ⁱ⁾,

Ying Qiao ^(D) ², Wei Xu ^(D) ^{2, 3, *}, Qin Li ^(D) ¹ and Zhongbin Cheng ^(D) ^{1,*}

¹School of Pharmacy, Henan University, Kaifeng 475004, People's Republic of China ²Key Laboratory of Tropical Marine Ecosystem and Bioresource, Fourth Institute of Oceanography, Ministry of Natural Resources, Beihai 536000, People's Republic of China ³Key Laboratory of Marine Biogenetic Resources, Third Institute of Oceanography, Ministry of Natural Resources, Xiamen 361005, People's Republic of China

(Received April 22, 2022; Revised May 11, 2022; Accepted May 15, 2022)

Abstract: Chemical investigation of the EtOAc extract of a deep-sea-derived fungus *Penicillium thomii Maire* YPGA3 cultured on rice solid medium led to the isolation of three new chromone analogs, named penithochromones U–W (1–3), along with six known compounds (4–9). The structures were determined by extensive analyses of spectroscopic data (NMR and HRESIMS data). Compounds 1–3 are 5,7-dioxygenated chromone analogs bearing an aliphatic acid side chain. All isolated compounds were inactive toward the α -glucosidase at the concentration of 200 μ M.

Keywords: *Penicillium thomii*; deep-sea fungus; chromone derivative; penithochromone. © 2022 ACG Publications. All rights reserved.

1. Plant Source

The fungal strain YPGA3 was isolated from the deep-sea sediments that collected in the Yap Trench (West Pacific Ocean) at a depth of -4500 m. The strain was identified as *Penicillium thomii* based on microscopic examination and by internal transcribed spacer (ITS) sequencing. The ITS sequence has been deposited in GenBank (http://www.ncbi.nlm.nih.gov) with the accession number MG835903. The strain YPGA3 (MCCC 3A01052) was deposited at the Marine Culture Collection of China

2. Previous Studies

In the past ten years, marine-derived fungi have gained increasing attention from natural product chemists, lots of small molecules bearing diverse structures and significant pharmacological effects have been reported. The strains belonging to the genera *Penicillium* from marine environment are continually studied and proved to be productive, leading to the discovery of DMOA-based

The article was published by ACG Publications

http://www.acgpubs.org/journal/records-of-natural-products Month-Month 2022 EISSN:1307-6167

DOI: http://doi.org/10.25135/rnp.330.2204.2433

^{*} Corresponding author: E- Mail: <u>xuwei@tio.org.cn</u> (W. Xu). czb360@126.com (Z.B. Cheng).

Yang et al., Rec. Nat. Prod. (2023) 17:1 174-178

meroterpenoids with unprecedented skeletons [1, 2], chlorinated diphenyl ethers possessing significant inhibitory effects against α -glucosidase [3], *p*-terphenyls with promising anti-HSV activity [4], antimicrobial alkaloids [5, 6], terpenoids [7, 8], and steroids [9, 10]. In our continuing research for new or bioactive metabolites from marine-derived fungi [7, 11, 12], three new chromone analogs (1–3) and six known compounds (4–9) were identified from a deep-sea derived strain *P. thomii* (Figure 1). Herein, the isolation and structural elucidation of the metabolites were described.

3. Present Study

The fermentation was conducted in 30 Fernbach flasks (500 mL), each containing 70 g of rice. Artificial seawater (90 mL) was added to each flask, and the contents were soaked for 3 h before autoclaving at 15 psi for 30 min. After cooling to room temperature, each flask was inoculated with 3.0 mL of the spore inoculum and incubated at room temperature for 30 days. The fermented material was extracted successively with EtOAc (4000 mL) for three times. After evaporation under vacuum, the EtOAc extract (3.1 g) was subjected to a middle chromatogram isolated gel (MCI) with MeOH/H₂O (10:90 \rightarrow 100:0) as eluent to obtain 10 fractions (Fr. 1 to Fr. 10). Fr. 3 (280 mg) was chromatographed over RP-18 silica gel eluted with MeOH/H₂O (30:70 \rightarrow 90:10) to afford 14 subfractions (Fr. 3a–Fr. 3n). Fr. 3b (32 mg) was separated by HPLC (MeCN/H₂O = 34:66, 3 mL/min) to yield 1 (4.6 mg, $t_R = 9.5$ min), 2 (11.3 mg, $t_R = 11.2$ min) and 3 (1.9 mg, $t_R = 13.4$ min). Fr. 3c (23) mg) was purified by HPLC (MeCN/H₂O = 44:56, 3 mL/min) to give 6 (5.3 mg, t_R = 32.2 min). Fr. 3g (27 mg) was subjected to HPLC (MeCN/H₂O = 51:49, 3 mL/min) to afford 5 (6.7 mg, $t_R = 13.4$ min) and 7 (2.3 mg, $t_R = 9.8$ min). Fr. 7 (150 mg) was chromatographed over ODS silica gel CC eluted with MeOH/H₂O (30:70→100:0) to give seven subfractions (Fr. 7a–Fr. 7g). Fr. 7a (19 mg) was further purified by HPLC (MeCN/H₂O=15:85, 3 mL/min) to yield 4 (2.3 mg, $t_R = 31.1$ min) and 8 (3.2 mg, t_R = 27.3 min). Fr. 7b (52 mg) was purified by HPLC (MeCN/H₂O=20:80, 3 mL/min) to give 9 (19.6 mg, $t_{R} = 33.7$ min).

Penithochromone U (1): light yellow oil; UV (MeOH) λ_{max} 249 (3.93), 289 (3.64) nm; ¹H and ¹³C NMR data, see Table 1; Negative HRESIMS m/z 291.0863 [M – H] – (calcd for C₁₅H₁₅O₆⁻, 291.0874).

Penithochromone V (2): light yellow oil; UV (MeOH) λ_{max} 247 (3.92), 290 (3.65) nm; ¹H and ¹³C NMR data, see Table 1; Negative HRESIMS m/z 277.0704 [M – H] – (calcd for C₁₄H₁₃O₆⁻, 277.0718).

Penithochromone W (3): light yellow oil; UV (MeOH) λ_{max} 250 (4.08), 294 (3.67) nm; ¹H and ¹³C NMR data, see Table 1; Negative HRESIMS m/z 249.0392 [M – H] ⁻ (calcd for C₁₂H₉O₆⁻, 249.0405).



Figure 1. Structures of compounds 1-9 from P. thomii YPGA3

Three new chromone derivatives from Penicillium thomii

The molecular formula of compound 1 was defined as $C_{15}H_{16}O_6$, based on analysis of the HRESIMS (m/z 291.0863 [M – H]⁻ (calcd 291.0874)) and the ¹³C NMR data, requiring eight double bond equivalents. The ¹H NMR spectrum displayed the signals of two aromatic protons for a metacoupling system [$\delta_{\rm H}$ 6.34 (1H, d, J = 1.9 Hz), 6.35 (1H, d, J = 1.9 Hz)], an aromatic singlet ($\delta_{\rm H}$ 5.87, s), an aromatic methoxyl group [$\delta_{\rm H}$ 3.75 (3H, s)], as well as several aliphatic protons (Table 1). Analysis of the ¹³C NMR (Table 1) and HSQC spectra revealed a total of 15 carbon resonances, which were attributed to a benzene ring, a double bond, two carbonyl carbons (δ_c 175.7, 174.5), a methoxyl carbon ($\delta_{\rm C}$ 55.9), and four methylene carbons ($\delta_{\rm C}$ 33.4, 32.3, 25.6, 23.9). The above-mentioned functional groups accounted for seven double bond equivalents, the remaining one revealed an additional ring in the structure. The presence of meta-coupled aromatic protons ($\delta_{\rm H}$ 6.34 and 6.35) indicated a 1.2,3,5-tetra substituted benzene ring. The HMBC correlations (Figure 2) from H-3 ($\delta_{\rm H}$ 5.87) to C-2 ($\delta_{\rm C}$ 165.8), C-4 ($\delta_{\rm C}$ 175.7), C-4a ($\delta_{\rm C}$ 107.0), H-6 ($\delta_{\rm H}$ 6.34) to C-4a, and H-8 ($\delta_{\rm H}$ 6.35) to 4a, 8a (δ_c 159.5), in combination with the additional ring enabled the construction of a 5,7dioxygenated chromone nucleus. The methoxy group was located at C-5 by the HMBC correlation from the methoxyl protons at $\delta_{\rm H}$ 3.75 to the oxygenated aromatic carbon C-5 ($\delta_{\rm C}$ 160.7). Analysis of the COSY spectrum for 1 revealed the presence of a spin system including four methylenes (CH₂-9/ CH₂-10/ CH₂-11/ CH₂-12) (Figure 2). The HMBC correlations from the protons H₂-11 ($\delta_{\rm H}$ 1.53) and H₂-12 ($\delta_{\rm H}$ 2.24) to the carbonyl carbon C-13 ($\delta_{\rm C}$ 174.5) allowed a pentanoic acid side chain to be defined, which was anchored to the chromone nucleus at C-2 by HMBC correlations of H₂-9 to C-2 $(\delta_{\rm C} 165.8)$ and C-3 ($\delta_{\rm C} 110.4$). Thus, the structure of **1** was determined as 7-hydroxy-5-methoxy-2pentanoic acid-chromone and was named penithochromone U.

No	1 ^{<i>a</i>}		2 ^b		3 ^b	
	$\delta_{ m H}$	$\delta_{ m C}$	$\delta_{ m H}$	$\delta_{ m C}$	$\delta_{ m H}$	$\delta_{ m C}$
2		165.8		167.6		170.9
3	5.87, s	110.4	6.00, s	111.1	6.09, s	108.5
4		175.7		179.9		183.9
4a		107.0		108.1		105.2
5		160.7		162.5		163.3
6	6.34, d (1.9)	95.1	6.39, d (1.9)	96.2	6.19, d (2.1)	100.1
7		162.6		165.0		166.0
8	6.35, d (1.9)	96.4	6.39, d (1.9)	97.4	6.33, d (2.1)	94.9
8a		159.5		161.6		159.8
9	2.53, t (7.0)	32.3	2.90, t (7.1)	29.6	2.93, t (7.2)	30.3
10	1.61, m	25.6	2.75, t (7.1)	31.4	2.73, t (7.2)	31.6
11	1.53, m	23.9		174.0		175.4
12	2.24, t (7.2)	33.4				
13		174.5				
5-OCH ₃	3.75, s	55.9	3.86, s	56.4		
11-OCH ₃			3.69, s	52.4		

Table 1. ¹H (400 Hz) and ¹³C NMR (100 Hz) Data of 1-3 (δ in ppm)

^{*a*} in DMSO-*d*₆, ^{*b*} in methanol-*d*₄

The molecular formula of **2** was determined to be $C_{14}H_{14}O_6$ by the HRESIMS data. The NMR data showed similar structural features as those of **1**, presenting the signals for the same chromone moiety [$\delta_H 6.00$ (s), 6.39 (d, J = 1.9 Hz, 1H), 6.51 (d, J = 1.9 Hz, 1H); δ_C 96.2, 97.4, 108.1, 111.1, 161.6, 162.5, 165.0, 167.6, 179.9] as that of **1**. The structure of **2** was established by 2D NMR analyses (Figure 2). Analysis of the COSY spectrum for **2** revealed the connectivity of the two methylenes [$\delta_H 2.90$ (t, J = 7.1 Hz, 2H), 2.75 (t, J = 7.1 Hz, 2H)], both showed HMBC correlations to the ester carbonyl carbon C-11 (δ_C 174.0). The HMBC correlation from the remaining methoxyl protons (δ_H 3.69) to C-11 assigned a methyl propionate side chain, which was attached to the

chromone unit at C-2 by HMBC correlations of H₂-9 to C-2 (δ_{C} 167.6) and C-3 (δ_{C} 111.1). Thus, the structure of **2** was determined as 7-hydroxy-5-methoxy-2-methyl propionate-chromone and was named penithochromone V.



Figure 2. Key $^{1}\text{H}^{-1}\text{H}$ COSY (—) and HMBC (\rightarrow) correlations of 1–3

Compound **3** had a molecular formula of $C_{12}H_{10}O_6$, as determined by the HRESIMS along with ¹³C NMR data. The ¹H NMR and ¹³C NMR spectra of **3** were very similar to those of **2** with differences due to the absence of the two methoxy groups in **2**, indicating **3** was demethylated derivative of **2**. The structure of **3** was confirmed by the 2D NMR analyses (Figure 2). Compound **3** was established as 5,7-dihydroxy-2-propanoic acid-chromone and was named penithochromone W.

Additionally, the known compounds 4-9 were identified to be isoeugenin (4) [13], 6,8dihydroxy-3-methylisocoumarin (5) [14], methyl indole-3-acetate (6) [15], methyl indole-3-lactate (7) [16], crithminic acid (8) [17], and methyl (4-hydroxyphenyl)acetate (9) [18] by comparing their NMR data with those in the literature.

The isolated compounds were screened for their inhibitory effect toward the α -glucosidase at the initial concentration of 200 μ M, while all compounds were inactive with inhibition rate less than 30%.

Supporting Information

Supporting Information accompanies this paper on <u>http://www.acgpubs.org/journal/records-of-natural-products</u>

Acknowledgments

This work was supported by the Fund of the Key Laboratory of Tropical Marine Ecosystem and Bioresource MNR (2021QN04) and the National Natural Science Foundation of China (81903536)

ORCID 💷

Fan Yang: 0000-0001-6608-0371 Yu Liu: 0000-0001-9687-3153 Xiaoqian Zhang: 0000-0001-6767-3275 Wan Liu: 0000-0001-5612-9434 Ying Qiao: 0000-0002-4356-0927 Wei Xu: 0000-0002-3265-7475 Qin Li: 0000-0001-8295-6230 Zhongbin Cheng: 0000-0003-0942-6422

References

[1] X. Cheng, X. Liang, Z.H. Zheng, X. X. Zhang, X. H. Lu, F. H. Yao and S. H. Qi (2020). Penicimeroterpenoids A–C, meroterpenoids with rearrangement skeletons from the marine-derived fungus *Penicillium* sp. SCSIO 41512, *Org. Lett.* **22**, 6330-6333.

- [2] J. Zhang, Y. Wu, B. Yuan, D. Liu, K. Zhu, J. Huang, P. Proksch and W. Lin (2019). DMOA-based meroterpenoids with diverse scaffolds from the sponge-associated fungus *Penicillium brasilianum*, *Tetrahedron* **75**, 2193-2205.
- [3] J. F. Wang, L. M. Zhou, S. T. Chen, B. Yang, S. R. Liao, F. D. Kong, X. P. Lin, F. Z. Wang, X. F. Zhou and Y. H. Liu (2018). New chlorinated diphenyl ethers and xanthones from a deep-sea-derived fungus *Penicillium chrysogenum* SCSIO 41001, *Fitoterapia* 125, 49-54.
- [4] W. Chen, J. Zhang, X. Qi, K. Zhao, X. Pang, X. Lin, S. Liao, B. Yang, X. Zhou, S. Liu, J. Wang, X. Yao and Y. Liu (2021). *p*-Terphenyls as anti-HSV-1/2 agents from a deep-sea-derived *Penicillium* sp., *J. Nat. Prod.* 84, 2822-2831
- [5] Y. H. Li, X. M. Li, X. Li, S. Q. Yang, X. S. Shi, H. L. Li and B. G. Wang (2020). Antibacterial alkaloids and polyketide derivatives from the deep sea-derived fungus *Penicillium cyclopium* SD-413, *Mar. Drugs*, **18**, 553.
- [6] D. Tian, X. Gou, J. Jia, J. Wei, M. Zheng, W. Ding, H. Bi, B. Wu and J. Tang (2022). New diketopiperazine alkaloid and polyketides from marine-derived fungus *Penicillium* sp. TW58-16 with antibacterial activity against *Helicobacter pylori*, *Fitoterapia* 156, 105095.
- [7] Y. Li, W. Liu, S. Han, J. Zhang, W. Xu, Q. Li and Z. Cheng (2020). Penitholabene, a rare 19-nor labdane-type diterpenoid from the deep-sea-derived fungus *Penicillium thomii* YPGA3 *Fitoterapia* 146, 104691.
- [8] F. Li, W. Sun, S. Zhang, W. Gao, S. Lin, B. Yang, C. Chai, H. Li, J. Wang, Z. Hu and Y. Zhang (2020). New cyclopiane diterpenes with anti-inflammatory activity from the sea sediment-derived fungus *Penicillium* sp. TJ403-2, *Chin. Chem. Lett.* **31**, 197-201.
- [9] T. P. T. Hoang, C. Roullier, G. Genta-Jouve, M. C. Boumard, T. Robiou du Pont, H. Nazih, Y. F. Pouchus and O. Grovel (2019). C25 steroids from the marine mussel-derived fungus *Penicillium ubiquetum* MMS330, *Phytochem. Lett.* 34, 18-24.
- [10] S. S. Gao, X. M. Li, C. S. Li, P. Proksch and B. G. Wang (2011). Penicisteroids A and B, antifungal and cytotoxic polyoxygenated steroids from the marine alga-derived endophytic fungus *Penicillium chrysogenum* QEN-24S, *Bioorg. Med. Chem. Lett.* 21, 2894-2897.
- [11] Z. Cheng, Y. Li, W. Xu, W. Liu, L. Liu, D. Zhu, Y. Kang, Z. Luo and Q. Li (2019). Three new cyclopiane-type diterpenes from a deep-sea derived fungus *Penicillium* sp. YPGA11 and their effects against human esophageal carcinoma cells, *Bioorg. Chem.* **91**, 103129.
- [12] Q. Li, W. Xu, R. Fan, J. Zhang, Y. Li, X. Wang, S. Han, W. Liu, M. Pan and Z. Cheng (2020). Penithoketone and penithochromones A–L, polyketides from the deep-sea-derived fungus *Penicillium thomii* YPGA3, *J. Nat. Prod.* 83, 2679-2685.
- [13] H. J. An, A. Nugroho, B. M. Song and H. J. Park (2015). Isoeugenin, a novel nitric oxide synthase inhibitor isolated from the rhizomes of *Imperata cylindrica*, *Molecules* **20**, 21336-21345.
- [14] J. K. Kendall, T. H. Fisher, H. P. Schultz and T. P. Schultz (1989). An improved synthesis of 6,8dimethoxy-3-methylisocoumarin, a fungal metabolite precursor, *J. Org. Chem.* **54**, 4218-4220.
- [15] Y. Liu, J. H. Jung and S. Zhang (2006). Indole alkaloids from a sponge Sarcotragus species, Biochem. Syst. Ecol. 34, 453-456.
- [16] R. Uchida, K. Shiomi, T. Sunazuka, J. Inokoshi, A. Nishizawa, T. Hirose, H. Tanaka, Y. Iwai and S. Omura (1996). Kurasoins A and B, new protein farnesyltransferase inhibitors produced by *Paecilomyces* sp. FO-3684. II. Structure elucidation and total synthesis, *J. Antibiot.* 49, 886-889.
- [17] A. K. C. Schmidt and C. B. W. Stark (2011). TPAP-catalyzed direct oxidation of primary alcohols to carboxylic acids through stabilized aldehyde hydrates, *Org. Lett.* **13**, 4164-4167.
- [18] D. S. Bose and A. V. Narsaiah (2005). An efficient asymmetric synthesis of (S)-atenolol using hydrolytic kinetic resolution, *Bioorg. Med. Chem.* **13**, 627-630.

