

# Green synthesis of 3,4-disubstituted isoxazol-5(4*H*)-one using Gluconic acid aqueous solution as an efficient recyclable medium

Ganesh V. Shitre <sup>1</sup>, Anisha R. Patel <sup>2</sup> and Ramesh S. Ghogare <sup>2\*</sup>

<sup>1</sup>Department of Chemistry, Vaishnavi Mahavidyalaya, Wadwani, Dist-Beed- 431144, India

<sup>2</sup>Department of Chemistry, B. N. N. College Bhiwandi, Dist-Thane-421305, India

(Received April 11, 2023; Revised May 24, 2023; Accepted May 29, 2023)

**Abstract:** Green synthesis of 3,4-disubstituted isoxazol-5(4*H*)-ones has been achieved via Knoevenagel condensation using three-component coupling reaction of aromatic aldehyde, ethyl acetoacetate and hydroxylamine hydrochloride in Gluconic acid aqueous solution (50 wt % GAAS). In this methodology, Gluconic acid aqueous solution used as reaction medium as well as catalyst and it can be recycled and reused several times without loss of its a significant efficacy. All compounds have been confirmed by <sup>1</sup>H NMR, <sup>13</sup>C NMR, IR spectroscopy and mass spectrometry.

**Keywords:** Green synthesis; gluconic acid aqueous solution; bio-based green solvent; Knoevenagel condensation; 3,4-disubstituted isoxazol-5(4*H*)-ones. © 2023 ACG Publications. All rights reserved.

## 1. Introduction

Nowdays, environmental benign green synthesis is important perspective in chemical and pharmaceutical productions, because uses of conventional organic solvents as well as reagents are harmful and toxic to environments<sup>1-2</sup>. In addition, some other organic solvents are also responsible to depletion of ozone layer. Therefore, use of secure and less toxic chemicals as the solvent is one of the most important criterions among the green chemistry principles. The purpose of using “green reaction medium” is to reduce or eliminate the toxic and harmful effects of chemicals on the environment. Hence, emerging interest in the development of novel methodologies in organic synthesis, researchers have need to find out an alternative non-polluting as well as bio-degradable green reaction medium<sup>3-10</sup> for organic transformations. Recently, Gluconic acid aqueous solution (50 wt % GAAS) is used as bio-based green solvent for the organic syntheses<sup>11-16</sup>. This solvent shows various characteristics like non-volatility, bio-degradability, recyclability, eco-friendly and cost-effective organic acid solvent. Present work has performed using Gluconic acid aqueous solution as reaction medium, which has been recycled and reused several times.

Isoxazole and its derivatives are significant class of heterocyclic compounds containing a potent pharmacophore and these pharmacophore plays important role in medicinal chemistry for building blocks of drugs<sup>17-19</sup>. Several substituted isoxazoles have been shown a broad range of biological activities such as anti-inflammatory<sup>20</sup>, immunosuppressive<sup>21</sup>, β-Adrenergic receptor antagonists<sup>22</sup>, Androgen antagonists<sup>23</sup>, anti-viral<sup>24</sup>, anti-HIV<sup>25-26</sup>, antiprotozoal<sup>27</sup>, HDAC inhibitors<sup>28</sup>, anti-tubercular<sup>29</sup>,

\* Corresponding author: E-mail: [rsghogare05@gmail.com](mailto:rsghogare05@gmail.com)

## Green synthesis of 3,4-disubstituted isoxazol-5(4*H*)-one

anti-bacterial<sup>30</sup>, anti-fungal<sup>31</sup>, anticancer<sup>32-33</sup> and antioxidant activity<sup>34</sup>. Some of isoxazoles are also used as fungicidal in agricultural field<sup>35</sup>. In addition to these, many 3,4-disubstituted isoxazol-5(4*H*)-ones are used in development of optical storage devices, optical storage and nonlinear optical research<sup>36-37</sup>, filter dyes in photographic films<sup>38</sup> and light conversion in molecular devices<sup>39</sup>.

In view of the emerging importance of 3,4-disubstituted isoxazol-5(4*H*)-ones, various modified methods were developed for their synthesis using several reagents, catalysts and strategies. Amongst them, some methods were carried out using different catalysts like acidic catalyst<sup>40-50</sup>, basic catalyst<sup>51-68</sup>, nano materials<sup>69-75</sup> and Synthetic enzyme<sup>76</sup>. Additionally, various bio-based solvents<sup>77-80</sup>, deep eutectic solvents<sup>81-82</sup>, ion exchange resins<sup>83-84</sup>, ionic liquids<sup>85-86</sup> as well as metal and its complexes<sup>87-89</sup> were also used. In similar manner, some of the newest techniques such as microwave irradiation<sup>90-92</sup> ultrasound irradiation<sup>93-94</sup> and visible light<sup>95</sup> have been reported. However, several of these reported methods suffer from one or two drawbacks such as harsh reaction conditions, expensive catalysts, toxic solvents, prolonged reaction time, poor yields and tedious workup process. Thus, the need for the development of an alternate route to synthesis of 3,4-disubstituted isoxazol-5(4*H*)-ones is in high demand.

During our recent studies directed towards the development of novel methodology using alternative procedures<sup>96-98</sup> and by considering the significance of 3,4-disubstituted isoxazol-5(4*H*)-ones, herein we wish to report, a green approach for the catalyst-free synthesis of 3,4-disubstituted isoxazol-5(4*H*)-one using Gluconic acid aqueous solution as an efficient recyclable medium.

## 2. Experimental

### 2.1. General Methods

Solvents and reagents were purchased from commercially and used without purification. Melting points were recorded on the Buchi R-535 apparatus and are uncorrected. <sup>1</sup>H NMR spectra were recorded on Gemini-300 spectrometer in CDCl<sub>3</sub> using TMS as an internal standard. IR spectra were recorded on a Bruker FT-IR spectrophotometer using neat or KBr disk and mass spectra were recorded on a Finnigan MAT 1020 mass spectrometer with operating at 70 eV.

### 2.2. General Procedure

A mixture of benzaldehyde (1 mmol), ethyl acetoacetate (1 mmol) and hydroxylamine hydrochloride (1 mmol) were stirred in gluconic acid aqueous solution (GAAS) (5 mL) at 70 °C. The progress of the reaction was monitored by thin layer chromatography (TLC) and the reaction was completed within 45 minutes. Then, the reaction mixture was cooled and extracted with ethyl acetate. The extracted ethyl acetate was concentrated under reduced pressure to obtained crude solid product, which was purified by crystallization from ethanol. All the pure products were confirmed by comparing their physical and spectral data.

### 2.3. Spectral Data for Compounds

(*Z*)-4-benzylidene-3-methylisoxazol-5(4*H*)-one (**4a**): Pale yellow solid (90 %). m.p. 143-145°C. IR (KBr):  $\nu_{\max}$  = 3225, 2370, 1740, 1635, 1215, cm<sup>-1</sup>; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta$  7.92 (d, *J* = 8.8 Hz, 2H), 7.49-7.61 (m, 3H), 7.36 (s, 1H), 2.40 (s, 3H) ppm; <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  175.7, 164.9, 152.1, 133.1, 127.8, 127.5, 127.4, 126.0, 17.4 ppm; ESIMS: *m/z* 188 [M+1]<sup>+</sup>.

(*Z*)-3-methyl-4-(4-methylbenzylidene)isoxazol-5(4*H*)-one (**4b**): Pale yellow solid (90%). m.p. 128-130°C. IR (KBr):  $\nu_{\max}$  = 3055, 2940, 2865, 1735, 1612, 1110 cm<sup>-1</sup>; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta$  7.92 (d, *J*=8.2 Hz, 2H); 7.48 (s, 1H); 7.36 (d, *J*=8.1Hz, 2H); 2.48 (s, 3H); 2.30 (s, 3H) ppm; <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>):  $\delta$  172.4, 163.7, 150.3, 136.0, 130.5, 127.9, 126.0, 120.3, 22.8, 17.8 ppm; ESIMS: *m/z* 201 [M]<sup>+</sup>.

(Z)-4-(4-hydroxybenzylidene)-3-methylisoxazol-5(4H)-one (**4c**): Yellow Solid (92%). m.p. 214-216 °C. IR (KBr):  $\nu_{\max}$  = 3460, 3080, 2940, 1745, 1615, 1220,  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$ : 10.70 (s, 1H), 8.42 (d,  $J$  = 8.8 Hz, 2H), 7.60 (d,  $J$  = 8.2 Hz, 2H), 7.38 (s, 1H), 3.00 (s, 3H) ppm;  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  174.7, 164.5, 157.2, 149.7, 130.8, 125.4, 123.4, 115.5, 17.4 ppm; ESIMS:  $m/z$  226  $[\text{M}+23]^+$ , 203  $[\text{M}]^+$ .

(Z)-4-(4-methoxybenzylidene)-3-methylisoxazol-5(4H)-one (**4d**): Yellow Solid (90%). m.p. 180-182 °C. IR (KBr)  $\nu_{\max}$  = 3080, 2980, 2815, 1736, 1605, 1270, 1025  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.10 (d,  $J$  = 8.8 Hz, 2H); 7.42 (s, 1H), 6.98 (d,  $J$  = 8.1 Hz, 2H), 3.85 (s, 3H), 2.40 (s, 3H) ppm;  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  174.4, 163.9, 160.6, 149.7, 130.8, 125.8, 114.1, 54.9, 17.3 ppm. ESIMS:  $m/z$  218  $[\text{M}+1]^+$ .

(Z)-4-(4-hydroxy-3-methoxybenzylidene)-3-methylisoxazol-5(4H)-one (**4e**): Pale yellow solid (92%). m.p. 214-216 °C. IR (KBr):  $\nu_{\max}$  = 3445, 3136, 2940, 1735, 1642, 1270  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.72 (s, 1H), 7.50 (s, 1H), 7.24-7.16 (m, 2H), 6.86 (d,  $J$  = 8.1 Hz, 1H), 3.42 (s, 3H), 2.31 (s, 3H) ppm;  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  174.7, 163.9, 151.2, 149.7, 148.3, 129.3, 125.4, 122.8, 116.5, 111.6, 56.0, 17.3 ppm. ESIMS:  $m/z$  251  $[\text{M}+18]^+$ .

(Z)-4-(3,4-dimethoxybenzylidene)-3-methylisoxazol-5(4H)-one (**4f**): Yellow solid (92%). m.p. 130-132 °C. IR (KBr):  $\nu_{\max}$  = 3092, 2935, 2830, 2355, 1730, 1565, 1266  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.50 (s, 1H), 7.320-7.28 (m, 2H), 6.90 (d,  $J$  = 8.1 Hz, 1H), 3.83 (s, 6H), 2.44 (s, 3H) ppm;  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  175.1, 167.6, 150.3, 149.4, 147.7, 127.9, 126.0, 122.6, 111.6, 111.1, 56.2, 17.4 ppm; ESIMS:  $m/z$  248  $[\text{M}+1]^+$ .

(Z)-4-(3,4,5-trimethoxybenzylidene)-3-methylisoxazol-5(4H)-one (**4g**): Yellow solid (94%). m.p. 138-140 °C. IR (KBr):  $\nu_{\max}$  = 3130, 2945, 1740, 1560, 1278, 1042,  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.40 (s, 1H), 7.08 (d,  $J$  = 8.8 Hz, 2H), 3.78 (s, 9H), 2.43 (s, 3H) ppm;  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  174.4, 164.5, 153.0, 150.3, 138.0, 129.7, 125.1, 103.7, 60.5, 56.3, 17.5 ppm; ESIMS:  $m/z$  278  $[\text{M}+1]^+$ , 277  $[\text{M}]^+$ .


(Z)-4-(4-(dimethylamino)benzylidene)-3-methylisoxazol-5(4H)-one (**4h**): pale red solid (90%). m.p. 212-214 °C. IR (KBr):  $\nu_{\max}$  = 1730, 1625, 1552, 1505, 1436, 1312, 1245, 1120  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.42 (d,  $J$  = 8.5 Hz, 2H), 7.34 (s, 1H), 6.71 (d,  $J$  = 9.2 Hz, 2H), 3.10 (s, 6H), 2.26 (s, 3H) ppm;  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  172.2, 162.4, 153.5, 149.3, 138.1, 122.6, 120.2, 111.6, 41.7, 13.0 ppm; ESIMS:  $m/z$  248  $[\text{M}+18]^+$ , 230  $[\text{M}]^+$ .

(Z)-3-methyl-4-(thiophen-2-ylmethylene)isoxazol-5(4H)-one (**4i**): Pale yellow solid (88%). m.p. 144-146 °C. IR (KBr):  $\nu_{\max}$  = 2930, 1736, 1695, 1510, 1330, 1145  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.15 (d,  $J$  = 3.6 Hz, 1H), 7.92 (d,  $J$  = 4.8 Hz, 1H), 7.60 (s, 1H), 7.35 (t,  $J$  = 4.8 Hz, 1H), 2.26 (s, 3H);  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  168.8, 160.6, 141.5, 140.7, 140.0, 136.8, 128.5, 114.9, 12.9 ppm; ESIMS:  $m/z$  193  $[\text{M}]^+$ .

### 3. Results and Discussion

In continuation our research interest in the development of green synthetic methodologies, we decided to explore the use of Gluconic acid aqueous solution (50 wt% GAAS) as reaction medium for the synthesis of 3,4-disubstituted isoxazol-5(4H)-one.

In viewpoint of green chemistry, the model reaction was carried out between benzaldehyde (**1**) (1.0 mmol), ethyl acetoacetate (**2**) (1.0 mmol) and hydroxylamine hydrochloride (**3**) (1.0 mmol) under various solvents conditions using different reaction temperature. The results are shown in Table 1. Initially, the reaction was carried out under solvent free condition at room temperature as well as higher temperature (100 °C) for prolonged reaction time (120 minutes) but formation of the desired product was not observed (Table 1, entry 1-2). After that, we chose water as a green solvent and reaction was carried out at different temperatures for 120 minutes. Amongst them, reflux condition progresses towards the desired product up to 30 % yield (Table 1, entry 3-4).

Green synthesis of 3,4-disubstituted isoxazol-5(4*H*)-one**Scheme 1.** Gluconic acid aqueous solution catalyzed synthesis of 3,4-disubstituted isoxazol-5(4*H*)-one

With these obtained results in hand, our next goal was to improve the product yield with reducing the reaction time. It has needed to use an appropriate catalyst which acts green catalyst as well as recyclable reaction medium and increases the product yield with reducing the reaction time. Here, we decided to explore the use of Gluconic acid aqueous solution (50 wt % GAAS) system as bio-based reaction medium and catalyst for this reaction. Because, Gluconic acid aqueous solution (GAAS) has hydrophobic nature with organic substrate which induce favorable aggregation of organic substrates in water to leads the fast collisions of the reactants and formation of desired product in short reaction times.

Initially, the reaction was performed at room temperature using various concentrations of GAAS system from 3 mL to 6 mL for 120 minutes (Table 1, entry 5-8). The yield of product gradually increases with increasing the volume of solvent till 5 mL, and further increasing the volume of reaction medium, no improvement occurred in terms of product yield. Thus, the observation shows that a 5 mL volume of GAAS system was enough for obtaining moderate yield of product at the room temperature. After these results, the effects of temperature were studied, by carrying out the model reaction at different temperatures from 40 to 80 °C. In this optimization of reaction, increasing the reaction temperature from 40 to 70 °C, the product yield increases with gradually decreasing reaction time (Table 1, entry 9-12). Further increasing the reaction temperature up to 80 °C, but yield of the product was not improved (Table 1, entry 13). The observation shows that reaction condition at 70 °C is sufficient for the completion of reaction within 45 minutes with excellent yield.























**Table 1.** Optimization of reaction solvent and GAAS (50 wt%) at different conditions

| Sr. No. | Solvent          | Quantity Solvent (mL) | Temperature (°C) | Time (min.) | Isolated Yields (%) |
|---------|------------------|-----------------------|------------------|-------------|---------------------|
| 1       | Solvent-free     | --                    | RT <sup>a</sup>  | 120         | NR <sup>b</sup>     |
| 2       | Solvent-free     | --                    | 100              | 120         | NR <sup>b</sup>     |
| 3       | H <sub>2</sub> O | 5                     | RT <sup>a</sup>  | 120         | Trace               |
| 4       | H <sub>2</sub> O | 5                     | Reflux           | 120         | 30                  |
| 5       | GAAS             | 3                     | RT <sup>a</sup>  | 120         | 40                  |
| 6       | GAAS             | 4                     | RT <sup>a</sup>  | 120         | 54                  |
| 7       | GAAS             | 5                     | RT <sup>a</sup>  | 120         | 60                  |
| 8       | GAAS             | 6                     | RT <sup>a</sup>  | 120         | 60                  |
| 9       | GAAS             | 5                     | 50               | 70          | 72                  |
| 10      | GAAS             | 5                     | 60               | 60          | 85                  |
| 11      | GAAS             | 5                     | 70               | 45          | 92                  |
| 12      | GAAS             | 5                     | 80               | 45          | 92                  |

RT<sup>a</sup> = Room Temperature, NR<sup>b</sup> = No Reaction (Product not formed).

Based on the optimized reaction conditions, various aromatic and heterocyclic aldehydes were reacted with ethyl acetoacetate and hydroxylamine hydrochloride for the synthesis of 3,4-disubstituted isoxazol-5(4*H*)-ones (Table 2, **4a-4k**) to demonstrate the scope of GAAS as catalyst as well as reaction medium. All results are summarized in table 2.

**Table 2.** Synthesis of the 3-Methyl-4-arylmethylene isoxazol-5(4*H*)-one

| Sr No    | Aldehyde  | Product   | Reaction Time (min.) | Isolated Yield (%) | M.P. °C [Lit. M.P.] <sup>Ref</sup> |
|----------|---|---|----------------------|--------------------|------------------------------------|
| <b>a</b> |    |    | 45                   | 92                 | 143-145 [142-144] <sup>[50]</sup>  |
| <b>b</b> |    |    | 45                   | 90                 | 128-130 [132-134] <sup>[50]</sup>  |
| <b>c</b> |    |    | 45                   | 92                 | 214-216 [210-212] <sup>[50]</sup>  |
| <b>d</b> |    |    | 45                   | 90                 | 180-182 [176-178] <sup>[50]</sup>  |
| <b>e</b> |    |    | 45                   | 92                 | 214-216 [210-212] <sup>[50]</sup>  |
| <b>f</b> |   |   | 45                   | 92                 | 130-132 [134-136] <sup>[50]</sup>  |
| <b>g</b> |  |  | 45                   | 94                 | 138-140 [134-136] <sup>[50]</sup>  |
| <b>h</b> |  |  | 45                   | 90                 | 212-214 [206-209] <sup>[50]</sup>  |
| <b>i</b> |  |  | 45                   | 90                 | 144-146 [146-147] <sup>[66]</sup>  |
| <b>j</b> |  |  | 120                  | 00                 | --                                 |
| <b>k</b> |  |  | 120                  | 00                 | --                                 |

NR\*= No Reaction

Various effects of substituents present on the aromatic aldehyde were examined by using different electron donating and withdrawing functional groups. The aromatic aldehyde having electron donating groups reacted in appropriate time to achieve desired products with excellent yields, while electron withdrawing groups failed to obtain the required product even increasing reaction time. In general, it observed that the nature of the functional group on aromatic aldehyde have a various effect on this reaction.

Green synthesis of 3,4-disubstituted isoxazol-5(4*H*)-one

For the investigation of reusability of the reaction medium, reaction was carried out using model reaction (Scheme 1). After the completion of the reaction, the reaction mixture was cooled and the formed product extracted with ethyl acetate from the GAAS phase.

**Table 3.** Recyclability study of the GAAS (50 wt %)

| Sr. No. | Cycle | Reaction Time (min.) | Isolated Yields (%) |
|---------|-------|----------------------|---------------------|
| 1       | 1     | 45                   | 92                  |
| 2       | 2     | 45                   | 90                  |
| 3       | 3     | 50                   | 85                  |
| 4       | 4     | 60                   | 82                  |

The immiscibility of Gluconic acid aqueous solution (GAAS) with ethyl acetate, it's easy to separate out and reused up to four times for further reactions without loss of its a significant efficacy (Table 3).

After that, GAAS solvent system has also compared with some previously reported green catalysts or reaction mediums used for synthesis of 3,4-disubstituted isoxazol-5(4*H*)-ones (Table 4). The results show that the present method has more advantages from the viewpoint of product yield and reaction time.

**Table 4.** Comparison for different green catalysts or reaction mediums used for synthesis of 3,4-disubstituted isoxazol-5(4*H*)-ones

| Sr. No. | Catalyst   | Solvent               | Temperature (°C) | Time (min) | Yield (%)    | Ref                 |
|---------|--|-----------------------|------------------|------------|--------------|---------------------|
| 1       | DL-Tartaric acid (10 mole %)                         | H <sub>2</sub> O      | RT               | 100        | 88           | 41                  |
| 2       | Citric acid (1 mmol)                                 | H <sub>2</sub> O      | RT               | 480        | 90           | 43                  |
| 3       | Pyruvic acid   | H <sub>2</sub> O      | 100              | 60         | 85-92        | 49                  |
| 4       | Succinic acid (15 mol %)                             | H <sub>2</sub> O      | RT               | 90         | 92           | 50                  |
| 5       | Lemon juice (1 mL)                                   | H <sub>2</sub> O:EtOH | 90               | 55-60      | 94           | 77                  |
| 6       | Starch solution (4 mL)                               | --                    | 90               | 50-60      | 80-86        | 78                  |
| 7       | Fruit juice (10 mL)                                  | H <sub>2</sub> O:EtOH | RT               | 240-420    | 90-95        | 80                  |
| 8       | <b>Gluconic acid aqueous solution (50 wt % GAAS)</b> | <b>--</b>             | <b>70</b>        | <b>45</b>  | <b>70-92</b> | <b>Present work</b> |

In general, all the reactions were very neat in terms of conversion as well as isolation of products. All the products were characterized by their spectroscopy analysis such as <sup>1</sup>H, <sup>13</sup>C NMR, IR spectroscopy and mass spectrometry.

#### 4. Conclusion

In summary, we have represented a simple, inexpensive and novel one pot three-component methodology for the green synthesis of 3,4-disubstituted isoxazol-5(4*H*)-ones derivatives using various aromatic aldehydes, ethyl acetoacetate and hydroxylamine hydrochloride in Gluconic acid aqueous solution. Gluconic acid aqueous solution is recycled and reused several times without loss of its effectiveness. The present method offers significant advantages such as economical, nontoxic, shorter reaction time, catalyst-free condition and use of green reaction medium. All synthesized compounds were characterized by <sup>1</sup>H NMR, <sup>13</sup>C NMR, FT-IR spectroscopy, and mass spectrometry.

#### Acknowledgements

The authors are grateful to department of chemistry, B. N. N. College, Bhiwandi and DST-FIST Delhi for providing laboratory and instrumentation facility.

## Supporting Information

Supporting information accompanies this paper on <http://www.acgpubs.org/journal/organic-communications>

ORCID 

Ganesh V. Shitre [0009-0004-4882-7928](https://orcid.org/0009-0004-4882-7928)

Anisha Patel: [0009-0006-6074-9146](https://orcid.org/0009-0006-6074-9146)

Ramesh S. Ghogare: [0000-0003-0810-9636](https://orcid.org/0000-0003-0810-9636)

## References

- [1] Tucker, J.L. Green Chemistry, a Pharmaceutical Perspective. *Org. Process Res. Dev.*, **2006**, *10*(2), 315-319.
- [2] Gupta, P.; Mahajan, A. Green chemistry approaches as sustainable alternatives to conventional strategies in the pharmaceutical industry. *RSC Adv.* **2015**, *5*, 26686-26705.
- [3] Gu, Y. and Je'ro'me, F. Bio-based solvents: an emerging generation of fluids for the design of eco-efficient processes in catalysis and organic chemistry. *Chem. Soc. Rev.* **2013**, *42*, 9550-9570.
- [4] Narsaiah, A.V.; Ghogare, R.S.; Biradar, D.O. Glycerin as alternative solvent for the synthesis of Thiazoles. *Org. Commun.* **2011**, *4*(3) 75-81.
- [5] Ghogare, R.S.; Rajeshwari, K. and Narsaiah, A.V. Glycerol mediated strecker reaction: catalyst-free protocol for the synthesis of  $\alpha$ -amino nitriles. *Lett. Org. Chem.* **2014**, *11*, 688-692.
- [6] Dandia, A.; Jain, A.K. and Laxkar, A.K. Ethyl lactate as a promising bio based green solvent for the synthesis of spiro-oxindole derivatives via 1,3-dipolar cycloaddition reaction. *Tetrahedron Lett.* **2013**, *54*, 3929-3932.
- [7] Yang, J.; Tana, J-N. and Gu, Y. Lactic acid as an invaluable bio-based solvent for organic reactions. *Green Chem.* **2012**, *14*, 3304-3317.
- [8] Hazeri, N.; Maghsoodlou, M.T.; Mir, F.; Kangani, M.; Saravani, H.; Molashahi, E. An efficient one-pot three-component synthesis of tetrahydrobenzo[b]pyran and 3,4-dihydropyrano[c]chromene derivatives using starch solution as catalyst. *Chinese J. Catal.* **2014**, *35*, 391-395.
- [9] Ma, J.; Zhong, L.; Peng, X.; Sun, R.; d-Xylonic acid: a solvent and an effective biocatalyst for a three-component reaction. *Green Chem.* **2016**, *18*, 1738-1750.
- [10] Campos, J. F.; Scherrmannb, M-C.; Berteina-Raboina, S. Eucalyptol: a new solvent for the synthesis of heterocycles containing oxygen, sulfur and nitrogen. *Green Chem.* **2019**, *21*, 1531-39.
- [11] Lim, H.Y.; Dolzhenko, A.V. Gluconic acid aqueous solution: A bio-based catalytic medium for organic synthesis. *Sustain. Chem. Pharm.* **2021**, *21*, 100443.
- [12] Li, B-L.; Li, P-H.; Fang, X-N.; Li, C-X.; Sun, J-L.; Mo, L-P.; Zhang, Z-H. One-pot four-component synthesis of highly substituted pyrroles in gluconic acid aqueous solution. *Tetrahedron* **2013**, *69*(34), 7011-7018.
- [13] Patel, C.; Thakur, A.; Pereira, G. and Sharma, A. Gluconic acid promoted cascade reactions of 2-phenylimidazo[1,2-a] pyridine-3-carbaldehyde with cyclohexane-1,3-dione to create novel fused bisheterocycles. *Syn. Commun.* **2019**, *49*(14), 1836-1846.
- [14] Guo, R-Y.; Wang, P.; Wang, G-D.; Moa, L-P.; Zhang Z-H. One-pot three-component synthesis of functionalized spirooxindoles in gluconic acid aqueous solution *Tetrahedron* **2013**, *69*(8) 2056-2061.
- [15] Xu, Z.; Zhou, P.; Tu, Y.; Liu, D.; Liao, W.; Liao, C. A Green and Highly Efficient Synthesis of 5,5 (Phenylmethylene)bis(1,3-dioxane-4,6-dione) derivatives in biobased gluconic acid aqueous solution. *Heterocycles* **2017**, *94* (6) 1115-1122.

## Green synthesis of 3,4-disubstituted isoxazol-5(4H)-one

- [16] Diwan, F.; Shaikh, M.H.; Fatema, S.; Farooqui, M. Gluconic acid aqueous solution: A bio-compatible media for one pot multicomponent synthesis of dihydropyrano [2,3-c] pyrazoles. *Org. Commun.* **2019**, *12*(4), 188-201.
- [17] Chikkula, K.V.; Raja, S.; Isoxazole-a potent pharmacophore. *Int. J. Pharm. Pharm. Sci.* **2017**, *9*, 13-24.
- [18] Zhu, J.; Mo, J.; Lin, H-p.; Chenb, Y.; Sun, H-p. The recent progress of isoxazole in medicinal chemistry. *Bioorg. Med. Chem.* **2018**, *26*(12), 3065-3075.
- [19] Sysak, A.; Obminska-Mrukowicz, B. Isoxazole ring as a useful scaffold in a search for new therapeutic agents. *Eur. J. Med. Chem.* **2017**, *137*(8), 292-309.
- [20] Kwon, T.; Heiman, A.S.; Oriaku, E.T.; Yoon, K.; Lee, H.J. New steroidal antiinflammatory antedugs: steroidal [16.alpha.,17.alpha.-d]-3'-carbethoxyisoxazolines. *J. Med. Chem.* **1995**, *38*(6), 1048-1051.
- [21] Gordaliza, M. Faircloth, G.T.; Castro, M.A.; Miguel del Corral, J.M., Lo'pez-Va'zquez M.L., Feliciano A.S. Immunosuppressive cyclolignans. *J. Med. Chem.* **1996**, *39*(14), 2865-2868.
- [22] Conti, P.; Dallanoce, C.; Amici, M.D.; Micheli, C.D.; Klotz, K-N. Synthesis of new  $\Delta^2$ -isoxazoline derivatives and their pharmacological characterization as  $\beta$ -adrenergic receptor antagonists. *Bioorg. Med. Chem.* **1998**, *6*(4), 401-408.
- [23] Ishioka, T.; Kubo, A.; Koiso, Y.; Nagasawa, K.; Itai, A.; Hashimotoa, Y. Novel non-steroidal/non-anilide type androgen antagonists with an isoxazolone moiety. *Bioorg. Med. Chem.* **2002**, *10*(5), 1555-1566.
- [24] Lee, Y-S.; Kim, B.H. Heterocyclic nucleoside analogues: design and synthesis of antiviral, modified nucleosides containing isoxazole heterocycles. *Bioorg. Med. Chem. Lett.* **2002**, *12*(10), 1395-1397.
- [25] Deng, B-L.; Cullen, M.D.; Zhou, Z.; Hartman, T. L.; Buckheit, Jr R.W.; Pannecouque, C.; Clercq, E. D.; Fanwickd, P.E.; Cushman, M. Synthesis and anti-HIV activity of new alkenyldiarylmethane (ADAM) non-nucleoside reverse transcriptase inhibitors (NNRTIs) incorporating benzoxazolone and benzisoxazole rings. *Bioorg. Med. Chem.* **2006**, *14*(7), 2366-2374.
- [26] Loh, B.; Vozzolo, L.; Mok, B.J.; Lee, C.C.; Fitzmaurice, R.J.; Caddick, S.; Fassati, A. Inhibition of HIV-1 replication by isoxazolidine and isoxazole sulfonamides. *Chem. Biol. Drug. Des.* **2010**, *75*(5) 461-474.
- [27] Patrick, D.A.; Bakunov, S.A.; Bakunova, S.M.; Suresh Kumar, E.V. K.; Lombardy, R.J.; Jones, S.K.; Bridges, A.S.; Zhirnov, O.; Hall, J.E.; Wenzler, T.; Brun, R.; Tidwell, R.R. Synthesis and in Vitro Antiprotozoal activities of dicationic 3,5-diphenylisoxazoles. *J. Med. Chem.* **2007**, *50*(10), 2468-2485.
- [28] Kozikowski, A.P.; Tapadar, S.; Luchini, D.N.; Kim, K. H.; Billadeau, D.D. Use of the nitrile oxide cycloaddition (noc) reaction for molecular probe generation: a new class of enzyme selective histone deacetylase inhibitors (HDACIs) showing picomolar activity at HDAC6. *J. Med. Chem.* **2008**, *51*(15) 4370-4373.
- [29] Mao, J.; Yuan, H.; Wang, Y.; Wan, B.; Pieroni, M.; Huang, Q.; Breemen, R.B.; Kozikowski, A.P.; Franzblau, S.G. From serendipity to rational antituberculosis drug discovery of mefloquine-isoxazole carboxylic acid esters. *J. Med. Chem.* **2009**, *52*(22) 6966-6978.
- [30] Lavanya, G.; Reddy, L.M.; Padmavathi, V.; Padmaja, A. Synthesis and antimicrobial activity of (1,4-phenylene)bis(arylsulfonylpyrazoles and isoxazoles). *Eur. J. Med. Chem.* **2014**, *73*(12) 187-194.
- [31] Vicentini, C.B.; Romagnoli, C.; Manfredini, S.; Rossi, D.; Mares, D. Pyrazolo[3,4-c]isothiazole and isothiazolo[4,3-d]isoxazole derivatives as antifungal agents. *Pharmaceut. Biol.* **2011**, *49*(5), 545-552.
- [32] Ghorab, M.M.; Bashandy, M.S.; Alsaid, M.S. Novel thiophene derivatives with sulfonamide, isoxazole, benzothiazole, quinoline and anthracene moieties as potential anticancer agents. *Acta Pharm.* **2014**, *64*(4), 419-431.
- [33] Reddy, K.R.; Rao, P.S.; Dev, G.J.; Poornachandra, Y.; Kumar C.G.; Rao P.S.; Narsaiah, B. Synthesis of novel 1,2,3-triazole/isoxazole functionalized 2H-Chromene derivatives and their cytotoxic activity. *Bioorg. Med. Chem. Lett.* **2014**, *24*(7), 1661-1663.
- [34] Padmaja, A.; Rajasekhar, C.; Muralikrishna, A.; Padmavathi, V. Synthesis and antioxidant activity of oxazolyl/thiazolylsulfonylmethyl pyrazoles and isoxazoles. *Eur. J. Med. Chem.* **2011**, *46*(10), 5034-5038.
- [35] Hallenbach, W.; Guth, O.; Seitz, T.; Wroblowsky, H.J.; Desbordes, P.; Wachendorff-Neumann, U.; Dahmen, P.; Voerste, A.; Lo'sel, P.; Malsam, O.; Rama, R.; Hadano, H. 3-Aryl-4-(2,6-Dimethylbenzylidene) Isoxazol-5(4H)-Ones As Fungicides. US Patent, **2012**, Pub. No.: US2012/0065063A1



- [36] Zhang, X.H.; Wang, L.Y.; Zhan, Y.H.; Fu, Y.L.; Zhai, G.H.; Wen, Z.Y. Synthesis and structural studies of 4-[(5-methoxy-1H-indole-3-yl)-methylene]-3-methyl-isoxazole-5-one by X-ray crystallography, NMR spectroscopy, and DFT calculations. *J. Mol. Struct.*, **2011**, 994(1-3), 371-378.
- [37] Zhang, X-H.; Zhan, Y-H.; Chen, D.; Chen, D.; Wang, F.; Wang, L-Y. Merocyanine dyes containing an isoxazolone nucleus: Synthesis, X-ray crystal structures, spectroscopic properties and DFT studies. *Dyes Pigment.* **2012**, 93(1-3), 1408-1415.
- [38] Aret, E.; Meekes, H.; Vlieg, E.; Deroover, G. Polymorphic behavior of a yellow isoxazolone dye. *Dyes Pigment.* **2007**, 72(3), 339-344.
- [39] Biju, S.; Reddy, M.L.P.; Freire, R.O. 3-Phenyl-4-aryl-5-isoxazolonate complexes of Tb<sup>3+</sup> as promising light-conversion molecular devices. *Inorg. Chem. Commun.* **2007**, 10(4), 393-396.
- [40] Kiyani, H.; Darbandi, H.; Mosallanezhad, A.; Ghorbani, F. 2-Hydroxy-5-sulfobenzoic acid: an efficient organocatalyst for the three-component synthesis of 1-amidoalkyl-2-naphthols and 3,4-disubstituted isoxazol-5(4H)-ones. *Res. Chem. Intermed.* **2015**, 41, 7561-7579.
- [41] Khandebharad, A.U.; Sarda, S.R.; Gill, C.H.; Agrawal, B.R. Synthesis of 3-methyl-4-arylmethylene-isoxazol-5(4H)-ones catalyzed by Tartaric acid in aqueous media. *Res. J. Chem. Sci.* **2015**, 5(5), 27-32.
- [42] Maddila, S.N.; Maddila, S.; van Zyl, W.E.; Jonnalagadda, S.B. Ag/SiO<sub>2</sub> as a recyclable catalyst for the facile green synthesis of 3-methyl-4-(phenyl)methylene-isoxazole-5(4H)-ones. *Res. Chem. Intermed.* **2016**, 42, 2553-2566.
- [43] Rikani, A.B.; Setamdideh, D. One-pot and three-component synthesis of isoxazol-5(4h)-one derivatives in the presence of citric acid. *Orient. J. Chem.* **2016**, 32(3), 1433-1437.
- [44] Patil, M.S.; Mudaliar, C.; Chaturbhuj, G.U. Sulfated polyborate catalyzed expeditious and efficient three-component synthesis of 3-methyl-4-(hetero)arylmethylene isoxazole-5(4H)-ones. *Tetrahedron Lett.* **2017**, 58(33), 3256-3261.
- [45] Mosallanezhad, A.; Kiyani, H.; Green synthesis of 3-substituted-4-arylmethylideneisoxazol-5(4h)-one derivatives catalyzed by salicylic acid. *Curr. Organocatal.* **2019**, 6(1), 28-35.
- [46] Basak, P.; Dey, S.; Ghosh, P. Sulfonated graphene-oxide as metal-free efficient carbocatalyst for the synthesis of 3-methyl-4-(hetero)arylmethylene isoxazole-5(4h)-ones and substituted pyrazole. *ChemistrySelect* **2020**, 5(2), 626-636.
- [47] Barkule, A.B.; Gadkari, Y.U.; Telvekar, V.N. One-Pot multicomponent synthesis of 3-methyl-4-(hetero)arylmethylene isoxazole-5(4h)-ones using guanidine hydrochloride as the catalyst under aqueous conditions. *Polycycl. Aromat. Compd.* **2022**, 42(9), 5870-5881.
- [48] Asadi, S.K.; Aleaba, G.; Daneshvar, N.; Shirini, F. Sustainable and green synthesis of 3-methyl-4-arylmethylene-isoxazole-5(4H)-one derivatives under mild conditions using a novel phosphoric acid-based molten salt as catalyst. *Sustain. Chem. Pharm.* **2021**, 21, 100442.
- [49] Deshmukh, S.R.; Nalkar, A.S.; Thopate, S.R. Pyruvic acid-catalyzed one-pot three-component green synthesis of isoxazoles in aqueous medium: a comparable study of conventional heating versus ultrasonication. *J. Chem. Sci.* **2022**, 134, 15.
- [50] Ghogare, R.S.; Patankar-Jain, K.; Momin, S.A.H. A Simple and efficient protocol for the synthesis of 3,4-disubstituted isoxazol-5(4h)-ones catalyzed by succinic acid using water as green reaction medium. *Lett. Org., Chem.* **2021**, 18, 83-87.
- [51] Liu, Q.; Zhang, Y-N. One-pot synthesis of 3-methyl-4-arylmethylene-isoxazol-5(4h)-ones catalyzed by sodium benzoate in aqueous media: a green chemistry strategy. *Bull. Korean Chem. Soc.* **2011**, 32, 3559-3560.
- [52] Liu, Q.; Wu, R-T. Facile synthesis of 3-methyl-4-arylmethylene-isoxazol-5(4H)-ones catalysed by sodium silicate in an aqueous medium. *J. Chem. Res.* **2011**, 35, 598-599.
- [53] Mirzazadeh, M.; Mahdavinia, G.H. Fast and efficient synthesis of 4-Arylidene-3-phenylisoxazol-5-ones. *J. Chem.* **2012**, 9(1), 425-429.
- [54] Liu, Q.; Hou, X. One-pot three-component synthesis of 3-methyl-4-arylmethylene- isoxazol-5(4h)- ones catalyzed by sodium sulfide. *Phosphor. Sulfur Sil. Rel. Element.* **2012**, 187(4), 448-453.
- [55] Kiyani, H.; Ghorbani, F. Synthesis of arylmethylidene-isoxazol-5(4h)-ones in water catalyzed by sodium citrate. *Heteroletters* **2013**, 3(2), 145-153.
- [56] Kiyani, H.; Ghorbani, F. Sodium saccharin as a clean and efficient catalyst for the synthesis of 4-arylidene-3-methylisoxazol-5(4H)-ones via one-pot three-component reaction in aqueous medium. *Heteroletters* **2013**, 3(3), 359-369.
- [57] Kiyani, H.; Ghorbani, F. Efficient tandem synthesis of a variety of pyran-annulated heterocycles, 3,4-disubstituted isoxazol-5(4H)-ones, and  $\alpha,\beta$ -unsaturated nitriles catalyzed by potassium hydrogen phthalate in water. *Res. Chem. Intermed.* **2015**, 41, 7847-7882.

## Green synthesis of 3,4-disubstituted isoxazol-5(4H)-one

- [58] Ablajan, K.; Xiamuxi, H. Efficient one-pot synthesis of  $\beta$ -unsaturated isoxazol-5-ones and pyrazol-5-ones under ultrasonic irradiation. *Synth. Commun.* **2012**, *42*(8), 1128.
- [59] Safari, J.; Ahmadzadeh, M.; Zarnegar, Z. Ultrasound-assisted method for the synthesis of 3-methyl-4-arylmethylene isoxazole-5(4h)-ones catalyzed by imidazole in aqueous media. *Org. Chem. Res.* **2016**, *2*(2), 134-139.
- [60] Khodja, I.A.; Boulcina, R.; Boumoud, T.; Boumoud, B.; Debache, A.  $\text{NaH}_2\text{PO}_4$  catalyzed a three-component 4-arylidene-3-methylisoxazol-5(4H)-ones synthesis in solvent-free conditions. *Der Pharm. Chem.* **2016**, *8*(8), 97-101.
- [61] Dekamin, M.G.; Peyman, S.Z. Phthalimide-N-oxyl salts: efficient organocatalysts for facile synthesis of (Z)-3-methyl-4-(arylmethylene)-isoxazole-5(4H)-one derivatives in water. *Monatsh Chem.* **2016**, *147*, 445-450.
- [62] Kiyani, H.; Ghorban, i F. Potassium phthalimide as efficient basic organocatalyst for the synthesis of 3,4-disubstituted isoxazol-5(4H)-ones in aqueous medium. *J. Saudi Chem. Soc.* **2017**, *21*, S112-S119.
- [63] Laroum, R.; Debache, A. New eco-friendly procedure for the synthesis of 4-arylmethylene-isoxazol-5(4H)-ones catalyzed by pyridinium p-toluenesulfonate (PPTS) in aqueous medium. *Synth. Commun.* **2018**, *48*(14), 1876-1882.
- [64] Laroum, R.; Boulcina, R.; Bensouici, C.; Debache, A. Facile synthesis and antioxidant evaluation of 4-arylmethylideneisoxazol-5(4h)-ones. *Org. Prep. Proced. Int.* **2019**, *51*(6), 583-588.
- [65] Faramarzi, Z.; Kiyani, H. Organocatalyzed three-component synthesis of isoxazol-5(4h)-ones under aqueous conditions. *Heterocycles* **2021**, *102*(9), 1779-1790.
- [66] Daroughezadeh, Z.; Kiyani, H. Efficient and aqueous synthesis of 3,4-disubstituted isoxazol-5(4h)-one derivatives using piperazine under green conditions. *Heterocycles* **2022**, *104*(9), 1625-1640.
- [67] Gharehassanlou, S.; Kiyani, H. A catalytic three-component synthesis of isoxazol-5(4H)-ones under green conditions. *Ind. J. Chem.* **2022**, *61*, 515-520.
- [68] Faramarzi, Z.; Kiyani, H. Steglich's base catalyzed three-component synthesis of isoxazol-5-ones. *Polycycl Aromat. Compd.* **2023**, *43*(4), 3099-3121.
- [69] Kiyani, H.; Ghorbani, F. Expeditious green synthesis of 3,4-disubstituted isoxazole-5(4H)-ones catalyzed by nano-MgO. *Res. Chem. Intermed.* **2016**, *42*, 6831-6844.
- [70] Mashhadinezhad, M.; Shirini, F.; Mamaghani, M. Nanoporous  $\text{Na}^+$ -montmorillonite perchloric acid as an efficient heterogeneous catalyst for synthesis of merocyanine dyes based on isoxazolone and barbituric acid. *Micropor. Mesopor. Material.* **2018**, *262*, 269-282.
- [71] Maleki, B.; Chahkandi, M.; Tayeb, R.; Kahrobaei, S.; Alinezhad, H.; Hemmati, S. Synthesis and characterization of nanocrystalline hydroxyapatite and its catalytic behavior towards synthesis of 3,4-disubstituted isoxazole-5(4H)-ones in water. *Appl Organometal Chem.* **2019**, *33*(10), e5118.
- [72] Ghorbani, F.; Kiyani, H.; Ali Pourmousavi, S. Facile and expedient synthesis of  $\alpha,\beta$ -unsaturated isoxazol-5(4H)-ones under mild conditions. *Res. Chem. Intermed.* **2020**, *46*, 943-959.
- [73] Saadati-Moshtaghin, H.R.; Maleki, B.; Tayeb, R.; Kahrobaei, S.; Abbasinohoji, F. 6-methylguanamine-supported  $\text{CoFe}_2\text{O}_4$ : an efficient catalyst for one-pot three-component synthesis of isoxazol-5(4h)-one derivatives. *Polycycl Aromat. Compd.* **2020**, *42*(3), 885-896.
- [74] Kalhor, M.; Sajjadi, S.M.; Dadras, A. Cu/TCH-pr@SBA-15 nano-composite: a new organometallic catalyst for facile three-component synthesis of 4-arylidene-isoxazolidinones. *RSC Adv.* **2020**, *10*, 27439-27446.
- [75] Kalhor, M.; Samiei, S.; Mirshokraie, S.A.  $\text{MnO}_2$ @Zeolite-Y nanoporous: preparation and application as a high efficient catalyst for multi-component synthesis of 4-arylidene-isoxazolidinones. *Silicon* **2021**, *13*, 201-210.
- [76] Neto, B.A.D.; Oliveira, G.; Machado Ramos, L.; Paiva, R.; Passos, S.; Simões, M.; Machado, F.; Correa, J.R. Synthetic enzyme-catalyzed multicomponent reaction for Isoxazol-5(4H)-one Syntheses, their properties and biological application; why should one study mechanisms? *Org. Biomol. Chem.* **2021**, *19*(7), 1514-1531
- [77] Vekariya, R.H.; Patel, K.D.; Patel, H.D. Fruit juice of Citrus limon as a biodegradable and reusable catalyst for facile, eco-friendly and green synthesis of 3,4-disubstituted isoxazol-5(4H)-ones and dihydropyrano[2,3-c]-pyrazole derivatives. *Res. Chem. Intermed.* **2016**, *42*, 7559-7579.
- [78] Vekariya, R.H.; Patel, H.D. Facile, eco friendly and one pot synthesis of 3,4-disubstituted isoxazol-5(4H)-ones using starch solution as a reaction media. *Ind. J. Chem.* **2017**, *56B*, 890-896.
- [79] Patil, B.M.; Shinde, S.K.; Jagdale, A.A. Fruit extract of avertroha bilimbi: a green neoteric micellar medium for isoxazole and biginelli-like synthesis. *Res. Chem. Intermed.* **2021**, *47*, 4369-4398.

- [80] Gulati, S.; Singh, R.; Sangwan, S. Fruit juice mediated multicomponent reaction for the synthesis of substituted isoxazoles and their in vitro bio-evaluation. Fruit juice mediated multicomponent reaction for the synthesis of substituted isoxazoles and their in vitro bio-evaluation. *Sci Rep.* **2021**, *11*, 23563.
- [81] Atharifar, H.; Keivanloo, A.; Maleki, B. Greener synthesis of 3,4-disubstituted isoxazole-5(4*h*)-ones in a deep eutectic solvent. *Org. Prep. Proced. Int.* **2020**, *52*, 517-523.
- [82] Damghani, F.K.; Kiyani, H.; Pourmousavi, S.A. Green three-component synthesis of merocyanin dyes based on 4-arylideneisoxazol-5(4*H*)-ones. *Curr. Green Chem.* **2020**, *7*, 217-225.
- [83] Setamdideh, D. DOWEX(R)50WX4/H<sub>2</sub>O: A green system for a one-pot and three-component synthesis of isoxazol-5(4*h*)-one derivatives. *J. Mex. Chem. Soc.* **2015**, *59*(3), 191-197.
- [84] Setamdideh, D. One-pot green synthesis of isoxazol-5(4*H*)-one derivatives using Dowex1-x8OH in water. *J. Serb. Chem. Soc.* **2016**, *81*(9), 971-978.
- [85] Ahad, A.; Farooqui, M. Choline chloride and urea based deep eutectic solvent promoted synthesis of arylmethylidene-isoxazol-5(4*H*)-one. *Int. J. Chemtech. Res.* **2017**, *10*(6), 269-273.
- [86] Farahi, S.; Nowrouzi, N.; Irajzadeh, M. three-component synthesis of isoxazolone derivatives in the presence of 4-(*N,N*-dimethylamino)pyridinium acetate as a protic ionic liquid. *Iran J. Sci. Technol. Trans. Sci.* **2018**, *42*, 1881-1887.
- [87] Pourmousavi, S.A.; Fattahi, H.R.; Ghorbani, F.; Kajaani, A.; Ajloo, D. A green and efficient synthesis of isoxazol-5(4*H*)-one derivatives in water and a DFT study. *J. Iran. Chem. Soc.* **2018**, *15*, 455-469.
- [88] Kiyani, H.; Samimi, H.A. Nickel-catalyzed one-pot, three-component synthesis of 3,4-disubstituted isoxazole-5(4*h*)-ones in aqueous medium. *Chiang Mai J. Sci.* **2017**, *44*(3), 1011-1121.
- [89] Hatvate, N.T.; Ghodse, S.M. One-pot three-component synthesis of isoxazole using ZSM-5 as a heterogeneous catalyst. *Synth. Commun.* **2020**, *50*(23), 3676-3683.
- [90] Fozooni, S.; Hosseinzadeh, N.G.; Hamidi, H.; Akhgar, M.R., Nano Fe<sub>2</sub>O<sub>3</sub>, Clinoptilolite and H<sub>3</sub>PW<sub>12</sub>O<sub>40</sub> as efficient catalysts for solvent-free synthesis of 5(4*h*)-isoxazolone under microwave irradiation conditions. *J. Braz. Chem. Soc.* **2013**, *24*(10), 1649-1655.
- [91] Kolsepatil, S.R.; Pahade, N.; Lingampalle, D. An Efficient monosodium glutamate (msg) catalyzed multicomponent synthesis of isoxazolone in water under microwave irradiation. *Der Pharm. Chem.* **2017**, *9*(15), 52-55.
- [92] Kulkarni, P. An efficient solvent-free synthesis of 3,4-disubstituted isoxazole-5(4*H*)-ones using microwave irradiation. *J. Ind. Chem. Soc.* **2021**, *98* (1), 100013.
- [93] Safari, J.; Ahmadzadeh, M.; Zarnegar, Z. Sonochemical synthesis of 3-methyl-4-arylmethylene isoxazole-5(4*H*)-ones by amine-modified montmorillonite nanoclay. *Catal. Commun.* **2016**, *86*, 91-95.
- [94] Ahmadzadeh, M.; Zarnegar, Z.; Safari, J. Sonochemical synthesis of methyl-4-(hetero)arylmethylene isoxazole-5(4*H*)-ones using SnII-montmorillonite. *Green Chem. Lett. Rev.* **2018**, *11*(2), 78-85.
- [95] Saikh, F.; Das, J.; Ghosh, S. Synthesis of 3-methyl-4-arylmethylene isoxazole-5(4*H*)-ones by visible light in aqueous ethanol. *Tetrahedron Lett.* **2013**, *54*(35), 4679-4682.
- [96] Wadavrao, S.B.; Ghogare, R.S.; Narsaiah, A.V., A Simple and efficient protocol for the synthesis of quinoxalines catalyzed by pyridine. *Org. Commun.* **2013**, *6*(1) 23-30.
- [97] Ghogare, R.S. Succinic acid: a novel and efficient organo-catalyst for synthesis of  $\alpha$ -amino nitriles under solvent free condition. *Org. Commun.* **2020**, *13*(3) 103-113.
- [98] Ghogare, R.S.; Narsaiah, A.V., Mandelic acid: an efficient and green organo-catalyst for synthesis of 2,4,5-trisubstituted imidazoles under solvent free condition. *Org. Commun.* **2022**, *15*(1) 44-58.