

An efficient synthesis of quinoxaline derivatives using HCTU as catalyst in DMF

Nitin A. Sasane ^{1,2,*}, Bhushan B. Popatkar ²
and Gangadhar A. Meshram ²

¹Department of Chemistry, Mahatma Phule Arts, Science and Commerce College, Panvel, Dist-Raigad. 410 206 (MS), India

²Department of Chemistry, University of Mumbai, Vidyanagari, Kalina, Santacruz (E), Mumbai 400 098 (MS) India

(Received August 16, 2023; Revised November 29, 2023; Accepted December 02, 2023)

Abstract: We have successfully developed a novel and efficient method for the synthesis of quinoxaline and its derivatives. This method involves the reaction between various 1,2-diamines and substituted phenacyl bromides in the presence of a catalytic amount of *O*-(1H-6-Chlorobenzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HCTU) in dimethylformamide (DMF) as the solvent. This protocol offers several advantages, including mild reaction condition, short reaction time and moderate to high yield of the desired product(s).

Keywords: Quinoxalines; HCTU; 1,2-Diamines; phenacyl bromides. ©2023 ACG Publication. All right reserved.

1. Introduction

Quinoxalines are nitrogen-containing heterocyclic compounds. The functional derivatives of these heterocycles are recognized for their extensive biological applications such as antifungal,¹⁻² anticancer,³⁻⁴ antiviral,⁵ kinase inhibitor,⁶ biocidal⁷ and anti-HIV activity against anti-HIV IIIB strains.⁸ These heterocyclic compounds also find their importance in the other fields such as pharmaceuticals,⁹ sensing of ions,¹⁰⁻¹² dye industries,¹³ agro-chemicals,¹⁴ organic semiconductors¹⁵ and chemically controllable switches¹⁶. Looking to the applications of quinoxaline derivatives, enormous methodologies are developed for their synthesis. The most common method for the synthesis relies on the condensation of an aryl 1,2-diamines with 1,2-dicarbonyl compounds,¹⁷⁻²¹. Similarly, 1,2-keto hydroxyl compounds undergo reaction *via* tandem oxidation procedure involving catalysts such as Pd(OAc)₂,²²⁻²³ MnO₂,²⁴ I₂/DMSO,²⁵⁻²⁶. Quinoxalines and its derivatives also synthesized by oxidative cyclization of α -haloketones and 1,2-diamines from epoxides,²⁷⁻²⁸ and CeCl₃·7H₂O.²⁹ Several well-known catalysts and reagents were used in the synthesis of quinoxalines which include RuCl₂(PPh₃)₃,³⁰ CuSO₄·5H₂O,³¹ PTSA,³² TMSCl,³³ *N*-Bromosuccinimide,³⁴⁻³⁶ T3PDMSO or T3P,³⁷ PEG-400,³⁸ ZnI₂,³⁹ InCl₃,⁴⁰ Ga(OTf)₃,⁴¹ ZrCl₄,⁴² SbCl₃,⁴³ NbCl₅,⁴⁴ and ZnFe₂O₄.⁴⁵ Similarly the synthetic methodology for quinoxalines using recyclable nano-catalysts have been reported in the literature⁴⁶. Many of the established methods for synthesizing quinoxaline and its derivatives have faced certain limitations. These includes the use of costly transition metal-based catalysts, demanding reaction conditions, and

* Corresponding author: E-Mail: sasanenitin79@gmail.com; Phone: +91 8108877501.

HCTU-catalyzed, DMF mediated synthesis of quinoxalines

the incorporation of hazardous or toxic organic solvents to achieve a satisfactory product yield. Hence, the need for advancing methods to synthesize quinoxaline derivatives has become a focal point in current research.

In synthetic organic chemistry, uronium salt-based coupling agents like TBTU, TATU, COMU and HBPYU have gained considerable attention. The catalysts mentioned above have been effectively employed in organic reactions, primarily functioning as dehydrating agents, for instance synthesis of glycopeptide,⁴⁷ for esterification⁴⁸ and for condensation reaction.⁴⁹ Our research group specializes in developing environmental friendly methods for synthesis of heterocyclic compounds.⁵⁰⁻⁵³ We have discovered that uronium salts can serve as an efficient Lewis acid catalyst in heterocyclic compound synthesis.⁵⁴ Therefore, we intend to report the use of uronium salt i.e. *O*-(1*H*-6-Chlorobenzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (figure 1) as a Lewis acid catalyst to achieve the targeted compounds. Recently, catalysts have become increasingly significant, primarily due to their positive impact on both the environment and the economy. They have proven effective in various organic transformations, minimizing the production of undesirable waste materials.




Figure 1. *O*-(1*H*-6-Chlorobenzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate

2. Experimental

2.1. General Methods

All the chemicals, catalyst and reagents were purchased from Merck. Solvents were distilled before use. The progress of reactions was monitored by thin layer chromatography with TLC Silica gel 60 F₂₅₄ purchased from Merck. Column chromatography was performed on silica gel (60–120 mesh). Melting points were recorded by an open glass capillary sealed at one end melting point tube. The IR spectra were recorded on PerkinElmer Frontier FT-IR spectrophotometer. ¹H NMR (300 MHz) and ¹³C NMR (75 MHz) spectra were recorded on Bruker Ultra shield, Avance II model NMR spectrometer. Chemical shifts of ¹H and ¹³C NMR are reported in parts per million (ppm) from tetramethyl silane (TMS) as an internal standard in CDCl₃ /DMSO-*d*₆ as a solvent. Mass spectra were recorded on AB SCIEX QTRAP 3200 model LC-MS spectrophotometer.

2.2. General Procedure

In a 50 mL round-bottom flask, 1,2-diamine (1.0 mmol) and phenacyl bromide (1.0 mmol) were stirred at room temperature for 5 minutes. Then, HCTU (30 mol%) in 5 mL DMF was added to the reaction mixture. The progress of the reaction was monitored using TLC. After the reaction was completed, the product was extracted with ethyl acetate (EtOAc) three times (5 mL each), dried over anhydrous Na₂SO₄, filtered and evaporated the same under reduced pressure. The resulting material was further purified by silica gel column chromatography (60–120 mesh) using an EtOAc-petroleum ether (5:95) mixture to get the pure product.

2.3. Spectral Data for Compounds

*2-Phenylquinoxaline (3a)*³³: yellow solid; mp 75-76⁵⁵ °C; IR (solid, KBr, ν_{\max} , cm⁻¹) 2920, 2850, 2360, 1541, 1444, 1122, 1076, 954, 669, 549; ¹H NMR (500 MHz, CDCl₃) δ 9.33 (s, 1H), 8.26-8.04 (m, 4H), 7.77 (m, 2H), 7.63-7.45 (m, 3H); ¹³C NMR (126 MHz, CDCl₃) δ 151.86, 143.38, 142.32, 141.61, 136.80,

130.28, 130.19, 129.64, 129.54, 129.16, 127.56; LC-MS m/z [M-1]⁻ 205.1; Anal. Calcd for C₁₄H₁₀N₂: C, 81.53; H, 4.89; N, 13.58. Found: C, 81.45; H, 4.81; N, 13.49.

6-Nitro-2-phenylquinoxaline (3b)⁶²: brown solid; mp 165 -168 °C Lit.⁵⁵; IR (solid, KBr, ν_{\max} , cm⁻¹) 2920, 2358, 1521, 1350, 1078, 964, 690; ¹H NMR (500 MHz, CDCl₃) δ 9.50 (s, 1H), 9.03 (d, J = 2.5 Hz, 1H), 8.56 (dd, J = 9.2, 2.5 Hz, 1H), 8.31-8.24 (m, 3H), 7.62 (m, 3H); ¹³C NMR (126 MHz, CDCl₃) δ 154.34, 147.47, 145.51, 144.94, 140.37, 135.65, 131.42, 131.20, 129.44, 127.96, 125.69, 123.80; LC-MS m/z [M-1]⁻ 249.0; Anal. Calcd for C₁₄H₉N₃O₂: C, 66.93; H, 3.61; N, 16.73; O, 12.74. Found: C, 66.85; H, 3.55; N, 16.65; O, 12.69.

6-Methoxy-2-phenylquinoxaline (3c)⁶²: white crystalline solid; mp 73-76 °C Lit.⁵⁶; ¹H NMR (500 MHz, CDCl₃) δ 9.25 (s, 1H), 8.15 (s, 2H), 8.04 (d, J = 9.1 Hz, 1H), 7.54 (m, 4H), 7.12 (m, 1H), 4.00 (s, 3H); ¹³C NMR (126 MHz, CDCl₃) δ 161.2, 152.0, 144.1, 140.8, 137.9, 137.1, 130.1, 129.2, 127.6, 123.0, 107.0, 55.9.

6-Fluoro-2-phenylquinoxaline (3d)⁶³: light yellow solid; mp 96-98 °C, Lit.⁵⁷; IR (solid, KBr, ν_{\max} , cm⁻¹) 3026, 2922, 2852, 1618, 1541, 1161, 1024, 881, 835, 522; ¹H NMR (500 MHz, CDCl₃) δ 9.28 (s, 1H), 8.20-8.16 (m, 2H), 8.10 (d, J = 9.2, 1H), 7.76 (d, J = 9.3, 1H), 7.60-7.48 (m, 4H); ¹³C NMR (126 MHz, CDCl₃) δ 163.0 (d), 152.49, 143.22 (d), 142.65, 138.82 (d), 136.41, 130.50, 129.21, 127.63, 119.97, 113.15, 112.98; LC-MS m/z [M]⁺ 224.9; Anal. Calcd for C₁₄H₉FN₂: C, 74.99; H, 4.05; F, 8.47; N, 12.49. Found: C, 74.89; H, 3.99; F, 8.38; N, 12.40.

6-Chloro-2-phenylquinoxaline (3e)⁶²: light brown crystalline solid; mp 146-147 °C, Lit.⁵⁸; IR (solid, KBr, ν_{\max} , cm⁻¹) 3047, 2920, 2362, 1805, 1600, 1481, 1132, 1072, 958, 904, 588; ¹H NMR (500 MHz, CDCl₃) δ 9.30 (s, 1H), 8.21-8.16 (m, 2H), 8.14 (d, J = 2.3 Hz, 1H), 8.04 (d, J = 8.9 Hz, 1H), 7.67 (dd, J = 8.9, 2.3 Hz, 1H), 7.61-7.50 (m, 3H); ¹³C NMR (126 MHz, CDCl₃) δ 152.52, 143.41, 142.63, 140.09, 136.08, 130.55, 129.21, 128.49, 127.60; LC-MS m/z [M-1]⁻ 238.8; Anal. Calcd for C₁₄H₉ClN₂: C, 69.86; H, 3.77; Cl, 14.73; N, 11.64. Found: C, 69.81; H, 3.69; Cl, 14.68; N, 11.59.

2-(4-methylphenyl)quinoxaline (3f)³³: white solid, mp 97 °C, Lit.⁵⁹; IR (solid, KBr, ν_{\max} , cm⁻¹) 3676, 2970, 2920, 2360, 1615, 1577, 1489, 1124, 1045, 829, 555; ¹H NMR (500 MHz, CDCl₃) δ 9.30 (s, 1H), 8.12 (d, 8.0 Hz, 4H), 7.79-7.68 (m, 2H), 7.36 (d, J = 7.9 Hz, 2H), 2.44 (s, 3H); ¹³C NMR (126 MHz, CDCl₃) δ 151.82, 143.31, 142.33, 141.46, 140.49, 133.99, 130.18, 129.89, 129.54, 129.27, 129.10, 127.42, 21.42; LC-MS m/z [M+1]⁺ 221.3; Anal. Calcd for C₁₅H₁₂N₂: C, 81.79; H, 5.49; N, 12.72. Found: C, 81.71; H, 5.39; N, 12.65.

6-fluoro-2-(4-methylphenyl)quinoxaline (3g)⁶⁴: yellow solid, mp 135 °C, Lit.⁶⁰; IR (solid, KBr, ν_{\max} , cm⁻¹) 2920, 2850, 2360, 1606, 1543, 1425, 1184, 1159, 721; ¹H NMR (500 MHz, CDCl₃) δ 9.27 (s, 1H), 8.10 (m, 2H), 7.75 (d, J = 9.4, 1H), 7.61-7.42 (m, 2H), 7.38 (d, J = 7.9 Hz, 2H), 2.46 (s, 3H); LC-MS m/z [M-1]⁻ 237.3; Anal. Calcd for C₁₅H₁₁FN₂: C, 75.62; H, 4.65; F, 7.97; N, 11.76. Found: C, 75.58; H, 4.59; F, 7.91; N, 11.69.

2-(4-Chlorophenyl)quinoxaline (3h)³³: yellow solid, mp 134-135 °C, Lit.⁶⁰; IR (solid, KBr, ν_{\max} , cm⁻¹) 2918, 2850, 2360, 1591, 1209, 1122, 1089, 715; ¹H NMR (500 MHz, CDCl₃) δ 9.29 (s, 1H), 8.14 (d, 7.8 Hz, 4H), 7.81-7.69 (m, 2H), 7.54 (m, 2H); ¹³C NMR (126 MHz, CDCl₃) δ 150.59, 142.88, 142.23, 141.68, 136.59, 135.20, 130.48, 129.79, 129.61, 129.41, 129.19, 128.78; LC-MS m/z [M+1]⁺ 241.2; Anal. Calcd for C₁₄H₉ClN₂: C, 69.86; H, 3.77; Cl, 14.73; N, 11.64. Found: C, 69.78; H, 3.71; Cl, 14.65; N, 11.57.

6-fluoro-2-(4-chlorophenyl)quinoxaline (3i): light brown; mp 147-151°C; IR (solid, KBr, ν_{\max} , cm⁻¹) 3086, 2970, 2357, 1591, 1433, 1249, 1012, 713, 516; ¹H NMR (500 MHz, CDCl₃) δ 9.26 (s, 1H), 8.30-7.90 (m, 3H), 7.76 (d, J = 9.2, 1H), 7.55 (m, 3H); LC-MS m/z [M+1]⁺ 259.1; Anal. Calcd for C₁₄H₈ClFN₂: C, 65.00; H, 3.12; Cl, 13.70; F, 7.34; N, 10.83; N, 11.64. Found: C, 64.92; H, 3.05; Cl, 13.63; F, 7.29; N, 10.77.

HCTU-catalyzed, DMF mediated synthesis of quinoxalines

*6-methoxy-2-(4-chlorophenyl)quinoxaline (3j)*⁶²: yellow solid, mp 157-160 °C; IR (solid, KBr, ν_{\max} , cm^{-1}) 2848, 2360, 1620, 1462, 1219, 1051, 719; ¹H NMR (400 MHz, CDCl_3): δ 9.19 (s, 1H), 8.09 (d, $J = 8.0$ Hz, 2H), 8.00 (d, $J = 8$ Hz, 1H), 7.50-7.51 (m, 2H), 7.43 (d, $J = 8.0$ Hz, 1H), 7.38 (s, 1H), 3.98 (s, 3H); ¹³C NMR (101 MHz, CDCl_3): δ 160.74, 148.30, 143.20, 142.69, 138.33, 135.99, 135.39, 130.53, 129.30, 128.38, 123.80, 106.51, 55.84; LC-MS m/z $[\text{M}]^+$ 270.7; Anal. Calcd for $\text{C}_{15}\text{H}_{11}\text{ClN}_2\text{O}$: C, 66.55; H, 4.10; Cl, 13.09; N, 10.35; O, 5.91. Found: C, 66.49; H, 4.08; Cl, 13.00; N, 10.28; O, 5.84.

*7-Bromo-2-phenylpyrido[2,3-*b*]pyrazine (3k)*⁶⁰: mp 154.5-155.0 °C, Lit. ⁶¹; IR (solid, KBr, ν_{\max} , cm^{-1}) 3321, 2972, 2883, 2358, 1541, 1377, 1087, 879, 669; ¹H NMR (500 MHz, CDCl_3) δ 7.73 (d, $J = 4.2$ Hz, 1H), 7.54 (d, $J = 8.6$ Hz, 2H), 7.37-7.33 (m, 3H), 7.13 (m, 2H); LC-MS m/z $[\text{M}-1]^-$ 285.5; Anal. Calcd for $\text{C}_{13}\text{H}_8\text{BrN}_3$: C, 54.57; H, 2.82; Br, 27.93; N, 14.69. Found: C, 54.51; H, 2.76; Br, 27.85; N, 14.61.

3. Results and Discussion

As a part of our efforts to develop efficient and environment friendly synthetic method,³⁸ we have explored the synthesis of quinoxaline and its derivatives. This is achieved by reacting different 1,2-diamines and substituted phenacyl bromides in the presence of HCTU, which serves as a cost-effective and eco-friendly Lewis acid catalyst for this organic transformations.


Initially, a model reaction was designed involving *O*-phenylenediamine (1) (1.0 mmol, 0.108 g) and Phenacyl bromide (2) (1.0 mmol, 0.199 g) in 5 mL of ethanol. After stirring at room temperature for five minutes, a catalytic amount of HCTU (*O*-(1H-6-Chlorobenzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate) (30 mol%, 0.123 g) was added to the reaction mixture (Scheme 1). Subsequently, the reaction contents were stirred at room temperature. After 30 minutes, the completion of the reaction was verified using TLC. To optimize the reaction conditions, various solvents and different catalyst loadings were tested on the model reaction. Initially, the impact of various solvents on the reaction conditions was examined and it was observed that as the polarity of the solvent increased, the product yield also increased with a shorter reaction time (Table 1, entries 1, 3, 4 & 5). In the case of water, the product yield remained very low even after 100 minutes of stirring at room temperature, likely due to the limited solubility of the starting materials in water (Table 1, entry 6).

Table 1. Study of an effect of solvents and catalyst on the reaction condition

Entry ^[a]	Solvents	Catalyst (mol %)	Time (min)	Yield ^[b] (%)
1	Ethanol	5	50	52
2	DCM	5	80	38
3	THF	5	48	56
4	DMSO	5	45	58
5	DMF	5	42	61
6	Water	5	100	30
7	DMF	10	39	65
8	DMF	15	35	68
9	DMF	20	35	72
10	DMF	25	33	80
11	DMF	30	30	90
12	DMF	35	30	91
13	DMF	40	28	91


^[a] The reaction was carried out at room temperature

^[b] Isolated yield of the product.



Scheme 1. Model reaction between *O*-phenylenediamine and Phenacyl bromide

After conducting solvent studies it has been observed that DMF was determined to be the most suitable solvent for this reaction (Table 1, entry 11). Similarly, the impact of different percentages of catalyst loading was assessed in the DMF solvent. It was found that as the percentage of the catalyst increased, the chemical yield of the reaction also increased, and the required reaction time decreased (Table 1, entries 7-11). Interestingly, no significant effect on reaction time and overall yield of the product was observed when the catalyst loading exceeded 35 mol% and 40 mol%, respectively (Table 1, entries 12 & 13). Therefore, the optimized reaction condition was established, resulting in a 90% yield of the product (Table 1, entry 11).









Scheme 2. Synthesis of various quinoxaline derivatives

Table 2. Synthesis of quinoxaline derivatives using 1,2-diamines and phenacyl bromides under optimized reaction condition.


Entry ^[3a-j]	1,2-diamines (1)	Phenacyl bromide (2)	Product (3a-j)	Time (min)	Yield ^[b] %
3a	R ¹ = H	R ² = H		45	90
3b	R ¹ = NO ₂	R ² = H		42	93
3c	R ¹ = OMe	R ² = H		45	83
3d	R ¹ = F	R ² = H		42	92

HCTU-catalyzed, DMF mediated synthesis of quinoxalines

3e	R ¹ = Cl	R ² = H		45	91
3f	R ¹ = H	R ² = Me		45	90
3g	R ¹ = F	R ² = Me		42	94
3h	R ¹ = H	R ² = Cl		45	90
3i	R ¹ = F	R ² = Cl		42	94
3j	R ¹ = OMe	R ² = Cl		48	82




^[a] Dissolving 1,2-diamines **1** (1.0 mmol) and various phenacyl bromides **2** (2.0 mmol) in 5 ml of DMF, we achieved the final product with a good yield after 42-45 minutes of continuous stirring. All the products (**3a-j**) were identified using IR, ¹H, and ¹³C NMR.

^[b] Isolated yield of the product.



Scheme 3. Synthesis of 7-bromo-2-phenylpyrido[2,3-b]pyrazine (**3k**)

Table 3. Reaction of 5-bromo 2,3-diaminopyridine and 3,3' diaminobenzidine with phenacyl bromide


Entry ^[3k-1]	1,2, diamine (1)	Phenacyl bromide (2)	Product (3k)	Time (min)	Yield ^[b] %
3k				45	90

^[a] By dissolving 5-Bromopyridine-2,3-diamine (1.0 mmol) and phenacyl bromide in 5 ml of DMF, the desired product was obtained after 45 minutes of continuous stirring. The product (**3k**) was identified through IR, ¹H, and ¹³C NMR.

^[b] Isolated yield of the product.

For generality of this protocol and to examine the substrate scope, various aromatic, heteroaromatic and substituted 1,2-diamines were reacted with substituted phenacyl bromides under the optimized reaction condition (Scheme 2 and 3). The result shows that *O*-phenylenediamine reacts efficiently with substituted phenacyl bromides to afford final product in moderate yield (Table 2, entry 3a, 3f & 3h). When, 1,2-diamine and phenacyl bromide holds electron withdrawing groups the reaction proceeds quickly with an excellent product yield (Table 2, entry 3i) with minimal reaction time. The presence of electron-withdrawing groups attached to the 1,2-diamine molecule seems to accelerate the reaction with phenacyl bromide, leading to a higher yield of the product. (Table 2, entry 3b, 3d, 3e, & 3g). Similarly, when 1,2-diamine possesses electron donating groups gave comparably lesser yield of the desired product (Table 2, entry 3c & 3j). However, 5-bromopyridine-2,3-diamine reacted smoothly with phenacyl bromide under the optimized reaction condition and gives 7-bromo-2-phenylpyrido[2,3-*b*]pyrazine in the moderate yield (Table 3, entry 3k).

The plausible reaction mechanism is illustrated in (Scheme 4). In Step 1, HCTU attacks the active methylene carbon of phenacyl bromide, forming an HCTU bromonium salt. In Step 2, one of the amine groups from *O*-phenylenediamine attacks the active methylene carbon, weakening the HCTU bond. In Step 3, another amine group from *O*-phenylenediamine attacks the carbonyl carbon, leading to the elimination of water and subsequent oxidation, resulting in the formation of quinoxaline as the product.



Scheme 4. Plausible reaction mechanism for the synthesis of quinoxaline

4. Conclusion

To conclude, we have successfully developed a method for synthesizing quinoxaline derivatives utilizing HCTU as a catalyst in DMF solvent under mild and straightforward reaction condition. This method yields the desired product with good to moderate efficiency when various 1,2-diamines and phenacyl bromides are employed. Notable advantages of this methodology include shorter reaction time, ease of catalyst handling and the ability to carry out reactions at room temperature.

Acknowledgements

Authors are thankful to the Head, Department of Chemistry, University of Mumbai and Principal, Mahatma Phule Arts, Science and Commerce College, Panvel for their kind support. The authors are also thankful to laboratory staff of Central Instrumentation Centre of the Department of Chemistry, University of Mumbai for their assistance in spectral analysis.

Supporting Information

Supporting information accompanies this paper on <http://www.acgpubs.org/journal/organic-communications>

HCTU-catalyzed, DMF mediated synthesis of quinoxalines

ORCID Nitin A. Sasane: [0000-0001-7049-6425](https://orcid.org/0000-0001-7049-6425)Bhushan B. Papatkar: [0000-0002-4402-291X](https://orcid.org/0000-0002-4402-291X)Gangadhar A. Meshram: [0000-0001-9490-2539](https://orcid.org/0000-0001-9490-2539)

References

- [1] Seitz, L. E.; Suling, W. J.; Reynolds, R. C. Synthesis and antimycobacterial activity of pyrazine and quinoxaline derivatives. *J. Med. Chem.* **2002**, *45*, 5604–5606.
- [2] Lee, C. H.; Wu, W. C.; Dangate, P. S.; Shen, L. C.; Chung, W. S.; Sun, C. M. Skeletally diverse synthesis of innovative [2,1-c]-1,4-oxazepine and [1,4]-quinoxaline systems. *ACS Comb. Sci.* **2015**, *17*, 623-630.
- [3] Jaso, A.; Zarranz, B.; Aldana, I.; Monge, A. Synthesis of new quinoxaline-2-carboxylate 1,4-dioxide derivatives as anti-mycobacterium tuberculosis agents. *J. Med. Chem.* **2005**, *48*, 2019-2025.
- [4] Lindsley, C. W.; Zhao, Z.; Leister, W. H.; Robinson, R. G.; Barnett, S. F.; Jones, D. D.; Jones, R. E.; Hartman, G. D.; Huff, J. R.; Huber, H. E.; Duggan, M. E. Allosteric Akt (PKB) inhibitors: discovery and SAR of isozyme selective inhibitors. *Bioorganic Med. Chem. Lett.* **2005**, *15*, 761-764.
- [5] Shibinskaya, M. O.; Lyakhov, S. A.; Mazepa, A. V.; Andronati, S. A.; Turov, A. V.; Zhobak, N. M.; Spivak, N. Y. Synthesis, cytotoxicity, antiviral activity and interferon inducing ability of 6-(2-aminoethyl)-6H-indolo [2, 3-b] quinoxalines. *Eur. J. Chem.* **2010**, *45*, 1237-2043.
- [6] Kim, Y. B.; Kim, Y. H.; Park, J. Y.; Kim, S. K. Synthesis and biological activity of new quinoxaline antibiotics of echinomycin analogues. *Bioorganic Med. Chem. Lett.* **2004**, *14*, 541-544.
- [7] Sarges, R.; Howard, H. R.; Browne, R. G.; Lebe, L. A.; Seymour, P. A.; Koe, B. K. 4-Amino[1,2,4]triazolo[4,3-a]quinoxalines: A novel class of potent adenosine receptor antagonists and potential rapid-onset antidepressants. *J. Med. Chem.* **1990**, *33*, 2240-2254.
- [8] Patel, S. B.; Patel, B. D.; Pannecouque, C.; Bhatt, H. G. Design, synthesis and anti-HIV activity of novel quinoxaline derivatives. *Eur. J. Med. Chem.* **2016**, *117*, 230-240.
- [9] Seitz, L. E.; Suling, W. J.; Reynolds, R. C. Synthesis and antimycobacterial activity of pyrazine and quinoxaline derivatives," *J. Med. Chem.* **2002**, *45*, 5604-5606.
- [10] Dey, S. K.; Kobaisi, M. A.; Bhosale, S. V. Functionalized quinoxaline for chromogenic and fluorogenic anion sensing. *ChemistryOpen.* **2018**, *7*, 934-952.
- [11] More, Y. W.; Padghan, S. D.; Bhosale, R. S.; Pawar, R. P.; Puyad, A. L.; Bhosale, S. V.; Bhosale, S. V. Proton triggered colorimetric and fluorescence response of a novel quinoxaline comprising a donor-acceptor system. *Sensors.* **2018**, *18*, 3433.
- [12] Srivani, D.; Gupta, A.; La, D. D.; Bhosale, R. S.; Puyad, A.L.; Xiang, W.; Li, J.; Bhosale, S. V.; Bhosale S. V. Small molecular non-fullerene acceptors based on naphthalenediimide and benzoisoquinoline-dione functionalities for efficient bulk-heterojunction devices. *Dyes Pigm.* **2017**, *143*, 1-9.
- [13] Chang, D. W.; Lee, H. J.; Kim, J. H.; Park, S. Y.; Park, S. M.; Dai, L.; Beak, J. B. Novel quinoxaline-based organic sensitizers for dye-sensitized solar cells. *Org. Lett.* **2011**, *13*, 3880-3883.
- [14] Liu, X. H.; Yu, W.; Min, L. J.; Wedge, D. E.; Tan, C. X.; Weng, J. Q.; Wu, H. K.; Cantrell, C. L.; Hirschel, J. B.; Hua, X. W.; Duke, S. O. Synthesis and pesticidal activities of new quinoxalines. *J. Agric. Food Chem.* **2020**, *68*, 7324-7232.
- [15] Dailey, S.; Feast, W. J.; Peace, R. J.; Sage, I. C.; Till, S.; Wood, E. L. Synthesis and device characterization of side-chain polymer electron transport materials for organic semiconductor applications. *J. Mater. Chem.* **2001**, *11*, 2238-2243.
- [16] Crossley, M. J.; Johnston, L. A. Laterally-extended porphyrin systems incorporating a switchable unit. *Chem comm.* **2002**, *10*, 1122-1123.
- [17] Darabi, H. R.; Mohandessi, S.; Aghapoor, K.; Mohsenzadeh, F. A recyclable and highly effective sulfamic acid/MeOH catalytic system for the synthesis of quinoxalines at room temperature. *Catal. Commun.* **2007**, *8*, 389-392.
- [18] Huang, T. K.; Wang, R.; Shi, L.; Lu, X. X. Montmorillonite K-10: An efficient and reusable catalyst for the synthesis of quinoxaline derivatives in water. *Catal. Commun.* **2018**, *9*, 1143-1147.
- [19] More, S. V.; Sastry, M. N. V.; Yao, C. F. Cerium (IV) ammonium nitrate (CAN) as a catalyst in tap water: A simple, proficient and green approach for the synthesis of quinoxalines. *GreenChem* **2006**, *8*, 91-95.

- [20] Zhao, Z.; Wisnoski, D. D.; Wolkenberg, S. E.; Leister, W. H. Wang, Y.; Lindsley, C. W. General microwave-assisted protocols for the expedient synthesis of quinoxalines and heterocyclic pyrazines. *Tetrahedron Lett.* **2004**, *45*, 4873-4876.
- [21] Brown, D. J. The Chemistry of Heterocyclic Compounds Quinoxalines: Supplement II, John Wiley & Sons: New Jersey, **2004**.
- [22] Robinson, R. S.; Taylor, R. J. K. Quinoxaline synthesis from α -hydroxy ketones via a tandem oxidation process using catalysed aerobic oxidation. *Synlett.* **2005**, *6*, 1003-1005.
- [23] (a) Raw, S. A.; Wilfred, C. D.; Taylor, R. J. K. Preparation of quinoxalines, dihydropyrazines, pyrazines and piperazines using tandem oxidation processes. *Chem. Comm.* **2003**, *18*, 2286-2287.
- [24] Kim, S. Y.; Park, K. H.; Chung, Y. K. Manganese (iv) dioxide-catalyzed synthesis of quinoxalines under microwave irradiation. *Chem Comm.* **2005**, *10*, 1321-1323.
- [25] Bhosale, R. S.; Sarda, S. R.; Ardhapure, S. S.; Jadhav, W. N.; Bhusare, S. R.; Pawar, P. R. An efficient protocol for the synthesis of quinoxaline derivatives at room temperature using molecular iodine as the catalyst. *Tetrahedron Lett.* **2005**, *46*, 7183-1786.
- [26] More, S. V.; Sastry, M.; Wang, C. C.; Yao, C. F.; Molecular iodine: a powerful catalyst for the easy and efficient synthesis of quinoxalines. *Tetrahedron Lett.* **2005**, *46*, 6345-6348.
- [27] Cho, C. S.; Oh, S. G. Copper-catalyzed oxidative cyclization of α -hydroxyketones with o-phenylenediamines leading to quinoxalines. *J. Mol. Catal. A Chem.* **2007** *276*, 205-207.
- [28] Antoniotti, S.; Dunach, E. Direct and catalytic synthesis of quinoxaline derivatives from epoxides and ene-1, 2-diamines. *Tetrahedron Lett.* **2002**, *43*, 3971-3973.
- [29] Narsaiah, A. V.; Kumar, J. K. Glycerin and $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$: A new and efficient recyclable reaction medium for the synthesis of quinoxalines. *Synth. Commun.* **2012**, *42*, 883-892.
- [30] Cho, C. S.; and Oh, S. G. A new ruthenium-catalyzed approach for quinoxalines from o-phenylenediamines and vicinal-diols. *Tetrahedron Lett.* **2006**, *47*, 5633-5636.
- [31] Heravi, M. M.; Taheri, S.; Bakhtiari, K.; Oskooie, H. A. On Water: A practical and efficient synthesis of quinoxaline derivatives catalyzed by $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$. *Synth. Commun.* **2007**, *8*, 211-214.
- [32] Kumbhar, A.; Kamble, S.; Barge, M.; Rashinkar, G.; Salunkhe, R. Bronsted acid hydrotrope combined catalyst for environmentally benign synthesis of quinoxalines and pyrido[2,3-b]pyrazines in aqueous medium. *Tetrahedron Lett.* **2012**, *53*, 2756-2760.
- [33] Wan, J. P.; Gan, S. F.; Wu, J. M.; Pan, Y. Water mediated chemoselective synthesis of 1, 2-disubstituted benzimidazoles using O-phenylenediamine and the extended synthesis of quinoxalines. *Green Chem.* **2009**, *11*, 1633-1637.
- [34] Pardeshi, S. D.; Sathe, P. A.; Vadagaonkar, K. S.; Chaskar, A. C. One-pot protocol for the synthesis of imidazoles and quinoxalines using N-bromosuccinimide. *Adv. Synth. Catal.* **2017**, *359*, 4217-4226.
- [35] Kumar, A.; Dhameliya, T. M.; Sharma, K.; Patel, K. A.; Hirani, R. V.; Bhatt, A. J. Sustainable approaches towards the synthesis of quinoxalines: An update. *J. Mol. Struct.* **2022**, *1259*, 132732.
- [36] Avula, B.; Reddivari, C. K. R.; Muchumarri, R. M. R.; Eraganaboyina, S.; Zyryanov, G. V.; Nemallapudi B. R. Recent advances in the synthesis of quinoxalines. a mini review. *Polycycl. Aromat. Compd.* **2023**, DOI: 10.1080/10406638.2023.2167215
- [37] Harsha, K. B.; Rangappa, K. S. One-step approach for the synthesis of functionalized quinoxalines mediated by T3P@-DMSO or T3P@ via a tandem oxidation-condensation or condensation reaction. *RSC Adv.* **2016**, *6*, 57154-57162.
- [38] Zhang, X. Z.; Wang, J. X.; Sun, Y. J.; Zhan, H. W. Synthesis of quinoxaline derivatives catalyzed by PEG-400. *Chin Chem Lett.* **2010**, *21*, 395-398.
- [39] Sangshetti, J. N.; Kokare, N. D.; Shinde, D. B. Efficient one-pot synthesis of quinoxalines in the presence of zinc iodide as catalyst. *Russ. J. Org. Chem.* **2009**, *45*, 1116-1118.
- [40] Hazarika, P.; Gogoi, P.; Konwar, D. Efficient and green method for the synthesis of 1, 5-benzodiazepine and quinoxaline derivatives in water. *Synth. Commun.* **2007**, *37*, 3447-3454.
- [41] Cai, J. J.; Zou, J. P.; Pan, X. Q.; Zhang, W. Gallium (III) triflate-catalyzed synthesis of quinoxaline derivatives. *Tetrahedron Lett.* **2008**, *49*, 7386-7390.
- [42] Aghapoor, K.; Darabi, H. R.; Mohsenzadeh, F.; Balavar, Y.; Daneshyar, H. Zirconium (IV) chloride as versatile catalyst for the expeditious synthesis of quinoxalines and pyrido[2,3-b]pyrazines under ambient conditions. *Transit. Met. Chem.* **2010**, *35*, 49-53.
- [43] Aghapoor, H. D. K.; Mohsenzadeh, F.; Taala, F.; Asadollahnejad, N.; Badiei, A. Silica-supported antimony (III) chloride as highly effective and reusable heterogeneous catalyst for the synthesis of quinoxalines. *Catal Letters.* **2009**, *133*, 84-89.
- [44] Hou, J. T.; Liu, Y. H.; Zhang, Z. H. NbCl_5 as an efficient catalyst for rapid synthesis of quinoxaline derivatives. *J. Heterocycl. Chem.* **2010**, *47*, 703-706.

HCTU-catalyzed, DMF mediated synthesis of quinoxalines

- [45] Pund, G.B.; Dhupal, S.T.; Hebade, M.J; Farooqui, M.; Dobhal, B. S. Zinc ferrite as reusable and green catalyst for synthesis of quinoxaline derivatives. *J Chem Sci.* **2022**, *134*, 81.
- [46] Keri, R. S.; Reddy, D.; Budagumpi S.; Adimule V. Reusable nano-catalyzed green protocols for the synthesis of quinoxalines: an overview. *RSC Adv.* **2023**, *13*, 20373.
- [47] Habermann, J.; Kunz, H. Solid-phase synthesis of a glycopeptide from the homophilic recognition domain of epithelial cadherin 1 using a *O*-pentafluorophenyluronium salt. *Tetrahedron Lett.* **1998**, *39*, 265-268.
- [48] Twibanire, J.; Grindley, T. B. Efficient and controllably selective preparation of esters using uronium-based coupling agents. *Org. Lett.* **2011**, *13*, 2988-2991.
- [49] Oleg, M.; Youval, S.; Fernando, A.; HCTU and TCTU. New coupling reagents: development and industrial aspects. *Chim. Oggi.* **2002**, *20*, 37-41.
- [50] Sasane, N. A.; Meshram, G. A.; Bhise, K. S.; Popatkar, B. B. Facile, one pot synthesis of indoloquinoxaline and its derivatives using aqueous orange peel extract at room temperature. *IJARST.* **2022**, *2*, 129-133.
- [51] Popatkar, B. B.; Mane, A. A.; Meshram, G. A. Tomato fruit extract: an environmentally benign catalytic medium for the synthesis of isoxazoles derivatives. *Indian J Chem Sect [B].* **2021**, *60*, 1362-1367.
- [52] Popatkar, B. B.; Sasane, N. A.; Meshram, G. A. [BMPTFB]-ionic liquid as an efficient catalyst for the rapid, and eco-friendly synthesis of benzimidazole, 2-substituted benzimidazole, and benzothiazole derivatives at room temperature. *Synth. Commun.* **2022**, *52*, 2249-2259.
- [53] Popatkar, B. B.; Sasane, N. A.; Meshram, G. A. [EMIM]AlCl₄-ionic liquid catalyzed mechanochemically assisted synthesis of 3,4-dihydropyrimidin-2-(1H)-one and thione derivatives. *J. Heterocycl. Chem.* **2023**, *60*, 1199-1209.
- [54] Popatkar, B. B.; Meshram, G. A. HBTU-catalyzed simple and mild protocol for the synthesis of quinoxaline derivatives. *Heterocycles* **2020**, *100*, 1009-1018.
- [55] Jeganathan, M.; Dhakshinamoorthy, A.; Pitchumani, K. One-pot synthesis of 2-substituted quinoxalines using K10-montmorillonite as heterogeneous catalyst. *Tetrahedron Lett.*, **2014**, *55*, 1616-1620.
- [56] Singh, S.; Mishra, P.; Srivastava, M.; Singh, S. B.; Singh, J.; Tiwari, K. P. N-heterocyclic carbene (thiamine) promoted eco-friendly synthesis of quinoxalines under mild reaction conditions. *Green Chem. Lett. Rev.* **2012**, *5*, 587-593.
- [57] Shen, G.; Wang, Z.; Huang, X.; Gong, S.; Zhang, J.; Tang, Z.; Sun, M.; Lv, X. Bonded- and discreted-Lindqvist hexatungstate-based copper hybrids as heterogeneous catalysts for the one-pot synthesis of 2-phenylquinoxalines via 2-haloanilines with vinyl azides or 3-phenyl-2H-azirines. *Dalton Trans.* **2020**, *49*, 13993-13998.
- [58] Rock, J.; Garcia, D.; Espino, O.; Shetu, S. A.; Bacab, M. J. C.; Puc, R. M.; Patel, N. B.; Rivera, G.; Bandyopadhyay, D. Benzopyrazine-based small molecule inhibitors as trypanocidal and leishmanicidal agents: green synthesis, in vitro, and in silico evaluations. *Front. Chem.* **2021**, *9*, 725892.
- [59] Maichrowski, J.; Gjikaj, M.; Hubner, E.; Bergmann, G. B.; Muller, I. B.; Kaufmann, D. E. Efficient synthesis of quinoxaline derivatives by selective modification of 3-chloro-6-fluoroquinoxalin-2(1H)-one 4-oxide. *Eur. J. Org. Chem.* **2013**, *11*, 2091-2105.
- [60] Lv, W.; Xiong, B.; Tan, Z.; Jiang, H.; Zhang, M. Straight forward synthesis of 2-arylquinoxalines from 2-nitroanilines with an equimolar amount of monoalcohols via a base-catalyzed hydrogen-transfer coupling reaction. *Asian J. Org. Chem.* **2015**, *4*, 1127-1131.
- [61] Kekesi, L.; Dancso, A.; Illyes, E.; Boros, S.; Pato, J.; Greff, Z.; Nemeth, G.; Garamvolgyi, R.; Baska, F.; Orfi, L.; Keri, G. Preparation of pyrido[2,3-b]pyrazine ring system via regioselective condensation reaction. *Lett. Org. Chem.* **2014**, *11*, 651-656.
- [62] Chaubey, T. N.; Borpatra, P. J.; Pandey, S. K. Elemental sulfur-mediated synthesis of quinoxalines from sulfoxonium ylides. *Org. Lett.* **2023**, *25*, 5329-5332.
- [63] Ma, J.; Zou, Q.; Yin, G.; Li, F. Base-promoted facile synthesis of quinoxalines from β , β -dibrominated secondary enamides and *o*-phenylenediamines. *Tetrahedron Lett.* **2023**, *126*, 154646.
- [64] Lv, W.; Xiong, B.; Tan Z.; Jiang H.; Zhang M. Straightforward synthesis of 2-arylquinoxalines from 2-nitroanilines with an equimolar amount of monoalcohols via a base-catalyzed hydrogen-transfer coupling reaction. *Asian J. Org. Chem.* **2015**, *4*, 1127-1131.

ACG
publications

© 2023 ACG Publications