

## A Novel Limonoid from the Seeds of *Chisocheton macrophyllus*

Nurlelasari <sup>id</sup> <sup>1\*</sup>, Muhammad Badrul Huda <sup>id</sup> <sup>1</sup>, Wahyu Safriansyah <sup>id</sup> <sup>1</sup>,  
Unang Supratman <sup>id</sup> <sup>1,2</sup>, Yudha P. Budiman <sup>id</sup> <sup>1</sup> Desi Harneti P. Huspa <sup>id</sup> <sup>1</sup>,  
Rani Maharani <sup>id</sup> <sup>1</sup>, Tri Mayanti <sup>id</sup> <sup>1</sup>, Kindi Farabi <sup>id</sup> <sup>1</sup> and Sofa Fajriah <sup>id</sup> <sup>3</sup>

<sup>1</sup>Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Padjadjaran,  
Jl. Raya Bandung-Sumedang Km 21, Jatinangor 45363, West Java, Indonesia

<sup>2</sup>Central Laboratory of Universitas Padjadjaran, Jl. Raya Bandung-Sumedang Km 21, Jatinangor  
45363, West Java, Indonesia

<sup>3</sup>Research Center for Pharmaceutical Ingredients and Traditional Medicine, National Research and  
Innovation Agency (BRIN), Complex Cibinong Science Center – BRIN., Cibinong 16911, Bogor,  
West Java, Indonesia

(Received November 10, 2023; Revised February 26, 2024; Accepted March 06, 2024)

**Abstract:** This study aimed to isolate a novel limonoid compound, (6R,7S, 8R, 9R, 10R, 11R, 13S, 17R)-17-(furan-3-yl)-4, 4, 8, 10, 13-pentamethyl-3-oxo-4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 16, 17-dodecahydro-3H-cyclopenta[a]phenanthrene-6,7,11-triyltriacetate (11 $\alpha$ -acetoxydysobinin) (**1**), and 3 pre-existing limonoids, namely dysobinin (**2**), dysobinol (**3**), and 7-deacetylepoxызadiradione (**4**) from the seeds of *Chisocheton macrophyllus* (Meliaceae). In addition, the structure of the isolated compounds was determined using various spectroscopic techniques, including UV, IR, HRTOFMS, 1D, and 2D NMR. The cytotoxic effects of each compound were then evaluated against breast cancer cells of the Michigan Cancer Foundation-7 (MCF-7), but no significant activity was observed.

**Keywords:** *Chisocheton macrophyllus*; Meliaceae; limonoid; cytotoxic activity; MCF-7. © 2024 ACG Publications. All rights reserved.

### 1. Plant Source

*Chisocheton* is a genus in the family Meliaceae, consisting of 53 species, which are widely distributed in Thailand, India, Malaysia, and Indonesia [1]. This current study aimed to reports the isolation and structural analysis of a novel limonoid (**1**) and 3 pre-existing limonoids (**2-4**) from *C. macrophyllus* seeds. The cytotoxic effects of the isolated compounds on MCF-7 breast cancer cells were also examined.

### 2. Previous Studies

The seeds of *Chisocheton macrophyllus* have been shown to contain bioactive limonoids, such as dysobinin, dysobinol, 7 $\alpha$ -hydroxyneotricilenone, nimonol and other limonoids [2-6].

\* Corresponding author: E-Mail: [nurlelasari@unpad.ac.id](mailto:nurlelasari@unpad.ac.id)


A novel limonoid from *Chisocheton macrophyllus***3. Present Study**

In this study, *Chisocheton macrophyllus* (Miq.) seeds were gathered in June 2022 from the Bogor Botanical Garden, Bogor, Indonesia (-6.601812, 106.799269). The identification of the plant was confirmed by Bogoriense Herbarium staff member, Mr Harto, and a voucher specimen (No. Bo-1295452) was subsequently lodged in the herbarium.

A total of 3.5 kg of *C. macrophyllus* dried seeds were macerated for 6 days in methanol. The seeds were then subjected to extraction, leading to the production of 414 g of crude extract by evaporating the solvent. Subsequently, the crude extract obtained was fractionated with *n*-hexane, ethyl acetate, and *n*-butanol to produce 197g, 41.7g, and 15g of concentrated extract, respectively. The *n*-hexane extract was subjected to vacuum liquid chromatography on silica gel G60, and the eluent system was a gradient of 10% stepwise *n*-hexane-ethyl acetate-methanol. The resulting process produced 9 fractions (A-I), with fraction D (50 mg) being recrystallized using chloroform-methanol to yield fraction **2** (30 mg). Fraction E (2.04 g) underwent chromatography on silica gel CC (70-230 mesh) with elution using a 2.5% gradient of ethyl acetate in *n*-hexane. This process yielded 7 subfractions (E1-E12), and E9 (289.4 mg) underwent chromatography on a silica gel column, followed by recrystallization with chloroform, leading to the production of **1** (14 mg). In addition, fractions E2 to E5 were pooled (142.5 mg) and purified on a silica gel column to produce **3** (16.1 mg). Fractions E9-E12 (96 mg) were combined and subjected to chromatography on a silica gel column, leading to the isolation of **4** (18 mg).

(6*R*, 7*S*, 8*R*, 9*R*, 10*R*, 11*R*, 13*S*, 17*R*)-17-(furan-3-yl)-4, 4, 8, 10, 13-pentamethyl-3-oxo-4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 16, 17-dodecahydro-3*H*-cyclopenta[*a*]phenanthrene-6, 7, 11-triyl triacetate (11  $\alpha$ -acetoxydysobinin) (**1**): Colorless crystals; mp: 203–205°C;  $[\alpha]_{\text{D}}^{26.0} = -1.4^{\circ}$  (c 0.05, CHCl<sub>3</sub>); UV (CH<sub>3</sub>OH):  $\lambda_{\text{max}}$  (log  $\epsilon$ ): 212 (1.42); IR (KBr)  $\nu_{\text{max}}$  2979, 1737, 1666, 1502, 1366, 1377, 1244 cm<sup>-1</sup>; <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz):  $\delta_{\text{H}}$  (ppm) = 0.87 (3H, s, H-18), 1.17 (3H, s, H-28), 1.23 (3H, s, H-19), 1.24 (3H, s, H-29), 1.34 (3H, s, H-30), 2.01 (3H, s, H-1''), 2.03 (3H, s, H-1'), 2.06 (3H, s, H-1'''), 2.47 (1H, m, H-16), 2.56 (1H, d, *J* = 12.4 Hz, H-5), 2.57 (1H, m, H-9), 2.63 (2H, d, *J* = 12.4 Hz, H-12), 2.81 (1H, dd, *J* = 11.0, 7.3 Hz, H-17), 5.40 (1H, d, *J*=7.7 Hz, H-15), 5.43 (1H, d, *J*= 2.5 Hz, H-7), 5.45 (1H; d, *J*=7.7 Hz), 5.46 (1H, m, H-6), 5.86 (1H, d, *J* = 10.3 Hz, H-2), 6.23 (1H, dd, *J*= 1.8, 0.9 Hz, H-22), 7.23 (1H, s, H-21), 7.36 (1H, d, *J*= 10.3 Hz, H-1), 7.43 (d, *J* = 10.3 Hz, H-23); <sup>13</sup>C-NMR (CDCl<sub>3</sub>, 125 MHz):  $\delta_{\text{C}}$  (ppm) = 20.1 (CH<sub>3</sub>-1'''), 20.5 (CH<sub>3</sub>-28), 20.9 (CH<sub>3</sub>-18), 21.1 (CH<sub>3</sub>-1''), 21.4 (CH<sub>3</sub>-1'), 22.0 (CH<sub>3</sub>-29), 28.4 (CH<sub>3</sub>-30), 31.7 (CH<sub>3</sub>-19), 34.5 (CH<sub>2</sub>, C-16), 41.5 (C, C-10), 42.0 (CH, C-9), 42.6 (CH<sub>2</sub>, C-12), 42.8 (C, C-8), 45.3 (CH, C-5), 46.4 (C, C-13), 47.3 (C, C-4), 51.4 (CH, C-17), 69.5 (CH, C-6), 70.6 (CH, C-11), 73.9 (CH, C-7), 111.0 (CH, C-22), 120.0 (CH, C-15), 123.9 (C, C-20), 125.4 (CH<sub>2</sub>, C-2), 139.9 (CH, C-21), 142.9 (CH, C-23), 157.1 (C, C-14), 158.3 (CH, C-1), 170.1 (C, C-2'), 170.1 (C, C-2'') 170.3 (C, C-2'''), 204.1 (C, C-3); HRTOFMS (positive ion mode): *m/z* 553.2816 [M+H]<sup>+</sup> (calcd. for C<sub>32</sub>H<sub>41</sub>O<sub>8</sub>; 553.2801).

Compound **1** appeared as colorless crystals in chloroform, with a molecular formula of C<sub>32</sub>H<sub>40</sub>O<sub>8</sub>, which was determined by HRTOFMS investigation of the positive ion peak at *m/z* 553.2801 [M+H]<sup>+</sup>, necessitating 12 degrees of unsaturation. The IR spectrum revealed absorption spectra, suggesting the presence of aliphatic (2979 cm<sup>-1</sup>), 2 carbonyl for ester and  $\alpha,\beta$ -unsaturated ketone (1737 cm<sup>-1</sup> and 1666 cm<sup>-1</sup>), alkene (1502 cm<sup>-1</sup>), and *gem*-dimethyl groups (1366 and 1377 cm<sup>-1</sup>). The <sup>1</sup>H-NMR spectrum exhibited the presence of 8 tertiary methyls, with 5 in upfield chemical shift ( $\delta_{\text{H}}$  0.87, 1.17, 1.23, 1.24 and 1.34), and 3 in downfield chemical shift at ( $\delta_{\text{H}}$  2.01, 2.03, and 2.06, each 3H). In addition, the results showed the presence of oxymethine protons for H-6, H-7, and H-11 at  $\delta_{\text{H}}$  5.46 (1H, m), 5.43 (1H, d, *J*=2.6 Hz) and 5.45 (1H, m, *J*=7.7 Hz), respectively. Furan skeleton was also observed for H-21, H-22, and H-23 at  $\delta_{\text{H}}$  6.23 (1H, d, *J*=1.45 Hz), 7.23 (1H, s), and 7.36 (1H, d, *J*=1.45 Hz), respectively. A total of 4 olefinic methine group resonating at  $\delta_{\text{H}}$  5.40 (1H, d, *J*=7.7 Hz), 5.86 (1H, d, *J*=10.3 Hz), 6.23 (1H, dd, *J*= 1.8, 0.9 Hz) and 7.36 (1H, d, *J*=10.3 Hz) each for H-15, H-2, H-1 and H-22 were displayed in the <sup>1</sup>H-NMR spectrum.



**Figure 1.** Structures of Compounds 1-4


The  $^{13}\text{C}$ -NMR, DEPT135, and (HSQC) spectra showed the presence of 32 carbons containing carbonyl for ketone and ester group ( $\delta_{\text{C}}$  204.1, 170.1, 170.1, 170.3) as well as 8 methyl tertiary ( $\delta_{\text{C}}$  21.4, 21.1, 20.1, 20.5, 20.9, 22.0, 28.4, and 31.7). The  $^{13}\text{C}$  NMR spectra also displayed 2 methylenes ( $\delta_{\text{C}}$  42.6 and 34.5), along with 3 methines aliphatic ( $\delta_{\text{C}}$  42.0, 45.3, and 51.4). In addition, there were 4 methines olefinic ( $\delta_{\text{C}}$  111.0, 120.0, 125.4, and 158.3), 3 oxymethines ( $\delta_{\text{C}}$  69.5, 70.6, and 73.9), 2 oxygenated  $sp^2$  methine carbons ( $\delta_{\text{C}}$  139.9, and 142.9), and 6 quaternary carbons including  $sp^3$  and  $sp^2$  quaternary carbon ( $\delta_{\text{C}}$  41.5, 42.8, 47.3, 46.4, 123.9, and 157.1). In this study, 7 out of 12 degrees of unsaturation could be attributed to the functionalities, and the pentacyclic limonoid structure accounted for the remaining 5 degrees [3,4].

The NMR data of compound **1** were similar to those reported for dysobinin [5], except for the presence of acetoxy signals [ $\delta_{\text{H}}$  5.45 (1H, d,  $J=7.7$  Hz),  $\delta_{\text{C}}$  70.6], indicating that **1** was an acetoxy analog of dysobinin. In addition, 3 acetoxy groups were located at C-6 ( $\delta_{\text{C}}$  69.5), C-7 ( $\delta_{\text{C}}$  73.9), and C-11 ( $\delta_{\text{C}}$  70.6), respectively elucidated through ( $^1\text{H}$ - $^1\text{H}$  COSY) and (HMBC) experiments (Figure 2). HMBC correlations were observed between H-6 ( $\delta_{\text{H}}$  5.46, m), H-7 ( $\delta_{\text{H}}$  5.43, d,  $J=2.5$  Hz), and H-11 ( $\delta_{\text{H}}$  5.45, d,  $J=7.7$  Hz) and each corresponding acetyl carbons, namely  $\delta_{\text{C}}$  170.1 (C-2'), 170.1 (C-2''), 170.3 (C-2'''). The positions of 3 acetoxy were also revealed by  $^1\text{H}$ - $^1\text{H}$  COSY from H<sub>5</sub>/H<sub>6</sub>/H<sub>7</sub> and H<sub>9</sub>/H<sub>11</sub>/H<sub>12</sub>. The results showed that the position of an  $\alpha,\beta$ -unsaturated ketone was assigned to be in C-1/C-2/C-3 from the correlations of H-1 ( $\delta_{\text{H}}$  7.36, t, 10.3 Hz) to carbonyl ketone  $\delta_{\text{C}}$  204.1 (C-3) with  $^1\text{H}$ - $^1\text{H}$  COSY correlation H<sub>1</sub>/H<sub>2</sub>.


The double bond in ring D was confirmed using HMBC correlations from H-15 ( $\delta_{\text{H}}$  5.40, d,  $J=7.7$  Hz) to similar configuration with dysobinin. The optical rotation of compound **1**, [ $\alpha$ ] $^{26.0}_{\text{D}}$   $-1.4^\circ$  (c 0.05,  $\text{CHCl}_3$ ) was similar to that of dysobinin, as previously reported ([ $\alpha$ ] $^{20.0}_{\text{D}}$   $+150^\circ$ ) [7]. Consequently, the structure of compound **1** had been determined as a novel azadirone type of limonoid derivative and was named *11* $\alpha$ -acetoxydysobinin. C-13, C-16, and C-17 ( $\delta_{\text{C}}$  46.4, 34.5, 51.4) combined with  $^1\text{H}$ - $^1\text{H}$  COSY correlation H<sub>15</sub>/H<sub>16</sub>/H<sub>17</sub>. The furan skeleton was identified by HMBC correlations, from H-22 ( $\delta_{\text{H}}$  6.23, dd, 1.8, 0.9 Hz) correlated to oxygenated methine  $sp^2$  C-23 ( $\delta_{\text{C}}$  142.9) and from H-21 ( $\delta_{\text{H}}$  7.23, s) to C-20, C-22, and C-23 ( $\delta_{\text{C}}$  123.9, 111.0, and 142.9), along with  $^1\text{H}$ - $^1\text{H}$  COSY correlation H<sub>22</sub>/H<sub>23</sub>. The relative configuration of the acetoxy group (C-11) in compound **1** was determined by the NOESY experiment (Figure 3), as well as by comparing the NMR data with

A novel limonoid from *Chisocheton macrophyllus*

dysobinin. The key correlation observed at H-11/CH<sub>3</sub>-19 indicated that the acetoxy group at C-11 was  $\alpha$  oriented and 2 acetoxy at C-6 and C-7 had had been determined as a novel azaridone-type of limonoid derivative and was named *11* $\alpha$ -acetoxydysobinin.



**Figure 2.** Selected HMBC and <sup>1</sup>H-<sup>1</sup>H COSY correlations for **1**



**Figure 3.** Selected <sup>1</sup>H-<sup>1</sup>H NOESY correlations of **1**

Based on the established procedures [4], all compounds were evaluated for their cytotoxicity against the MCF-7 breast cancer cell line, using cisplatin as a positive control. The results showed that compound **1** did not show any significant activity towards MCF-7 breast cancer cells, with IC<sub>50</sub> (inhibitory concentration, 50%) values >250 μg/mL, as shown in Table 1. The findings indicated that the presence of acetoxy at C-11 could reduce cytotoxic activity. This was evidenced by compound **2**, which had better activity compared to compound **1** due to the absence of acetoxy group at C-11.

**Table 1.** Cytotoxicity of compounds **1–4** against MCF-7 breast cancer cell.

Compound	IC <sub>50</sub> (μg/mL)
11 $\alpha$ -acetoxydysobinin ( <b>1</b> )	>250.00
Dysobinin ( <b>2</b> )	68.15
Dysobinol ( <b>3</b> )	243.70
7-deacetylepoxызadiradione ( <b>4</b> )	94.62
Cisplatin (positive control)	38.06

### Acknowledgments

The authors are grateful to Mr. Kansu Haikal and Mrs. Meita Kartika at the Center Laboratory of Universitas Padjadjaran for conducting the HRTOFMS measurement and cytotoxic assay. This study collaboration was conducted by the Research Center of Universitas Padjadjaran and BRIN, 2023.

## Funding Statement

This study was funded by the PDUPT grant, No. 1834/UN6.3.1/PT.00/2023, awarded to Nurlelasari from Universitas Padjadjaran, and the Academic Leadership Grant (No: 1549/UN6.3.1/PT.00/2023), granted to Unang Supratman.

## Supporting Information

Supporting Information accompanies this paper on <http://www.acgpubs.org/journal/records-of-natural-products>

## ORCID

Nurlelasari : [0000-0002-9317-2607](https://orcid.org/0000-0002-9317-2607)

Muhammad Badrul Huda: [0009-0003-0464-5543](https://orcid.org/0009-0003-0464-5543)

Wahyu Safriansyah: [0000-0002-3243-4001](https://orcid.org/0000-0002-3243-4001)

Unang Supratman: [0000-0002-1104-2321](https://orcid.org/0000-0002-1104-2321)

Yudha P. Budiman: [0000-0002-3929-1891](https://orcid.org/0000-0002-3929-1891)

Desi Harneti P. Huspa: [0000-0002-8120-7892](https://orcid.org/0000-0002-8120-7892)

Rani Maharani: [0000-0001-8156-9773](https://orcid.org/0000-0001-8156-9773)

Tri Mayanti: [0000-0003-1628-6288](https://orcid.org/0000-0003-1628-6288)

Kindi Farabi: [0000-0001-5552-3827](https://orcid.org/0000-0001-5552-3827)

Sofa Fajriah: [0000-0002-1758-2102](https://orcid.org/0000-0002-1758-2102)

## References

- [1] J.A. Shilpi, S. Saha, S.L. Chong, L. Nahar, S.D. Sarker and K. Awang (2016). Advances in chemistry and bioactivity of the genus *Chisocheton Blume*, *Chem. Biodivers.* **13**, 483–503.
- [2] I.A. Najmuldeen, A.H.A. Hadi, K. Mohamad, K. Awang, K.A. Ketuly, M.R. Mukhtar, H. Taha, N. Nordin, M. Litaudon, F. Guéritte, A.E. Nugroho and H. Morita (2012). Chisomicines D and E, two new limonoids from *Chisocheton ceramicus*, *Heterocycles* **84**, 1265–1270.
- [3] Wong, C.P., Shimada, M., Nugroho, A.E., Hirasawa, Y., Kaneda, T., Hadi, A.H.A., Osamu, S., & Morita, H. (2012). Ceramicines J-L, new limonoids from *Chisocheton ceramicus*, *J. Nat. Medicin.* **66**, 566–570.
- [4] Nurlelasari, I. Rahmayanti, S. Salam, A. Safari, D. Harneti, R. Maharani, A.T. Hidayat, M. Tanjung, R. Retnowati, Y. Shiono and U. Supratman (2021). A new havanensin-type limonoid from *Chisocheton macrophyllus*, *Appl. Biol. Chem.* **64**. 2-6.
- [5] U. Supratman, S. Salam, W. Naibaho, M. Fajar, Nurlelasari, D.G. Katja, D. Harneti, R. Maharani, A.T. Hidayat, R. Lesmana, M. Azlan Nafiah and Y. Shiono (2020). New cytotoxic limonoids from the stem bark of *Chisocheton pentandrus* (Blanco) Merr, *Phytochemistry Lett.* **35**, 63–67.
- [6] Nurlelasari, D.G. Katja, D. Harneti, M.M. Wardayo, U. Supratman and K. Awang (2017). Limonoids from the seeds of *Chisocheton macrophyllus*, *Chem. Nat. Compound.* **53**, 83–87.
- [7] S. Singh, H. S. Garg, and N. M. Khanna (1976). Dysobinin, A New Tetranortriterpene from *Dysoxylum binectariferu*, *Phytochemistry* **15**, 2001-2002.

**ACG**  
publications

© 2024 ACG Publications