

A New Lignan with Potential Anti-inflammatory Activity from *Notopterygium incisum*

Li-Lian Zhao ¹, Su Hu ¹, Xin-Yu Li ¹, Yun Deng ¹,
Li-Jun Huang ^{*2} and Da-Le Guo ^{*1}

¹ School of Pharmacy, Chengdu University of Traditional Chinese Medicine, 611137, Chengdu, China

² School of Basic Medical Science, Chengdu University of Traditional Chinese Medicine, 611137,
Chengdu, China

(Received May 08, 2025; Revised July 01, 2025; Accepted July 04, 2025)

Abstract: *Notopterygium incisum* is widely used in clinical practice because of its complex chemical composition and remarkable pharmacological effects. In this study, a previously undescribed lignan (**1**) was isolated from *N. incisum*. Its structure was evaluated through examinations of the NMR, HREIMS data and ECD calculation. Compound **1** demonstrated significant anti-inflammatory potential by inhibiting the secretion of nitric oxide (NO) in lipopolysaccharide (LPS)-induced macrophages, with an IC₅₀ value of 3.57 μM.

Keywords: *Notopterygium incisum*; lignan; anti-inflammatory activity. © 2025 ACG Publications. All rights reserved.

1. Plant Source

The roots and rhizomes of *Notopterygium incisum* utilized in this study were sourced from Sichuan Neautus Pharmaceutical Co., Ltd. The authenticity of the plant material was validated by Prof. Guang-Zhi Wang, who is affiliated with Chengdu University of Traditional Chinese Medicine. Moreover, the specimen of *N. incisum* currently preserved in the plant specimen chamber with a collection number of 20221225/CDCM/SCA at the Herbarium of Chengdu University of Traditional Chinese Medicine.

2. Previous Studies

In our continuous pursuit to investigate natural products featuring novel structures and anti-inflammatory properties sourced from traditional Chinese medicines in southwestern China, a chemical analysis of *N. incisum* resulted in the extraction of several previously uncharacterized coumarins [1-3] exhibiting anti-inflammatory effects.

* Corresponding authors: E-Mail: guodale@cdutcm.edu.cn

A new lignan with potential anti-inflammatory activity

3. Present Study

A combined weight of 30 kg of roots and rhizomes from *N. incisum* was transformed into a powder, then extracted with 95% ethanol, resulting in the production of dried extract. Following this, this extract was then dissolved by distilled water and fractionated by a column D101 macroporous resin measuring (80 × 1200 mm) employing a methanol/water gradient. The gradient consisted of varying ratios, specifically starting with a composition of 0% methanol to 100% water, transitioning to 30% methanol and 70% water, followed by 60% methanol and 40% water, and finally reaching a concentration of 95% methanol and 5% water. The fraction eluted with 60% methanol, weighing 101.3 g, underwent silica gel column chromatography. A gradient of petroleum ether and acetone was employed, gradually changing from a 100:0 to a 50:50 volume-to-volume ratio. This process resulted in the collection of eight distinct fractions, labeled Fr.1 to Fr.8. Fr.3, which weighed 18.51 g, underwent further separation using a silica gel column. In this step, petroleum ether and ethyl acetate were used as mobile phases, also adjusted from a 100:0 to a 50:50 volume-to-volume ratio. This further purification resulted in the formation of sixteen sub-fractions (labeled Fr.3.1 to Fr.3.16). Subsequently, Fr.3.6 fractionated using a Sephadex LH-20 column chromatography. The elution process involved a mixture of dichloromethane and methanol in a 50:50 ratio. Five sub-fractions, (labeled Fr.3.6.1 to Fr.3.6.5) were successfully yielded. Fraction 3.6.4 was further purified by semi-preparative HPLC utilizing a C18 column (5 μ m, 10×250mm), and yielded compound **1** (methanol-water: 68:32; 3 mL/min; t_R : 51.0 min; 4.7 mg). Fr 3.6.1 was further fractionated through semi-preparative HPLC with a C18 column (5 μ m, 10×250mm) and resulted in nine sub-fractions, labeled Fr.3.6.1.1 to Fr.3.6.1.9. Fr.3.6.1.3 was subjected to semi-preparative HPLC with the same C18 column, using a methanol-water mixture at a ratio of 68:32 and a flow rate of 3 mL/min and yield 2.3 mg of compound **2**, which eluted at a retention time of 13.0 min. Similarly, Fr.3.6.1.4 was further purified to isolate compound **3** (0.8 mg) at the same methanol-water mixture and flow rate were used, with the compound eluting at a retention time of 18.0 min. Besides, Fr 3.6.2 was refined using semi-preparative HPLC on the same C18 column. This process employed a methanol-water mixture at a ratio of 65:35 and a flow rate of 3 mL/min. This refinement led to the isolation of compound **4** (4.0 mg), which eluted at a retention time of 57.0 min, and compound **5** (2.1 mg), which eluted at a retention time of 59.0 min.




Figure 1. The structures of compounds **1-5**

Compound 1: yellow gum; $[\alpha]_D^{20} = +9.1$ ($c = 0.22$, MeOH), IR (KBr): 3417, 2920, 2848, 1705, 1629, 1595, 1511, 1427, 1384, 1272, 1152, 1032, 700 cm^{-1} ; UV (MeOH) λ_{max} 203 (5.08), 237 (2.03), 288 (1.16), 327 (1.62) nm; CD ($c = 1.74$, MeOH) λ_{max} ($\Delta\epsilon$) 204 (20.31), 220 (-3.05), 240 (0.26) nm; HRESIMS m/z 478.2201 $[\text{M}+\text{NH}_4]^+$ (calcd. 478.2225). For ^1H and ^{13}C NMR data, see Table 1.

Compound **1** was obtained as a yellow gum. Its molecular formula was confirmed to be $\text{C}_{28}\text{H}_{28}\text{O}_6$, by HRESIMS adduct ion peak at m/z 478.2201 $[\text{H}+\text{HN}_4]^+$. The infrared (IR) spectrum revealed prominent absorption bands at 3417 and 1705 cm^{-1} , indicative of hydroxyl and carbonyl

functional groups, respectively. These findings suggest the presence of these functional groups within the compound's structure. The ^1H NMR spectrum provided further insights into the compound's architecture, displaying characteristic signals that are consistent with a polyaromatic framework. Notably, it revealed ten aromatic protons distributed across three distinct phenyl groups observed at δ_{H} 7.32 (H-2'', H-6''), 7.26-7.20 (H-3'', H-5'', H-4''), 6.94 (H-4), 6.91 (H-6), 6.85 (H-5'), 6.72 (H-2'), and 6.70 (H-6'), along with a disubstituted double bond [δ_{H} 6.23 (H-8), 7.56 (H-7)] and a terminal vinyl group displaying resonances at δ_{H} 6.27 (H-8'), 4.96 (H-9'a), and 5.23 (H-9'b). Oxygenated functionalities were identified as two methoxy groups at δ_{H} 3.84 (3-OMe) and 3.91 (4'-OMe), while two methylene groups resonated at δ_{H} 3.01 (H-7'') and 4.41 (H-8''). The ^{13}C NMR analysis of compound **1** displayed a signal at δ_{C} 167.3 (C-9) indicative of a carbonyl group. Signals corresponding to twenty sp^2 -hybridized carbon atoms, consistent with aromatic and olefinic carbons. Five sp^3 hybridized carbon atoms, supporting the presence of methoxy and methylene groups.




Figure 2. The key ^1H - ^1H COSY, HMBC and NOESY correlations of **1**

The analysis results from HSQC and HMBC for compound **1** illustrated the existence of three phenyl groups. Among these, one is identified as a tetrasubstituted phenyl, characterized by aromatic protons at δ_{H} 6.94 (1H, d, $J = 2.1$ Hz, H-4), 6.91 (1H, d, $J = 1.9$ Hz, H-6) correlated with carbons at δ_{C} 146.5 (C-3), 144.3 (C-4), 134.1 (C-1), 121.3 (C-5), 114.3 (C-6), and 111.3 (C-2). The significant coupling constants observed between H-7'' and H-8'' indicate the *E*-configuration of the disubstituted double bond. The HMBC correlations (Figure 2) show the relationships between H-7 and C-1, C-2, C-6, C-9, along with connections between H-8 and C-1 and C-9, developing the initial C6-C3 unit within the lignan framework. A trisubstituted phenyl group exhibiting an ABX coupling system [δ_{H} 6.85 (d, $J = 8.1$ Hz, H-5'), 6.72 (d, $J = 2.1$ Hz, H-2'), 6.70 (dd, $J = 8.1, 2.1$ Hz, H-6'); δ_{C} 146.8 (C-2'), 145.4 (C-3'), 129.6 (C-5'), 126.3 (C-1'), 123.5 (C-6'), 107.4 (C-4')] was linked to a second C6-C3 moiety via ^1H - ^1H COSY correlations (H-7'/H-8'/H-9') and HMBC correlations (H-7'/C-1', C-2', C-6'). The third substituent, a 2-hydroxyethylphenyl group, displayed aromatic signals at δ_{H} 7.32 (2H, t, $J = 7.3$ Hz, H-2'', H-6'') and 7.26–7.20 (3H, m, H-3'', H-5'', H-4''), with aliphatic protons at 3.01 (2H, t, $J = 7.1$ Hz, H-7'') and 4.41 (2H, t, $J = 7.1$ Hz, H-8'') correlated to δ_{C} 138.1 (C-1''), 129.1 (C-3''), 128.7 (C-2''), 126.7 (C-4''), 65.0 (C-8''), 35.9 (C-7''). The methoxy group positions were confirmed through HMBC correlations that connected 4'-OMe to C-4' and 3-OMe to C-3. Additionally, NOESY correlations involving 4'-OMe/H-5' and 3-OMe/H-4 were also considered. Key correlations were observed between the proton at position H-8'' and the carbon at position C-9. Additionally, interactions between the proton at H-7'' and the carbons at positions C-4, C-5, and C-6 as well as the proton at H-7'' and the carbon at position C-1'', C-2'', C-6'' provided further structural insights. Hence, the planar structure of compound **1** was established. The absolute configuration was determined to be 7'S. This conclusion was supported by subsequent ECD analyses [4-5], which is illustrated in Figure 3. As reported in the SciFinder database, compound **1** was identified as a new lignan, exhibiting a similarity of 92% with the most comparable compounds.

A new lignan with potential anti-inflammatory activity




Figure 3: The experimental ECD spectra and calculated curves of **1**

Compounds **2-5** were elucidated to be name Bergaptol[6], name Pabulenol[7], name Notopterol[8], name Notoptol[9] by comparing spectral data in literatures.

In a series of anti-inflammatory assays with dexamethasone (DXM) as positive control, compound **1** demonstrated significant efficacy in reducing inflammatory markers. Specifically, at a concentration of 7.5 μM , compound **1** also significantly downregulated key pro-inflammatory gene expression (IL-1 β and IL-6) in LPS-stimulated RAW 264.7 macrophages. Besides, it markedly decreased nitric oxide (NO) production with a IC_{50} of 3.57 μM . These effects are illustrated in Figure 4, highlighting the potential of compound **1** as a promising anti-inflammatory hit-compound.




Figure 4. Cell viability, RT-PCR test results and IC_{50} determination for compound **1**. A: Cell viability results for compound **1**. B: Expression levels of inflammatory cytokines IL-1 β for compound **1**. C: Expression levels of inflammatory cytokines IL-6 for compound **1**. D: IC_{50} of Compound **1** for NO production in RAW 264.7 macrophages. Data are expressed as the mean $\pm$ SD. ##### $p < 0.0001$, compared with the normal control group. **** $p < 0.0001$, compared with the LPS group.

Table 1. The NMR data of compound **1** in CDCl₃ (δ in ppm, J in Hz)

Compound 1	δ_C	δ_H J (Hz)
1	126.3	-
2	146.8	-
3	145.4	-
4	107.4	6.94 (d, 2.0)
5	129.6	-
6	123.5	6.91 (d, 2.0)
7	145.6	7.56 (d, 15.8)
8	115.4	6.23 (d, 15.8)
9	167.3	-
1'	134.1	-
2'	111.3	6.72 (d, 2.0)
3'	146.5	-
4'	144.3	-
5'	121.3	6.70 (dd, 8.0, 2.0)
6'	114.3	6.85 (d, 8.0)
7'	47.3	5.05 (d, 6.6)
8'	139.7	6.27 (ddd, 17.0, 10.1, 6.6)
9'	116.6	5.23 (dt, 10.1, 1.6), 4.96 (dt, 17.0, 1.6)
1''	138.1	-
2'', 6''	128.7	-
3'', 5''	129.1	7.32 (t, 7.3)
4''	126.7	7.26-7.20 (m)
7''	35.9	7.26-7.20 (m)
8''	64.9	3.01 (t, 7.2)
3-OMe	56.1	4.41 (t, 7.2)
4'-OMe	56.3	3.84 (s)
2-OH	-	3.91 (s)
3'-OH	-	5.99 (s)
		5.50 (s)

Acknowledgments

This work was partly supported by the National Natural Science Foundation of China (No. 82304887) and Sichuan Science and Technology Program (No. 2024ZYD0116).

Supporting Information

Supporting Information accompanies this paper on <https://www.acgpubs.org/journal/records-of-natural-products>

ORCID

Li-Lian Zhao: [0009-0002-0560-9205](https://orcid.org/0009-0002-0560-9205)

Su Hu: [0009-0003-2712-0264](https://orcid.org/0009-0003-2712-0264)

Xin-Yu Li: [0009-0007-5865-5762](https://orcid.org/0009-0007-5865-5762)

Yun Deng: [0000-0002-3428-8992](https://orcid.org/0000-0002-3428-8992)

Li-Jun Huang: [0000-0003-0563-6684](https://orcid.org/0000-0003-0563-6684)

Da-Le Guo: [0000-0003-3219-7066](https://orcid.org/0000-0003-3219-7066)

References

- [1] X. Y. Huo, M. D. Liu, Y. H. Chen, Y. J. Hu, C. Y. Yan, D. L. Guo, and Y. Deng. (2025). Three new coumarins from *Notopterygium incisum* with potential anti-inflammatory activity, *J. Asian Nat. Prod. Res.* **27**, 18–30.
- [2] Y. J. Hu, M. D. Liu, Y. T. Mu, C. C. Li, M. H. Zhao, D. L. Guo, and Y. Deng (2024). Two undescribed coumarins from *Notopterygium Incisum* with anti-inflammatory activity, *Chem. Biodiversity* **21**, e202401093.
- [3] L. S. Jiang, H. X. Li, C. Y. Zhang, Y. T. Mu, M. J. Zhang, Y. J. Hu, and Y. Deng (2025). Bioactive coumarins from *Notopterygium incisum* root: A step toward natural ant-inflammatory candidates, *Fitoterapia* **182**, 106491.
- [4] H. F. Du, L. Li, Y. H. Zhang, X. Wang, C. Y. Zhou, H. J. Zhu, C. U. Pittman, J. W. Shou, and F. Cao. (2025). The first dimeric indole-deterpenoids from a marine-derived *Penicillium sp.* Fungus and their potential for anti-obesity drugs, *Mar. Life. Sci. Technol.* **7**, 120–131.
- [5] Y. Jung, C. Kwon, T. Kim, J. W. Lee, M. K. Shin, and S.H. Shim (2024). Tetramic acid-motif natural products from a marine fungus *Tolypocladium cylindrosporum* FB06 and their anti-Parkinson activities, *Mar. Life Sci. Technol.* **6**, 84–92.
- [6] X. Luo, X. J. Wang, Y. W. Zhao, W. Z. Huang, Z. L. Xu, Z. Z. Wang, and W. Xiao (2016). Chemical constituents from *Notopterygium incisum*, *Zhongcaoyao* **47**, 1492-1495.
- [7] Y. B. Jiang, X. L. Lu, Z. Z. Yang, L. Ma, and Y. Y. Ma (2013). Studies on the chemical constituents of *Angelica dahurica* related to sulfur fumigated, *Zhongguo Shiyang Fangjixue Zazhi* **19**, 74-78.
- [8] X. W. Wu, and X. W. Yang (2020). Coumarins from *Notopterygium incisum* and their inhibitory effect against lipopolysaccharide-induced nitric oxide production in RAW 264.7 macrophage cells, *Zhongcaoyao* **51**, 3383-3392.
- [9] L. M. Li, B. D. Liang, S. W. Yu, and H. D. Sun (2007) Chemical constituents of *Notopterygium incisum*, *Chin. J. Nat. Med.* **5**, 351-354

A C G
publications

© 2025 ACG Publications