

Supporting Information

J. Chem. Metrol. 6:1 (2012) 1-8

Base hydrolytic forced degradation study of zolpidem tartrate by HPLC

**Prathikantam Pushpalatha^{1*}, Rajendra Kumar Sarin¹, Mohd Idris¹,
Mesineni Anand Rao², Tulsidas Ramachandra Rao Baggi³ and
M. Satish Varma⁴**

¹*Central Forensic Science Laboratory, Directorate of Forensic Science, Ministry of Home Affairs,
Hyderabad, India*


²*Director for PGRR Center for Distant Education, Osmania University, Hyderabad, India*

³*Department of Chemistry, University College of Science, Osmania University, Hyderabad, India.*


⁴*Dr. Reddy's Laboratories Ltd. Active pharmaceutical ingredients, IPDO, Bachupally, Hyderabad-
500072, A.P., India.*

Table of Contents	Page
S1: Standard chromatogram of Zolpidem	2
S2: Representative chromatogram showing separation of drug From base hydrolysis degradant.	2
S3: Overlaid UV spectrum's of drug and base hydrolysis degradant.extracts	2
S4: Mass spectra of peak 1(zolpidem tartrate, RT=4.63)	3


* Corresponding author: pushpas_710@yahoo.co.in




S1. Standard chromatogram of Zolpidem



S.2. Representative chromatogram showing separation of drug From base hydrolysis degradant.



S.3 Overlaid UV spectrum's of drug and base hydrolysis degradant.



S.4. Mass spectra of peak 1(zolpidem tartrate, RT=4.63)