

Supporting Information

Rec. Nat. Prod. 10:2 (2016) 168-175

Antinociceptive Effects of Turkish Endemic *Eryngium kotschyi* Boiss. Roots by Bioactivity Guided Fractionation

Sinem Aslan Erdem^{1*}, Okan Arıhan², Anne-Claire Mitaine-Offer³, Alper B. İskit⁴, Murat Kartal⁵ and Marie-Aleth Lacaille-Dubois³

¹*Department of Pharmacognosy, Faculty of Pharmacy, University of Ankara, Tandogan, 06100-Ankara, Türkiye*

²*Department of Physiology, Faculty of Medicine, Hacettepe University, Ankara 06100, Türkiye*

³*Laboratoire de Pharmacognosie, EA 4267(FDE/UFC), Faculté de Pharmacie, Université de Bourgogne, 7, Bd Jeanne d'Arc, BP 87900, 21079 Dijon Cedex, France,*


⁴*Department of Pharmacology, Faculty of Medicine, Hacettepe University, Ankara 06100, Türkiye*

⁵*Department of Pharmacognosy, Faculty of Pharmacy, and Center of Phytotherapy Education, Research and Practice, Bezmialem Vakıf University, 34093 Istanbul, Türkiye*


Table of Contents

Page


S1: Extraction and main fractionation diagram of <i>Eryngium kotschyi</i> roots	2
S2: Isolation diagram of Compound 1 (Pittangretoside C)	3
S3: FAB-Mass Spectrum of Pittangretoside C	4
S4: ¹ H-NMR Spectrum of Pittangretoside C	4
S5: Expansion of NOESY Spectrum of Pittangretoside C	5
S6: Expansion of NOESY Spectrum of Pittangretoside C	6
S7: HMBC Correlations of Pittangretoside C	6
S8: Expansion of HSQC Spectrum of Pittangretoside C	7




S1: Extraction and main fractionation diagram of *Eryngium kotschy* roots




S2: Isolation diagram of Compound 1




S3: FAB-Mass Spectrum of Pittangretoside C




S4: ¹H-NMR Spectrum of Pittangretoside C (600 MHz, Pyridine-*d*₅)




S5: Expansion of NOESY Spectrum of Pittangretoside C (600 MHz, Pyridine-*d*₅)



S6: Expansion of NOESY Spectrum of Pittangretoside C (600 MHz, Pyridine- d_5)



S7: HMBC Correlations of Pittangretoside C



S8: Expansion of HSQC Spectrum of Pittangretoside C