

## Supporting Information

*Rec. Nat. Prod.* 6:2 (2012) 94-100

### Phenolic Compounds and Terpenoids from *Hypericum lanceolatum*

Hippolyte K. Wabo<sup>1</sup>, Théodora K. Kowa<sup>1a</sup>, Antoine Honoré N. Lonfouo<sup>1a</sup>,  
Alembert T. Tchinda<sup>2</sup>, Pierre Tane<sup>1</sup>, Haruhisa Kikuchi<sup>3</sup>, Michel Frédérich<sup>4</sup>  
and Yoshiteru Oshima<sup>3</sup>

<sup>1</sup> Department of Chemistry, University of Dschang, P.O. Box 67, Dschang, Cameroon



<sup>2</sup> Institute of Medical Research and Medicinal Plants Studies (IMPM), P.O. Box 6163, Yaounde, Cameroon

<sup>3</sup> Graduate School of Pharmaceutical Sciences, Tohoku University, 6-3 Aoba, Aramaki, Aoba-ku, Sendai 980-8578, Japan



<sup>4</sup> Laboratory of Pharmacognosy, Department of Pharmacy, CIRM, University of Liege, B36, 4000 Liege, Belgium

Table of contents	Page
S1 : <sup>1</sup> H NMR (Acetone)	2
S2 : <sup>13</sup> C NMR (Acetone)	3
S3 : EIMS	4
S4 : <sup>1</sup> H NMR (DMSO)	5
S5 : APT	6
S6 : EIMS	7


<sup>1</sup>H NMR / Acetone





<sup>13</sup>C NMR/Acetone





EIMS



<sup>1</sup>H NMR/DMSO



APT



EIMS

