Supporting Information

Org. Commun. 10:4 (2017) 273-279

An İntermolecular Diels-Alder cycloaddition under various condition between 1,3-cyclohexadiene and 7-oxabicyclo[2.2.1]hepta-2,5-diene-2,3-dicarboxylate

Özgür Yılmaz and Nermin Şimşek Kuş

Department of Chemistry, Faculty of Arts and Sciences, Mersin University, 33343, Mersin, Türkiye,

| Table of Contents | Page |
|---|------|
| Figure S1. The ¹ H NMR spectrum of compound 6 | 2 |
| Figure S2. The ¹³ C NMR spectrum of compound 6 | 3 |
| Figure S3. The HMQC spectrum of compound 6 | 4 |
| Figure S4. The NOESY spectrum of compound 6 | 5 |
| Figure S5. The ¹ H NMR spectrum of compound 5 | 6 |
| Figure S6. The ¹³ C NMR spectrum of compound 5 | 7 |




Figure S1. ¹H NMR spectrum of compound 6




Figure S2. ¹³C NMR spectrum of compound 6




Figure S3. HMQC NMR spectrum of compound 6




Figure S4. NOESY NMR spectrum of compound 6




Figure S5. ¹H NMR spectrum of compound **5**




Figure S6. 13 C NMR spectrum of compound **5**