

Supporting Information

Rec. Nat. Prod. 17:2 (2023) 300-311

***Clinopodium nepeta* (L.) Kuntze from Bosnia and Herzegovina: Chemical Characterisation of Headspace and Essential Oil of Fresh and Dried Samples**

**Emina Boškailo^{1,4}, Huriya Džudžević-Čančar^{2,4*}, Alema Dedić², Zvonimir
Marijanović³, Amra Alispahić², Ismet Fatih Čančar², Danijela Vidic⁴ and
Igor Jerković⁵**

¹Department of Pharmacy, Faculty of Health Studies, University of Modern Sciences-CKM,
Mostar, Zalik bb, Bosnia and Herzegovina

²Department of Chemistry in Pharmacy, University of Sarajevo-Faculty of Pharmacy, Sarajevo,
Zmaja od Bosne 8 – Campus UNSA, Bosnia and Herzegovina

³Department of Food Technology and Biotechnology, Faculty of Chemistry and Technology,
University of Split, Split, Ruđera Boškovića 35, Croatia

⁴Department of Chemistry, University of Sarajevo-Faculty of Science, Sarajevo,
Zmaja od Bosne 33-35, Bosnia and Herzegovina

⁵Department of Organic Chemistry, Faculty of Chemistry and Technology, University of Split,
Split, Ruđera Boškovića 35, Croatia

Table of Contents	Page
Figure S1: GC-MS chromatogram of <i>C. nepeta</i> - fresh sample HS volatiles on F2- location A	3
Figure S2: GC-MS chromatogram of <i>C. nepeta</i> - dried sample HS volatiles on F1, location A	3
Figure S3: GC-MS chromatogram of <i>C. nepeta</i> - dried sample HS volatiles on F2, location A	4
Figure S4: GC-MS chromatogram of <i>C. nepeta</i> of EO volatiles, location A	4
Figure S5: GC-MS chromatogram of <i>C. nepeta</i> - fresh sample HS volatiles on F1, location B	5
Figure S6: GC-MS chromatogram of <i>C. nepeta</i> - fresh sample HS volatiles on F2, location B	5
Figure S7: GC-MS chromatogram of <i>C. nepeta</i> - dried sample HS volatiles on F1, location B	6
Figure S8: GC-MS chromatogram of <i>C. nepeta</i> - dried sample HS volatiles on F2, location B	6
Figure S9: GC-MS chromatogram of <i>C. nepeta</i> of EO volatiles, location B	7
Figure S10: GC-MS chromatogram of <i>C. nepeta</i> - fresh sample HS volatiles on F1, location C	7
Figure S11: GC-MS chromatogram of <i>C. nepeta</i> - fresh sample HS volatiles on F2, location C	8
Figure S12: GC-MS chromatogram of <i>C. nepeta</i> - dried sample HS volatiles on F1, location C	8

* Corresponding author: E-Mail: huriya.dzudzevic-cancar@ffsa.unsa.ba; Phone: +387-61-871-836, office: +387-33-586-189

Figure S13: GC-MS chromatogram of <i>C. nepeta</i> - dried sample HS volatiles on F2, location C	9
Figure S14: GC-MS chromatogram of <i>C. nepeta</i> - fresh sample HS volatiles on F1, location D	9
Figure S15: GC-MS chromatogram of <i>C. nepeta</i> - fresh sample HS volatiles on F2, location D	10
Figure S16: GC-MS chromatogram of <i>C. nepeta</i> - dried sample HS volatiles on F1, location D	10
Figure S17: GC-MS chromatogram of <i>C. nepeta</i> - dried sample HS volatiles on F1, location D	11
Figure S18: GS-MS chromatogram of <i>C. nepeta</i> of EO volatiles, location D	11

HS- Headspace, EO- essential oil;

Extraction fibers: F1- Divinylbenzene/Carboxen/Polydimethylsiloxane fiber (DVB/CAR/PDMS);

F2- Polydimethylsiloxane/Divinylbenzene fiber (PDMS/DVB);

Locations: A- Pirići ; B- Blagaj; C-Lokve; D- Počitelj




Figure S1: GC-MS chromatogram of *C. nepeta*- fresh sample HS volatiles on F2- location A




Figure S2: GC-MS chromatogram of *C. nepeta*- dried sample HS volatiles on F1, location A




Figure S3: GC-MS chromatogram of *C. nepeta*-dried sample HS volatiles on F2, location A




Figure S4: GC-MS chromatogram of *C. nepeta* of EO volatiles, location A




Figure S5: GC-MS chromatogram of *C. nepeta*- fresh sample HS volatiles on F1, location B




Figure S6: GC-MS chromatogram of *C. nepeta*- fresh sample HS volatiles on F2, location B




Figure S7: GC-MS chromatogram of *C. nepeta*- dried sample HS volatiles on F1, location B




Figure S8: GC-MS chromatogram of *C. nepeta*- dried sample HS volatiles on F2, location B




Figure S9: GC-MS chromatogram of *C. nepeta* of EO volatiles, location B




Figure S10: GC-MS chromatogram of *C. nepeta*- fresh sample HS volatiles on F1, location C




Figure S11: GC-MS chromatogram of *C. nepeta*- fresh sample HS volatiles on F2, location C




Figure S12: GC-MS chromatogram of *C. nepeta*- dried sample HS volatiles on F1, location C




Figure S13: GC-MS chromatogram of *C. nepeta*- dried sample HS volatiles on F2, location C




Figure S14: GC-MS chromatogram of *C. nepeta*- fresh sample HS volatiles on F1, location D




Figure S15: GC-MS chromatogram of *C. nepeta*- fresh sample HS volatiles on F2, location D




Figure S16: GC-MS chromatogram of *C. nepeta*- dried sample HS volatiles on F1, location D




Figure S17: GC-MS chromatogram of *C. nepeta*-dried sample HS volatiles on F1, location D




Figure S18: GC-MS chromatogram of *C. nepeta* EO volatiles, location D