

Supporting Information

Rec. Nat. Prod. 17:3 (2023) 509-515

Chemical Constituents from The Leaves and Twigs of *Magnolia Decidua*

Qi Zhou¹, Jiang Wan¹, Chun-Xiao Jiang^{1,2}, Ying-Peng Tong¹, Zhen-Da Xie¹, Ting Wang¹, Dao-An Xiao³, Zhi-Yong Xiao⁴ and Jin-Feng Hu^{1,2*}

¹ Institute of Natural Medicine and Health Products, School of Pharmaceutical Sciences, Zhejiang Provincial Key Laboratory of Plant Evolutionary Ecology and Conservation, Taizhou University, Taizhou, Zhejiang 318000, P.R. China

² Department of Natural Medicine, School of Pharmacy, Fudan University, Shanghai 201203, P.R. China

³ College of Chemistry and Bioengineer, Yichun University, Yichun, Jiangxi 336000, PR China

⁴ Yichun Academy of Sciences, Yichun, Jiangxi 336000, PR China

Table of Contents	Page
Figure S1: HR-ESI-MS spectrum of 1 (Manglycoside A)	2
Figure S2: ¹ H-NMR (400 MHz, DMSO) spectrum of 1 (Manglycoside A)	2
Figure S3: ¹³ C-NMR (100 MHz, DMSO) spectrum of 1 (Manglycoside A)	3
Figure S4: HSQC spectrum of 1 (Manglycoside A)	3
Figure S5: ¹ H- ¹ H COSY spectrum of 1 (Manglycoside A)	4
Figure S6: HMBC spectrum of 1 (Manglycoside A)	4
Figure S7: HMBC spectrum of 1 (Manglycoside A) (From δ H1.0 ppm to δ H 6.0 ppm)	5
Figure S8: HMBC spectrum of 1 (Manglycoside A) (From δ H7.0 ppm to δ H 7.5 ppm)	5
Figure S9: The Scifinder searching results of compound 1	6
Table S1: The structural comparison of similar compounds with 1	6
Table S2: The NMR data comparison of similar compounds with 1	7
The spectra data of compounds 2-17	8




Figure S1: HR-ESI-MS spectrum of **1** (Manglycoside A)




Figure S2: ¹H-NMR (400 MHz, DMSO) spectrum of **1** (Manglycoside A)




Figure S3: ^{13}C -NMR (100 MHz, DMSO) spectrum of **1** (Manglycoside A)




Figure S4: HSQC spectrum of **1** (Manglycoside A)




Figure S5: ^1H - ^1H COSY spectrum of **1** (Manglycoside A)




Figure S6: HMBC spectrum of **1** (Manglycoside A)




Figure S7: HMBC spectrum of **1** (Manglycoside A) (From δ_H 1.0 ppm to δ_H 6.0 ppm)




Figure S8: HMBC spectrum of **1** (Manglycoside A) (From δ_H 7.0 ppm to δ_H 7.5 ppm)







Score: 96 1. 1131878-62-0  Absolute stereochemistry. C₁₈ H₂₂ O₉ Benzoic acid, 4-((β-D-glucopyranosyloxy)-3-methoxy-, ethyl ester ~ Key Physical Properties Molecular Weight 358.34 Boiling Point (Predicted) Value: 574.1±50.0 °C Condition: Press: 760 Torr Density (Predicted) Value: 1.405±0.06 g/cm3 Condition: Temp: 20 °C Press: 760 Torr pKa (Predicted) Value: 12.77±0.70 Condition: Most Acidic Temp: 25 °C	Score: 95 2. 130774-07-1  Absolute stereochemistry. C₁₈ H₂₂ O₉ Benzoic acid, 4-[(6-deoxy-α-L-mannopyranosyloxy)-3,5-dimethoxy-, methyl ester ~ Key Physical Properties Molecular Weight 358.34 Boiling Point (Predicted) Value: 493.2±45.0 °C Condition: Press: 760 Torr Density (Predicted) Value: 1.358±0.06 g/cm3 Condition: Temp: 20 °C Press: 760 Torr pKa (Predicted) Value: 12.84±0.70 Condition: Most Acidic Temp: 25 °C Spectra	Score: 94 3. 163623-36-7  Absolute stereochemistry. C₁₈ H₂₀ O₉ Benzoic acid, 4-((β-D-glucopyranosyloxy)-3-hydroxy-, ethyl ester ~ Key Physical Properties Molecular Weight 344.31 Boiling Point (Predicted) Value: 621.6±55.0 °C Condition: Press: 760 Torr Density (Predicted) Value: 1.501±0.06 g/cm3 Condition: Temp: 20 °C Press: 760 Torr pKa (Predicted) Value: 8.49±0.35 Condition: Most Acidic Temp: 25 °C
Score: 94 4. 294847-45-3  Absolute stereochemistry. C₁₈ H₂₂ O₁₀ Benzoic acid, 3-((β-D-glucopyranosyloxy)-4,5-dimethoxy-, methyl ester ~ Key Physical Properties Molecular Weight 374.34 Boiling Point (Predicted) Value: 544.1±50.0 °C Condition: Press: 760 Torr Density (Predicted) Value: 1.426±0.06 g/cm3 Condition: Temp: 20 °C Press: 760 Torr pKa (Predicted) Value: 12.75±0.70 Condition: Most Acidic Temp: 25 °C	Score: 94 5. 502844-31-7  Absolute stereochemistry. C₁₈ H₂₀ O₁₀ Benzoic acid, 3-((β-D-glucopyranosyloxy)-4,5-dihydroxy-, ethyl ester ~ Key Physical Properties Molecular Weight 360.31 Boiling Point (Predicted) Value: 673.3±55.0 °C Condition: Press: 760 Torr Density (Predicted) Value: 1.581±0.06 g/cm3 Condition: Temp: 20 °C Press: 760 Torr pKa (Predicted) Value: 7.46±0.48 Condition: Most Acidic Temp: 25 °C	Score: 94 6. 586398-39-2  Rotation (-), Absolute stereochemistry. C₁₈ H₂₂ O₁₀ Benzoic acid, 4-((β-D-glucopyranosyloxy)-3,5-dimethoxy-, methyl ester ~ Key Physical Properties Molecular Weight 374.34 Melting Point (Experimental) Value: 91-93 °C Condition: Solv: methanol (67-56-1) Boiling Point (Predicted) Value: 544.1±50.0 °C Condition: Press: 760 Torr Density (Predicted) Value: 1.426±0.06 g/cm3 Condition: Temp: 20 °C Press: 760 Torr pKa (Predicted) Value: 12.71±0.70 Condition: Most Acidic Temp: 25 °C

Figure S9: The Scifinder searching results of compound **1**

Table S1: The structural comparison of similar compounds with **1**







No.	Similarity score	Chemical structure and CAS Chemical structure of number	Chemical structure of compound 1
1	96	 CAS NO. 1131878-62-0	
2	95	 CAS NO. 130774-07-1	
3	94	 CAS NO. 163623-36-7	

Table S2: The NMR data comparison of similar compounds with **1**

NO.	Compound 1 ^a	ethyl 4-β-D-glucopyranosyloxy-3-methoxybenzoate ^b (CAS NO. 1131878-62-0)	methyl syringate α-L-rhamnoside ^c (CAS NO. 130774-07-1)		
	δ _H	δ _C	δ _H	δ _C	
1		125.93	NMR data is not available		127.9
2, 6	7.24 (2H, s)	106.76	7.31 (2H, s)		108.5
3, 5		153.39			155.3
4		138.68			140.8
7		165.78			168.9
1'	4.31 (2H, q, <i>J</i> = 7.2 Hz)	61.34	3.89 (3H, s)		53.6
2'	1.32 (3H, t, <i>J</i> = 7.2 Hz)	14.70			
1''	5.19 (1H, s)	102.57	5.34 (1H, d, <i>J</i> = 2Hz)		104.2
2''	3.90 (1H, br s)	70.93	4.14 (1H, dd, <i>J</i> =4, 2Hz)		73.0
3''	3.65 (1H, dd, <i>J</i> = 3.2, 9.6 Hz)	70.68	3.90 (1H, dd, <i>J</i> =10, 4Hz)		72.8
4''	3.24 (1H, t, <i>J</i> = 9.6 Hz)	72.02	3.44 (1H, t, <i>J</i> = 10Hz)		74.5
5''	3.99 (1H, dd, <i>J</i> = 6.4, 9.6 Hz)	70.59	4.24 (1H, dq, <i>J</i> = 10, 7Hz)		72.1
6''	1.06 (3H, d, <i>J</i> = 6.4 Hz)	18.20	1.21 (3H, d, <i>J</i> =7Hz)		18.7
3, 5-OCH₃	3.80 (6H, s)	56.61	3.87 (6H, s)		57.4

^a in DMSO-*d*₆.^b H. T. Simonsen, M. S. Nielsen, N. J. Christensen, U. Christensen, T. V. La Cour, M. S. Motawia, B. P. A. Jespersen, S. B. Engelsens, B. L. Moller (2009). Molecular Interactions between Barley and Oat β-Glucans and Phenolic Derivatives, *J. Agric. Food Chem.* **57**, 2056-2064.^c R. C. Cambie, A. R. Lal, C. E. F. Rickard, and N. Tanaka (1990). Chemistry of Fijian plants. V.1) Constituents of *Fagraea gracilipes* A. Gray. *Chem Pharm Bull.* **38**, 1857-1861.

The spectra data of compounds 2-17

(-)-Syringaresinol (2) : colorless solid, $C_{22}H_{26}O_8$. 1H NMR (400 MHz, DMSO- d_6) δ : 3.05 (2H, m, H-8, 8'), 3.75 (12H, s, 3, 3', 5, 5'-OMe), 3.79 (2H, m, H-9a, 9'a), 4.16 (2H, dd, J = 8.8, 6.8 Hz, H-9b, 9'b), 4.61 (2H, d, J = 3.6 Hz, H-7, 7'), 6.60 (4H, s, H-2, 2', 6, 6'), 8.28 (2H, s, 4-OH', 4''-OH); ^{13}C -NMR (100 MHz, DMSO- d_6) δ : 53.7 (C-8, 8'), 56.0 (3, 3', 5, 5'-OMe), 71.1 (C-9, 9'), 85.3 (C-7, 7'), 103.6 (C-2', 2'', 6', 6''), 131.4 (C-1, 1'), 134.8 (C-4, 4'), 147.9 (C-3, 3', 5, 5').

(+)-Pinoresinol (3) : White amorphous powder, $C_{20}H_{22}O_6$. 1H NMR (400 MHz, DMSO- d_6) δ : 3.02 (2H, m, H-1, 5), 3.74 (2H, dd, J = 8.0, 3.6 Hz, H-4a, 8a), 3.78 (6H, s, 3, 3'-OMe), 4.14 (2H, dd, J = 8.0, 7.2 Hz, H-4b, 8b), 4.60 (2H, d, J = 4 Hz, H-2, 6), 6.70 (2H, d, J = 8.0 Hz, H-5', 5''), 6.74 (2H, dd, J = 8.0, 1.2 Hz, H-6', 6''), 7.00 (2H, d, J = 1.2 Hz, H-2', 2''); ^{13}C -NMR (100 MHz, DMSO- d_6) δ : 54.1 (C-1, 5), 56.1 (OMe $\times$ 2), 71.4 (C-4, 8), 85.6 (C-2, 6), 110.9 (C-2', 2''), 115.6 (C-5', 5''), 119.1 (C-6', 6''), 132.7 (C-1', 1''), 146.4 (C-4', 4''), 148.0 (C-3', 3'').

Pharsyringaresinol (4) : White amorphous powder, $C_{30}H_{38}O_{14}$, 1H NMR (400 MHz, DMSO- d_6) δ : 4.78 (1H, d, J = 4 Hz, H-2), 3.14 (1H, m, H-1), 4.30 (1H, dd, J = 16.0, 10.0 Hz, H-4a), 3.90 (1H, dd, J = 10.0, 4.0 Hz, H-4b), 3.16 (1H, m, H-5), 4.74 (1H, d, J = 4 Hz, H-6), 4.30 (1H, dd, J = 16.0, 10.0 Hz, H-8a), 3.90 (1H, dd, J = 10.0, 4.0 Hz, H-8b), 6.60 (1H, s, H-2', 6'), 6.66 (1H, s, H-2'', 6''), 3.85 (1H, s, 3'/5'-OCH₃), 3.86 (1H, s, 3''/5''-OCH₃), 4.81 (1H, m, H-1'), 3.36 (1H, m, H-2''), 3.42 (1H, m, H-3''), 3.35 (1H, m, H-4''), 3.50 (1H, m, H-5''), 4.28, (1H, m, H-6a'') 4.24 (1H, m, H-6b''), 1.94 (3H, s, H-OAc). ^{13}C -NMR (100 MHz, DMSO- d_6) δ : 131.8 (C-1), 104.1 (C-2, 6), 148.4 (C-3, 5), 135.3 (C-4), 85.8 (C-7), 54.2 (C-8), 71.7 (C-9), 170.5 (C-10), 21.0 (C-11), 133.8, (C-1'), 104.3 (C-2', 6'), 153.3 (C-3', 5'), 138.0 (C-4'), 85.5 (C-7'), 54.1 (C-8'), 71.6 (C-9'), 103.1 (C-1''), 74.1 (C-2''), 74.5 (C-3''), 76.7 (C-4''), 79.6 (C-5''), 63.9 (C-6'').

Syringaresinol- β -D-glucoside (5) : White amorphous powder, 1H -NMR (DMSO- d_6 , 400 MHz) δ : 6.67 (2H, s, H-2'', H-6''), 6.61 (2H, s, H-2', H-6'), 4.89 (1H, d, J = 4.4 Hz, H-gluc-1), 4.68 (1H, d, J = 4.4 Hz, H-2), 4.62 (1H, d, J = 4.2 Hz, H-6), 4.18 (2H, t, J = 6.4 Hz, H-8a, H-4a), 3.81 (1H, br. d, J = 3.6 Hz, H-4b), 3.77 (6H, s, 2 $\times$ OCH₃), 3.76 (6H, s, 2 $\times$ OCH₃), 3.20-3.22 (2H, m, H-1, H-5). ^{13}C NMR (100 MHz, DMSO- d_6) δ 153.1 (C-3'', C-5''), 148.4 (C-3', C-5'), 137.7 (C-4'), 135.3 (C-4''), 134.2 (C-1''), 131.8 (C-1'), 104.6 (C-2''), 104.6 (C-6''), 104.1 (C-2', C-6'), 103.2 (C-gluc-1), 85.8 (C-2), 85.5 (C-6),

77.7 (C-glc-5), 77.0 (C-glc-3), 74.6 (C-glc-2), 71.7 (C-4, C-8), 70.4 (C-glc-4), 61.4 (C-glc-6), 56.9 (3'',5''-OMe), 56.5 (3',5'-OMe), 54.1 (C-5), 54.0 (C-1).

Honokiol (6) : $C_{18}H_{18}O_2$, 1H -NMR (400 MHz, $CDCl_3$) δ : 3.27, 3.36 (each 2H, br d, $J = 7$ Hz, H-7, H-7'), 5.15 (4H; m, H-9, H-9'), 5.99 (2H, m, H-8, H-8'), 6.90 (2H, d, $J = 8$ Hz), 7.03 (2H, dd, $J = 8, 2$ Hz), 7.22 (2H, d, $J = 2$ Hz). ^{13}C -NMR (100 MHz, $CDCl_3$) δ : 132.2 (C-1), 130.2 (C-2), 126.5 (C-3), 150.8 (C-4), 115.5 (C-5), 131.1 (C-6), 39.4 (C-7), 137.8 (C-8), 115.6 (C-9), 127.8 (C-1'), 128.5 (C-2'), 129.5 (C-3'), 128.8 (C-4'), 116.5 (C-5'), 154.0 (C-6'), 35.0 (C-7'), 136.1 (C-8'), 116.8 (C-9'),

Magnolol diacetate (7): colorless powder, $C_{22}H_{22}O_4$. 1H NMR (400 MHz, DMSO) δ : 6.96 (2H, d, $J = 2.4$ Hz, H-6, 6'), 6.80 (2H, dd, $J = 2.4, 8$ Hz, H-4, 4'), 6.52 (2H, d, $J = 8$ Hz, H-3, 3'), 4.22 (2H, t, $J = 6.4$ Hz, H-8, 8'), 3.24 (2H, d, $J = 6.4$ Hz, H-9, 9'), 1.74 (6H, s, 2-OAc, 2'-OAc), 1.63 (2H, m, H-7a,7'a), 1.35 (2H, m, H-7b,7'b). ^{13}C -NMR (100 MHz, DMSO) δ : 130.3 (C-1, 1'), 129.2 (C-2, 2'), 118.6 (C-3, 3'), 127.7 (C-4, 4'), 132.0 (C-5, 5'), 129.1 (C-6, 6'), 30.5 (C-7, 7'), 65.5 (C-8, 8'), 39.6 (C-9, 9'), 174.6 (C=O), 24.1 (C-OAc)

Rel-(3R,3'S,4R,4'S)-3,3',4,4'-tetrahydro-6,6'-dimethoxy[3,3'-bi-2H-benzopyran]-4,4'-diol (8) : colorless oil, $C_{20}H_{22}O_6$. 1H -NMR ($CDCl_3$, 400 MHz) δ : 6.89 (2H, d, $J = 2.0$ Hz, H-5, H-5'), 6.88 (1H, d, $J = 7.8$ Hz, H-8, H-8'), 6.82 (2H, dd, $J = 1.8, 8.2$ Hz, H-7, H-7'), 4.73 (2H, d, $J = 4.0$ Hz, H-4, H-4'), 4.24 (2H, dd, $J = 6.8, 9.2$ Hz, H-2a, H-2a'), 3.90 (6H, s, OCH_3), 3.87 (2H, dd, $J = 3.8, 9.2$ Hz, H-2e, H-2e'), 3.10 (2H, m, H-3, H-3'). ^{13}C NMR (100 MHz, $CDCl_3$) δ 146.7 (C-6, C-6'), 145.2 (C-9, C-9'), 132.9 (C-10, C-10'), 119.0 (C-7, C-7'), 114.3 (C-5, C-5'), 108.6 (C-8, C-8'), 85.9 (C-4, C-4'), 71.7 (C-2, C-2'), 56.0 (OCH_3), 54.2 (C-3, C-3').

Rhemaneolignan A (9) : colorless amorphous powder, $C_{21}H_{24}O_8$, 1H NMR (400 MHz, $DMSO-d_6$) δ : 7.60 (1H, d, $J = 16$ Hz, H-7), 7.33 (1H, d, $J = 2$ Hz, H-3), 7.19 (1H, dd, $J = 2.0, 8.0$ Hz, H-5), 7.06 (1H, d, $J = 8$ Hz, H-6), 6.97 (1H, d, $J = 2$ Hz, H-3'), 6.75 (1H, dd, $J = 2.0, 8.0$ Hz, H-5'), 6.68 (1H, d, $J = 8$ Hz, H-6'), 6.53 (1H, d, $J = 16$ Hz, H-8), 4.71 (1H, t, $J = 4$ Hz, H-7'), 3.82 (1H, s, H- OCH_3), 3.73 (1H, s, H- OCH_3), 3.71 (1H, s, H- OCH_3), 4.40 (1H, m, H-8'), 3.56 (1H, m, H-9'a), 3.23 (2H, m, H-9'b). ^{13}C NMR (100 MHz, $DMSO-d_6$) δ : 150.1 (C-1), 151.3 (C-2), 111.5 (C-3), 127.2 (C-4), 123.1 (C-5), 115.0 (C-6), 145.2 (C-7), 115.2 (C-8), 167.5 (C-9), 145.99 (C-1'), 147.51 (C-2'), 111.7 (C-3'), 133.3 (C-4'), 119.5 (C-5'), 115.6 (C-6'), 71.5 (C-7'), 84.3 (C-8'), 60.7 (C-9'), 51.7 ($-OCH_3$), 56.0 ($-OCH_3$ '), 56.2 ($-OCH_3''$).

Melianoninol (10) : yellow oil, $C_{20}H_{20}O_6$, 1H -NMR (400 MHz, $CDCl_3$) δ : 7.60 (1H, s, H-6'), 7.04 (1H, s, H-3'), 6.90 (1H, d, $J = 2.4$ Hz, H-2), 6.88 (1H, dd, $J = 7.2, 2.4$ Hz, H-6), 6.87 (1H, d, $J = 7.2$ Hz, H-5), 4.97 (1H, d, $J = 7.2$ Hz, H-7'), 4.25 (1H, t, $J = 7.2, 3.6, 4.0$ Hz, H-8'), 3.70 (2H, m, $J = 3.6, 4.0$ Hz,

H-9'), 3.90 (6H, m, 7, H-13'). ^{13}C -NMR (100 MHz, CDCl_3) δ : 51.7 (C-8'), 56.0 (C-13'), 61.0 (C-7), 73.2 (C-9'), 86.2 (C-7'), 108.9 (C-5), 110.9 (C-6), 114.3 (C-2), 119.2 (C-6'), 120.0 (C-3'), 122.3 (C-11'), 127.8 (C-3), 129.8 (C-4'), 132.0 (C-1'), 145.3 (C-2'), 146.7 (C-1), 149.1 (C-4), 151.2 (C-5'), 152.2 (C-10'), 193.6 (C-12').

9-Acetoxy syringin (11) : white amorphous powder, $\text{C}_{14}\text{H}_{16}\text{O}_8$, ^1H -NMR (400 MHz, CDCl_3) δ : 6.78 (2H, s, H-2, 6), 6.58 (1H, d, J = 16 Hz, H-7), 6.33 (1H, dt, J = 6.4, 12.4, 16 Hz, H-8), 4.67 (1H, d, J = 9.6 Hz, H-9), 2.06 (3H, s, H-11), 4.93 (1H, d, J = 7.2 Hz, H-1'), 3.20 (1H, m, H-2'), 3.03 (1H, m, H-3'), 3.12 (1H, m, H-4'), 3.20 (1H, m, H-5'), 3.60 (1H, m, H-6'a), 3.39 (1H, m, H-6'b), 3.78 (6H, s, H-3, 5- OCH_3); ^{13}C -NMR (100 MHz, CDCl_3) δ : 132.04 (C-1), 105.29 (C-2,6), 153.19 (C-3,5), 134.83 (C-4), 133.55 (C-7), 123.64 (C-8), 64.78 (C-9), 170.63 (C-10), 21.23 (C-11), 102.89 (C-1'), 74.64 (C-2'), 77.72 (C-3'), 70.39 (C-4'), 77.03 (C-5'), 61.34 (C-6'), 56.84 (3, 5- OCH_3)

4-Allylcatechol (12) : brown powder, $\text{C}_9\text{H}_{10}\text{O}_2$, ^1H NMR (400 MHz, CDCl_3) δ : 5.74 (1H, d, J = 2 Hz, H-3), 5.68 (1H, dd, J = 8, 2 Hz, H-5), 5.55 (1H, d, J = 8 Hz, H-6), 4.95 (1H, m, H-8), 4.03 (2H, dd, J = 16, 8 Hz, H-9), 2.27 (2H, m, H-7). ^{13}C NMR (100 MHz, CDCl_3) δ : 144.23 (C-1), 142.51 (C-2), 137.63 (C-3), 114.71 (C-4), 119.38 (C-5), 114.20 (C-6), 39.03 (C-7), 115.16 (C-8), 131.46 (C-9).

p-Hydroxybenzaldehyde (13) : white powder, $\text{C}_7\text{H}_6\text{O}_2$, ^1H -NMR ($\text{DMSO}-d_6$, 400 MHz) δ : 9.78 (1H, s, H-CHO), 7.75 (2H, d, J = 8.4 Hz, H-2, H-6), 6.92 (2H, d, J = 8.5 Hz, H-3, H-5). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 191.4 (1-HO), 163.8 (C-4), 132.6 (C-1), 128.9 (C-2, C-6), 116.3 (C-3, C-5).

Liriodenine (14) : yellow amorphous powder, $\text{C}_{17}\text{H}_9\text{NO}_3$, ^1H -NMR ($\text{DMSO}-d_6$, 400 MHz) δ : 8.82 (1H, d, J = 7.6 Hz, H-5), 8.64 (1H, d, J = 7.2 Hz, H-11), 8.37 (1H, d, J = 7.2 Hz, H-8), 8.05 (1H, d, J = 6 Hz, H-4), 7.90 (1H, t, J = 8 Hz, H-9), 7.66 (1H, t, J = 8 Hz, H-10), 7.58 (1H, s, H-3), 6.52 (2H, s, $-\text{O}-\text{CH}_2-\text{O}-$).

N-Acetyldehydroanonaine (15) : $\text{C}_{18}\text{H}_{13}\text{NO}_3$, ^1H NMR (400 MHz, CDCl_3) δ : 8.97 (1H, m, H-11), 7.71 (1H, m, H-8), 7.55 (2H, m, H-9, 10), 7.23 (1H, s, H-7), 7.18 (1H, br s, H-7), 7.05 (1H, s, H-3), 6.16 (2H, s, OCH_2O), 4.10 (2H, t, J = 7.2 Hz, H-5), 3.07 (2H, t, J = 7.2 Hz, H-4), 2.28 (3H, s, $-\text{CH}_3$). ^{13}C -NMR (100 MHz, CDCl_3) δ : 142.2 (C-1), 145.6, (C-2), 109.0 (C-3), 126.1 (C-3a), 30.7 (C-4), 41.8 (C-5), (C-6a), 116.9 (C-7), 131.3 (C-7a), 127.8 (C-8), 126.8 (C-9), 127.0 (C-10), 127.2 (C-11), 127.4 (C-11a), 118.2 (C-11b), 120.0 (C-11c), 101.3 ($-\text{OCH}_2\text{O}-$), 169.9 ($\text{NC}=\text{O}$), 23.1 ($-\text{CH}_3$).

N-trans-feruloyltyramine (16) : $\text{C}_{18}\text{H}_{19}\text{NO}_4$, Colorless crystal, ^1H NMR (400 MHz, DMSO) δ 7.98 (1H, t, J = 5.5 Hz, NH), 7.32 (1H, d, J = 16 Hz, H-7), 7.11 (1H, d, J = 2 Hz, H-2), 7.01 (2H, d, J = 8 Hz, H-2', 6'), 6.99 (2H, dd, J = 2, 8 Hz, H-6), 6.79 (1H, d, J = 8 Hz, H-5), 6.68 (2H, d, J = 8.4 Hz, H-3', 5'), 6.44 (1H, d, J = 16 Hz, H-8), 3.80 (3H, s, OCH_3), 3.34 (2H, q, J = 6.7 Hz, H-8'), 2.65 (2H, t, J = 7.2 Hz, H-7'). ^{13}C NMR (100 MHz, DMSO) δ 129.9 (C-1), 111.2 (C-2), 148.3 (C-3), 148.7 (C-4),

115.6 (C-5), 122.0 (C-6), 139.3 (C-7), 119.5 (C-8), 165.8 (C-9), 56.0 (C-OCH₃), 126.9 (C-1'), 130.0 (C-2', 6'), 116.1 (C-3', 5'), 156.1 (C-4'), 34.9 (C-7'), 41.1 (C-8'),

N-trans-feruloyl-3-methoxytyrmine (17) : Colorless crystal, C₁₉H₂₁NO₅, ¹H-NMR (CDCl₃, 400 MHz) δ: 7.50 (1H, d, *J* = 16 Hz, H-7'), 7.31 (1H, d, *J* = 1.6 Hz, H-2'), 7.05 (1H, d, *J* = 1.6 Hz, H-2), 7.01 (1H, dd, *J* = 8.0, 1.6 Hz, H-6'), 6.85 (1H, d, *J* = 8.0 Hz, H-5'), 6.70 (1H, d, *J* = 8.0 Hz, H-5), 6.68 (1H, dd, *J* = 8.0, 1.6 Hz, H-6), 6.25 (1H, d, *J* = 15.6 Hz, H-8'), 3.59 (2H, m, H-8), 2.80 (2H, *J* = 8.0 Hz, H-7). ¹³C NMR (100 MHz, CDCl₃) δ 166.46 (C = O), 147.77 (C-4'), 147.30 (C-3'), 147.15 (C-3), 144.43 (C-4), 140.84 (C-7'), 130.67 (C-1), 127.15 (C-1'), 122.04 (C-6), 121.25 (C-6'), 118.19 (C-5), 115.06 (C-2), 114.72 (C-8'), 111.60 (C-2'), 109.96 (C-5'), 56.38 (OMe), 55.90 (OMe), 41.00 (C-8), 35.26 (C-7).