

Supporting Information

J. Chem. Metrol. 17:1 (2023) 15-24

Simultaneous estimation of remogliflozin etabonate and vildagliptin in a tablet formulation: UV spectrophotometric and HPLC-PDA method

Prachi Joshi ¹ and Rajendra Kotadiya ¹

¹ Department of Pharmaceutical Quality Assurance, Ramanbhai Patel College of Pharmacy, Charotar University of Science and Technology, Changa, Dist-Anand, Gujarat, India, Pin. 388421

Table of Contents	Page
Figure S1: Chemical structure of RE (a) and VLD (b)	2
Figure S2: Overlain spectra of RE and VLD	3
Figure S3: UV spectra for specificity studies: Spectra of blank (a), placebo (b), standard (c) and sample (d) mixture of RE and VLD	7
Figure S4: Chromatograms showing specificity investigation: a blank sample (a), a placebo sample (b), a standard sample (c), and a mixture of RE and VLD in the sample (d)	8
Figure S5: Peak purity index studies	8




Figure S1: Chemical structure of RE (a) and VLD (b)




Figure S2: Overlain spectra of RE and VLD




Figure S3: UV spectra for specificity studies: Spectra of blank (a), placebo (b), standard (c) and sample (d) mixture of RE and VLD




Figure S4: Chromatograms showing specificity investigation: a blank sample (a), a placebo sample (b), a standard sample (c), and a mixture of RE and VLD in the sample (d)




Figure S5: Peak purity index studies