Supporting Information

Rec. Nat. Prod. 17:5 (2023) 947-951

Callignan A, A New Neolignan from the Leaves of

Callicarpa kwangtungensis

Hancheng Ren and Weiqiang Fei*

Hangzhou Vocational and Technical College, Hangzhou 310018, China

Table of Contents	Page
Figure S1: H NMR Spectrum of compound 1 (400 MHz)	2
Figure S2: ¹³ C NMR Spectrum of compound 1 (100 MHz)	2
Figure S3: HSQC Spectrum of compound 1	3
Figure S4: HMBC Spectrum of compound 1	3
Figure S5: ¹ H- ¹ H COSY Spectrum of compound 1	4
Figure S6: NOESY Spectrum of compound 1	4
Figure S7: HRESIMS Spectrum of compound 1	5
Figure S8: ¹ H NMR Spectrum of 2 (400 MHz)	5
Figure S9: ¹ H NMR Spectrum of 3 (400 MHz)	6
Figure S10: Scifinder similarity report for compound 1	7




Figure S1: ¹H NMR spectrum of compound 1

4.5

4.0

15

158

3.5

3.0

2.5

2.0 ppm

5.5

5.0

2.06

6.0

1.17

7.0

3.05

6.5




Figure S2: ¹³C NMR spectrum of compound 1




Figure S3: HSQC spectrum of compound 1




Figure S4: COSY spectrum of compound 1




Figure S5: HMBC spectrum of compound 1




Figure S6: NOESY spectrum of compound 1

© 2023 ACG Publications. All rights reserved.




Figure S7: HRESIMS spectrum of compound 1






Figure S8: ¹H NMR spectrum of compound 2






Figure S9: 1 H NMR spectrum of compound 3




Figure S10: Scifinder similarity report for compound 1